

The Biology and Control of Localized Dry Spots on Sand Greens

By ROBERT A. HUDSON, Ph.D. and KARL DANNEBERGER, Ph.D.

Department of Agronomy
The Ohio State University

Hydrophobic (non-wettable) soils occurring on bentgrass sand greens constructed to USGA specifications have been previously described and partially characterized. These areas, which resist wetting, have been termed localized dry spots (LDS). The LDS syndrome starts with the turf turning a blue-green color followed by a loss of turgor and finally shoot die-back.

The symptoms observed are usually in irregularly shaped patches of variable size. Frog-eye type patches, characteristic of some turfgrass patch diseases, have been observed but are not predominant. Symptoms are most severe in hot, dry weather. Lower temperatures and adequate water will result in regrowth of the shoot system of plants that survive.

Management practices for the control of LDS are inconsistent at best, yet the following practices have aided in reducing symptom severity. Topdressing with sand that contains a minimal amount of "fine" particles. As will be discussed later, small particles (especially in the silt-clay size) may tend to aggravate the problem over time. Repeated core cultivation, especially in the spring and fall, has helped reduce the severity of LDS. Wetting agents, which reduce the surface tension of water, have given some degree of control for LDS, but are best used in a preventative program. Syringing of the greens may

be used as a stop-gap measure, but primarily serve to lower the canopy temperature and rarely will alleviate symptoms. Frequently, various combinations of the above strategies are necessary, and a trial and error type approach is needed to achieve adequate control of LDS.

Previous studies have shown an organic coating is present on sand grains associated with LDS, and removal of the coating yielded substances with an Infrared (IR) spectra characteristic of fulvic acids. Fulvic acids are a diverse group of large molecules, common in most soils, that are extractable in solutions with a high pH and do not precipitate when the pH is lowered to approximately 1 or below. Previous studies did not include an extraction of wettable soil from bentgrass sand greens, and therefore it could not be determined if the fulvic acid associated with LDS was unique compared to fulvic acids in the wettable areas.

Studies were conducted at the Ohio State University from 1989 through 1991 to provide a more complete characterization of the organic matter and soil characteristics associated with LDS, and included samples from wettable areas for comparative purposes.

(Continued on Page 12)

TURF SUPPLY COMPANY

2797 Eagandale Blvd.

Eagan, MN 55121

"PYTHIUM"

Prevention is better than Desiccation.

 **NOR-AM**
NOR-AM CHEMICAL COMPANY

BANOL

PREVENTATIVE and
CURATIVE Products

Available.

 **RHONE-POULENC**
ALIETTE

CIBA-GEIGY[®]

**SUBDUE
PACE**

TURF SUPPLY COMPANY

*The most complete source of Pythium Management
Products for your Course.*

(612) 454-3106 --- 1-800-551-4857

Biology and Control—

(Continued from Page 10)

First, several common classes of soil organic matter were extracted from two different sites, using several extraction sequences, were quantified and analyzed structurally using several techniques. Structural analyses of lipids (compounds that are similar to oils) were accomplished by gas chromatography/mass spectroscopy (GC/MS) and the large molecules that were extracted in alkaline solution, i.e. fulvic and humic acids, were analyzed by IR and nuclear magnetic resonance (NMR) spectroscopy.

Second, particle size distributions were determined and the extent of non-wettability determined for each size range. While particle size distributions have been determined in previous studies, there were no reports of which fractions were the most hydrophobic, if any. Also, the area in the soil profile that displayed the most hydrophobicity was determined using soil columns collected from greens with LDS, allowed to dry down, re-wetted from the bottom, and the distance that was infiltrated recorded at one and two minutes. Finally, since bentgrass roots have been reported to be colonized by various fungi, both pathogenic and non-pathogenic, roots associated from wettable and non-wettable areas were stained and examined for the extent of fungal colonization present.

Results obtained from the organic matter extraction and analysis indicated that LDS soils had greater amounts of all organic matter fractions studied than soils that were wettable. The only structural difference observed was from LDS that occurred on greens that were three years old, and this was only detected following an initial extraction with methanol. It appears that there is either a unique structure, or interaction be-

tween several structures, occurring in the LDS sample. One possible scenario to explain these results is that a unique structure or structures act to "prime" the LDS areas, and then the syndrome is intensified by subsequent drying cycles, which after several years may mask the unique component that initiated the LDS. The origin of the organic compounds could not be determined in these studies, but it is probably derived from bentgrass roots, soil microflora, or both.

Particle size distribution analysis showed no significant differences between the wettable and non-wettable soils. Hydrophobicity, as determined by how long it took a water drop to penetrate the sample, indicated that particles less than 0.25 mm in size were the most hydrophobic. Since the greens are constructed with 85-90% sand, this size fraction has been largely ignored in previous studies on LDS, but since this is the most chemically reactive fraction, due to the presence of clays, it would not be surprising that this is where organic-inorganic interactions would be the most prevalent. The hydrophobicity was the greatest in the area immediately below the thatch-soil interface. This is the area in the soil profile with the most biological activity, especially in regards to root colonization and thatch degradation. Electron micrographs of soil particles that were approximately 0.1 mm in diameter showed that the particles in LDS samples had an extensive organic coating compared to particles from wettable soils.

Roots from both areas on the sand greens were heavily colonized by several fungi including vesicular-arbuscular mycorrhizae (VAM), *Phialophora* spp., *Pythium* spp., and *Polyomyxa graminis*. The VAM appeared to be more extensive in the roots associated with wettable areas, but definitive conclu-

(Continued on Page 13)



DISTRIBUTORS OF:

RAIN BIRD

Hunter

CRELINE

AND OTHER QUALITY TURF IRRIGATION PRODUCTS.

- CONTROLLERS
- HEADS
- VALVES



RAIN BIRD GOLF

- PIPE/FITTINGS
- PUMPS/METERS
- WIRE/VALVE BOXES

LOCATIONS:

15400 28th Avenue North
Suite 2
Plymouth, MN 55447
(612) 559-0057

415 Sundial Drive
St. Cloud, MN 56387
(612) 259-6448

1971 Seneca Road
Bay "H"
Eagan, MN 55122
(612) 687-0282

Biology and Control—

(Cont. from Page 12)

sions should be avoided since the soil was already exhibiting LDS when the samples were collected. Therefore a cause and effect relationship could not be determined. No attempt was made to rate the colonization by the other fungi they were just observed in roots from both areas.

Results from these studies indicate that the role of the bentgrass root system, and associated microflora, on the development of LDS should be investigated in more detail. Previous studies have attempted to characterize the chemical and physical properties of LDS soils, but the impact of biological influences on its development cannot be ignored.

HARTMAN TREE FARM

WHOLESALE NURSERY GROWERS



SINCE 1973
8099 BAVARIA RD.
VICTORIA, MN 55386

* BUY DIRECT FROM THE GROWER, QUALITY SELECTION OF B & B SHADE TREES, EVERGREENS, AND ORNAMENTALS

* WE INVITE YOU TO TOUR OUR GROWING RANGE

* PHONE: (612) 443-2990

* TOLL-FREE: 1-800-473-4812 * FAX: (612) 443-2835



WILLIAMS & GILL

GOLF COURSE ARCHITECTURE
LANDSCAPE ARCHITECTURE
LAND DEVELOPMENT PLANNING

GARRETT GILL

421 North Main Street
River Falls, WI 54022

(715) 425-9511

Now
Apron
Treated



THE CROWNING TOUCH

To
Protect
Your
Investment

CROWN SERIES: Formulation blended for the Upper Midwest.
Choose the following Crown Series of turf formulations that meets your needs.

Coach's Formula For Athletic Fields

- ★ "Wear tested" elite Kentucky Bluegrasses and Turf Perennial Ryegrasses.
- Developed with the turf superintendent's requirements in mind.
- Recommended for athletic complexes, playfields, stadiums, high traffic areas around golf course fairways, tees, shopping centers, picnic areas, and parks.

Elite Landscape Formula

- Meeting today's demands for a low maintenance quality turf with a balanced combination of Kentucky Bluegrass, fine Fescue, and Perennial Ryegrass.
- ★ For new lawns and damaged turf around homes, parks, playgrounds, cemeteries, commercial and government buildings.

Shade Premium Formula

- Shade tolerant balanced combination of fine Fescues and Kentucky Bluegrasses that tolerate shade and are able to tolerate shade diseases and maintain an attractive turf.
- Ten percent Ryegrass provides quick cover.

Sod Growers Premium Formula

- ★ Top Quality Kentucky Bluegrasses combined to produce a premium sod.
- At harvest, the turf is attractive, dense, and yet tough, producing a liftable sod.
- ★ Quartet, blend of four perennial ryegrasses for high wear and quick establishment.

Turf Renewal Formula

- ★ Designed to redevelop vigor and beauty of old turf by introducing modern day grasses. This formulation produces fast erosion-resistant cover and helps with "on-time" fulfillment of contracts.

Turf Pride Formula

- Developed to produce a "turf proud" lawn which will maintain its premium quality for many years with moderate care.
- Recommended for home lawns, condominiums, important commercial and government buildings that require high quality turf with moderate to full management programs.

Crown Series Turfgrass formulas are distributed in the Upper Midwest by Brayton Wilbur-Ellis.

ADVANTA



- Plant Protectants
- Top Quality Grass Seed
- Plant Foods
- Herbicides

FOR MORE INFORMATION CONTACT
LARRY THORNTON
800/642-2736 or 612/723-0151