(Continued from Page 10)

if the superintendent had collected 10 "Hoffer tube cores" in a random fashion, "What Kind of a Toad is a Nematode?" would have been concluded with Chapter

Paul Diegnau came to Keller as Keller's new superintendent in 1996 convinced that plant nematodes had not been responsible for the poor playing condi-

tions of several of the greens in 1995. My first meeting with Paul was polite, civil and definitely chilly. Dr. Stienstra and I visited Paul in his office in the old Ouonset hut maintenance building that Keller had at that time. He describes his 1996 attitude best on page 20 of the October, 2005, issue of Hole Notes by saying "I was of the same mind set" (nematodes are a 'southern' problem and not an issue in the northern climes) ten years ago. Nematodes??? Get serious!" Paul uses the term "holistic" to describe his approach to culturing quality turf. And that approach with no direct attention paid to plant nematodes seemed to work because in the early summer of 1996 a sports

columnist for the St. Paul Pioneer Press described Keller as "never looking better." And so what could I do but retreat back to my once again tarnished Ivory Tower at the "U"! Don White continued to include a lecture and laboratory devoted to plant nematodes in his fall quarter course. And Paul turned out to be a fairly agreeable fellow after all. Every fall we made a pilgrimage to Keller so that the students could learn first-hand what Paul was doing with the golf course that by all rights should have had a nematode problem but really(?) didn't. Paul explained (again as stated in October, 2005 Hole Notes) to the students in each of those classes that: "My theory has always been that any damage inflicted by nematodes can be overcome with proper cultural programs and by maximizing plant health." He always allowed the students to use Hoffer tubes to try to find the progeny of those infamous Keller nematodes. And they always found them even though some of the student-designed sampling protocols that were followed such as taking the soil cores only from ball marks were more than a bit questionable.

And then came the summer of 2005,

another "beast of a year for growing turfgrasses" when Paul and 6 other superintendents who responded to Paul's survey (Hole Notes, October 2005) acknowledged that there had been outbreaks of Take All Patch on their courses that summer. Paul wrote that "I have a feeling that the current golf season brought nematode damage to the forefront and under the spotlight. I find it hard to believe that the quantity and variety of chemistries that we - and others - threw at our TAP outbreak were so ineffective in arresting this



No. 8 at Keller Golf Course in St. Paul

pest. There must be other factors at work here. Instead of 'grasping at straws', I am hopeful this study (more intensive, less extensive nematode survey #2) will provide some answers." Please do not consider Paul Diegnau to be "a lone voice crying in the wilderness." He has some pretty impressive supporters like Mr. Oscar L. Miles, the recently retired superintendent of The Merit Club of Libertyville, Illinois. Mr. Miles has given us permission to share the following information that he provided to Paul on August 29, 2005. "Back in 2001 we had a similar problem and found TAP and very high numbers of parasitic nematodes, both stunt and ring. For us, these two problems seem to work together. Nematodes weakening the plant with TAP becoming the primary problem." On September 1, 2005, Mr. Miles supplied additional information about his TAP and plant nematode experiences: He had two labs confirm that he had plants collapsing from TAP. He went on to say: "I couldn't believe this was the complete answer to our problem so we literally dug deeper and took another set of samples to the University of Wisconsin to have a nematode check done. Several weeks later I

heard from the UW Nematologist. She was very excited. She had never seen Tylenchorhynchus - stunt (nematode) numbers this high this far north. They were at 940 per 100 cc's of soil sample. Remember, this is the following year after the 2000 United States Women's Open at the Merit Club, where we had subjected our PennLinks greens to stressful growing practices to maintain the degree of ball roll distance and smoothness required for championship play. That summer and fall after the Open and into 2001 we attempt-

> ed to maintain the Open speeds. THIS WAS A BIG MISTAKE, I am sure that we had TAP and harmful Nematodes before the greens were subjected to this stressing maintenance but, we were able to mask it with additional leaf tissue from higher HOC and fall and following spring TAP protectants drenched in to protect the roots."

Take All Patch, the real reason why this "What Kind of Toad is a Nematode?" series came into being, is caused by a soil-inhabiting, root-rotting fungus called Gaeumannomyces graminis. The fungus causes a root rot that kills the epidermal and cortical cells of the infected root. Presumably its activities create an environment that is detrimental to most

if not all plant nematodes because those nematodes have to feed on living plant cells and the root rot deprives them of such cells. Although various kinds of plant nematodes have been shown to be able to survive for a period of months without a food source, most of them can be expected to perish quite soon particularly in the heat of summer when 1) their heat-enhanced activities will cause them to rapidly utilize their stored food reserves and 2) the decomposition products released by the diseased roots could be expected to stimulate the activity of competing and perhaps antagonistic microorganisms and/or result in the release of natural nematicides from decomposing plant tissues. Although we all recognized that starting an intensive plant nematode study in September-October, 2005, had many similarities to the message conveyed by that old saying about "Locking the barn door after the horse has been stolen," three superintendents who had fought TAP in 2005 (Forest Hills, Keller, and Mankato) agreed to participate in the

(Continued on Page 12)

(Continued from Page 11)

study. Although the TAP-affected greens were slowly on the way back to the quality levels expected by the superintendents, those superintendents attempted to identify locations where TAP had been present and other areas that had remained healthy. They collected soil samples from specific sites that could be found again. They collected 5 one-inch diameter Hoffer tube cores from small areas of about 64 square inches. Soil from the root-zone at a given sampling site went into one plastic bag as a sample. Soil from below the root-zone to a depth of 6 inches was combined in another plastic bag as another sample. And, if it was physically feasible, soil cores from the 6 to 12 inch deep layer were combined in another plastic bag as a third sample from a given sampling site. The samples from 18 sampling sites at Forest Hills, 36 at Keller, and 19 at Mankato were stored in a refrigerator at St. Paul until they could be processed for nematode extraction by the Cornell piepan method.

Well, what did we prove by sampling TAP-infested greens for nematodes after

the disease had run its course? At Forest Hills there did not seem to be any relationship between the numbers and kinds of plant nematodes that were present AFTER THE EPIDEMIC and the susceptibility of the turfgrass to TAP. Each of the three severely-afflicted greens that were sampled for plant nematodes has its own built-in weaknesses. Green #16 is an elevated green with good drainage that is shaded and does not receive much morning sunshine. Soil samples collected from 6 locations on Green #16, which Superintendent Marlow Hansen identified as being the green that was most severelyaffected by TAP, contained very few plant nematodes. None contained enough stunt (300/100 cm-3) or any other kind of plant nematode to qualify as having met the population densities necessary to cause damage (all by themselves) to cool-season turfgrasses (Nelson, 1995). TAP was estimated to be equally severe on Greens #5 and #6. Green #5 has been a "good green" although it is poorly drained. Green #6 is disease-prone, low-lying and shaded. Three of the 6 sampling sites on Green #5 were found to contain large (910-1825) and potentially significant root-zone populations of stunt nematodes. The other 3 sampling sites contained "unimportant" root-

zone populations (0-264) of stunt nematodes. Two of the 6 sampling sites on Green #6 were found to have potentially significant root-zone stunt nematode populations of 445 and 997/100 cm-3 of soil. The other 4 sampling sites each contained "theoretically unimportant" root-zone populations of 37-98 stunt nematodes.

Although Superintendent Hansen has never seen any evidence in 18+ years that: 1) plant nematodes have adversely affected the growth of turfgrasses at Forest Hills and 2) no direct relationship between the incidence and severity of TAP and plant nematodes was discovered in 2005, he has chosen to continue with and significantly increase the scope of nematode study there in 2006. Each of the 3 greens for which we now have "nematode data sites" will be resampled in September and up to an additional 6 sampling sites per green will be added. An additional green will also be intensively sampled as described above. "Why," you say, "for what purpose?" Well, this is our current thinking: If the summer of 2007 turns out to be another "doosy" as Paul Diegnau chooses to describe a "tough year for

(Continued on Page 13)

Meeting Your Needs With Quality Seeds

SEED RESEARCH BLENDS

MAGNUM GOLD - Perennial Rye Grass Blend

BLUE GRASS BLENDS - Coaches and Elite Landscape

TRADITION - Fescue Blends

Improve Your Course With the Following Varieties of Bentgrass:









Fax: 952-949-3889

PHONE: 612-804-1692 Superior Unit Services, Inc.

LARRY THORNTON





Fall colors at Rutgers Bay Lake Lodge

(Continued from Page 12)

growing turfgrasses", then hopefully we will be in the position to determine if there is any kind of correlation between plant nematode populations and the incidence and severity of TAP or any other kind of plant disease that might appear. The TAP story from Keller in 2005 was quite similar to the one from Forest Hills. At least 3 greens were severely affected and TAP was also recognized as being present on additional greens. Paul Diegnau has already written about his efforts to deal with the disease. He tried a variety of different fungicidal chemistries with no effect. He increased the level of TLC that each infected green received. It was an extremely stressful summer for him, a "doosy" as he described it.

Superintendent Diegnau selected 12 sampling sites on each of 3 of Keller's greens that had been most severely affected by TAP. He selected sites where either symptoms were still visible or where he remembered that the disease had been present. He selected other sites where the bentgrass had remained healthy. The TAP status of the locations where the soil samples (root-zone, below root-zone to 6 inches, and 6 to 12 inches) were collected was not identified to the Plant Nematology Laboratory. Root quality-quantity at each sampling site was estimated on the basis of soil volume when the samples were processed. Five root-zone cores from an area of reasonably healthy golf green grasses will yield something more than 100 cm-3 (4/5th cup) of soil. On that basis, the turfgrasses of Keller Green #6 were the weakest (only one of 12 root-zone samples yielded more than 100 cm-3 of soil) of those of the 3 greens that were sampled. The turfgrasses of Keller Green #9 had considerably better roots than those of #6 with 7/12 root-zone samples yielding at least 100 cm-3 soil. The plants sampled on Green #18 (a seriously-affected green in 1995) were the strongest with 10/12 root-zone samples yielding at least 100 cm-3 soil. The total number of plant nematodes extracted from the root-zone samples from greens 6 and 9 were about equal at 4165 and 4332, respectively. Fewer plant nematodes, 3428, were extracted from the "better" root-zone samples from Green #18. Potentially significant populations of stunt nematodes of

300 or more/100 cm-3 soil were present in 5/12 and 6/12 root-zone samples from Greens #6 and #9, respectively. Only 2/12 root-zone samples from Green #18 were estimated to be infested with 300 or more stunt nematodes. The largest population of stunt nematodes (830) was estimated to be present in a root-zone sample from Green #6. It may be appropriate and useful to use the information given above to recognize a relationship that seems to exist for many plant nematodes. Severely stressed plants eventually can only support small populations of nematodes. Moderately stressed plants (Greens #6 and #9) will support larger populations. And less stressed or more vigorous plants (Green #18) will support fewer plant nematodes. Plant nematodes have to feed on living plant cells. But how, biochemically or otherwise, do the cells of a moderately stressed plant differ from those of either a severely stressed or especially a less stressed one? It is an intriguing but unanswered question with implications for future management of plant nematodes!

(Continued on Page 14)

(Continued from Page 13)

Please continue to remember that we were in a "reactive mode" in 2005 as we tried to extrapolate back to the role that undefined plant nematode populations might have had that subsequently facilitated the development of the symptoms of TAP later in the summer. This is what Paul Diegnau wrote after he had time to compare the plant nematode data with the incidence and severity of TAP for the three Keller greens that were sampled in September, 2005: "I just looked at my mapping and labeling system for our sample taking adventure last September. I was rather excited after looking at the first 6 sample sites as the high nematode populations correlated with the Take-All-Patch. Then everything went 'willy nilly,' so to speak. I think the TAP vs. non-TAP sampling site selection process was entirely too subjective. We most likely sampled the TAP sites at various stages of the disease (mostly later) on into various degrees of recovery. As I recall, SOME of the TAP sites I was selecting were not definitive; I was taking my best guess. I would assume that there are some ongoing population dynamics occurring with the nematodes as the TAP infections run their course through those localized bentgrass communities and their food sources decline??? So

the \$10,000 question becomes: "How much does the timing of the sampling relative to the progression of the TAP impact the sampling results? How quickly do these populations shrink once the bent-grass roots are no longer a viable food source? Or do they simply shift over to the Poa annua roots?"

Fred Taylor, Superintendent at the Mankato Golf Club, believes that Take-All-Patch was probably present on his course in 2004, and its presence was the reason why he first became interested in plant nematodes. The fungus was identified as being present in 2005 on several of Mankato's greens causing cosmetic effects that concerned him but apparently did not concern (no complaints) the players. The symptoms persisted and were still very apparent to him in early November. The three greens at Mankato for which we now have nematode data all seem to be both rather uniformly (in distinct contrast to Forest Hills) as well as heavily infested with plant nematodes. Mankato's Green #3 with 425-1599 stunt nematodes, #6 with 551-1188 stunt nematodes, and #14 with 637-3885 stunt nematodes/100 cm-3 of soil all seemed to be especially vulnerable to outbreaks of Take All Patch or any other disease caused by a fungus whose causal agent might benefit from the biochemical-mechanical damage caused by plant nematodes. Of those greens, #14 with approximately 39% more total plant

nematodes than #3 and 78% more total plant nematodes than #6 would seem to be most vulnerable.

In 2006, Take All Patch was judged to be entirely absent from Forest Hills and Keller and of very limited and only an early season occurrence at Mankato. Mankato's Green #3 was the only one where its symptoms were observed. Those three superintendents controlled TAP in 2006 with a preventative spray program that began with applications in the fall of 2005 and continued with additional applications early in the spring of 2006. Although any correlation between the incidence and severity of TAP and abundance of plant parasitic nematodes has not been strengthened and may have been weakened by the observations made during the past 12 months, we are enthusiastic about having the opportunity to make some further progress toward obtaining a better understanding of the possible significance of the "northern" turfgrass nematodes.

Chapter IV of "What Kind of a Toad is a Nematode?" will conclude the series with additional information that was obtained from the soil samples collected in 2005. We will also address a new concern about the health of turfgrass roots that has surfaced as the result of our being able to learn something about what is going on in the plant root-zones of so many different greens.

Make Plans to Attend the MGCSA Awards & Recognition Banquet



December 5, 2006

Rush Creek Golf Club Maple Grove

Watch your mail for your invitation!

Classified Ads

FOR SALE

Lots of ten beige colored 1996 Club Car gas golf cars ... \$1,500 ea. or best offer ... discount on multiple car purchases. 2004 beverage unit refreshment cart, pull behind or stand alone ... \$2900 or best offer. 1991 Toro Greensmaster 3000 triplex 11-blade reel mower, 3,733 hours, good shape, runs strong ... \$1,200 or best offer. Contact: Guy W. Leach Spring Valley Golf Course 715-778-5513

FOR SALE

Toro Mulcher Kit for 455-D Groundsmaster, 3 piece set, Toro OEM, Part# 30475 -- \$200 Contact: Rick LaPorte or Jeff Hanson Tartan Park Golf Course 651-736-8797

FOR SALE

1995 Toro Triplex, 2,400 hrs., new reel stock \$3,000 1987 Scotman Ice Machine \$500 16 Portable Ballwasher Stands

(Par Aide) \$40 each/\$600 for all 16 12 Mounting Pipes (Par Aide) \$15 each/\$150 for all 12 6 Divotmate Containers w/ low stand (Par Aide) \$25 each/\$125 for all 6 19 Tournament Flagsticks (8ft) \$200 for set of 19 Par Aide Protection net (new) \$100 Parts Washer \$75 Range Ballwasher \$100 Jacobsen 548 slit seeder (PTO) \$500 Jac 524 slit seeder (walk behind) \$350 Olathe Core Pulverizer 1990 \$750 Turfco 2.5ft walking topdresser \$100 Contact: Jeff Johnson or Ken Diethert The Minikahda Club 612-926-4167

FOR SALE

2004 DMI Speedroller (ride-on greens roller) like new, 105 hours, \$5,000 or best offer Contact: Eric Ritter Spooner Golf Club 715-635-3580

FOR SALE

(26) vt-4 Toro Controllers Includes clocks & pedestal - \$315 Reconditioned VT-4s for \$275 ea. Contact: Brad Hable Bunker Hills Golf Course 763-755-4150

FOR SALE

5 Ft. 3 pt. Aerway with sport tines, used very little -- \$1900 Cushman Core Harvester with workman kit -- \$2,175 Toro 3020 Sand Pro with blade and box scraper -- \$3,875 1,300 hrs., very good shape. Contact: Scott Wersal Ridges at Sand Creek 952-492-2644, ext. 24

FOR SALE

2005 Toro Flex 21 Greens Walk-Behind 200 hr. -- \$5,000 Contact: Butch Brown Birch Bay Golf Course, Nisswa 218-820-2158

Please contact the MGCSA office after items have been sold or obtained.





OVERALL PRODUCT IMPROVEMENTS


- Reduced overall weight to minimize turf compaction and maximize turf health.
- CrossTrax™, a four wheel version of our patented Series/Parallel drive system.
- More powerful models with Kubota 35.5 and 44.2 horsepower diesel engines.
- All routine service can be conducted from the left hand side of the unit.



MTI Distributing, Inc. 4830 Azelia Avenue N. Brooklyn Center, MN 55429 800-362-3665 763-592-5600 MTI Distributing, Inc. 4310 Main Avenue Fargo, ND 58103 800-782-1031 701-281-0775



Count on it.







Introducing the revolutionary Toro® Sand Pro. With its patent-pending Quick Attach System (QAS), operators can switch out most of the 17 front, rear and mid-mount attachments in seconds. Realize the true potential of your equipment with the right tool for the job. The new Sand Pro with QAS, this changes everything.

www.Toro.com/SandPro



Count on it.

It's in the Hole



MGCSA Service Awards

The following MGCSA members will be recognized at the MGCSA Awards and Recognition Banquet at Rush Creek Golf Club in Maple Grove on December 5.

40-Year Service Awards: Roger Kisch, Southview Country Club; Orland Maenke, Olympic Hills Golf Club; Joe Moris (retired), and Larry Vetter, Minnesota Turf & Grounds Foundation.

35-Year Service Awards: James Brooks, Riverside Town & Country Club; Douglas Dieter, Fox Hollow Golf Club, and Steve Young, Plaisted Companies.

30-Year Service Awards: Kerry Glader, Plaisted Companies; Douglas Hausman, CGCS, Dakota Dunes Country Club; Thomas Johnson, New Richmond Golf Club; Dave Krupp, Precision Turf & Chemical; Douglas Mahal, CGCS, The Jewel Golf Club; Charlie Pooch, Les Bolstad University of Minnesota Golf Course, and John Steiner, CGCS, White Bear Yacht Club.

25-Year Service Awards: Scott Austin, CGCS; James Crawford, Town & Country Club; David Ksiazek, Phalen Park Golf Course; Mike Leitner, Leitner Companies; Glen Lentner, CGCS, Inver Wood Golf Course; Guy Lohman; Thomas Meier, Le Sueur Country Club; John Monson, Long Prairie Country Club; Robert Reihe, Northway Irrigation; Craig Vigen, CGCS, Fargo Country Club, and William Whitworth, CGCS, Northfield Golf Club.

Golembiewski Hired as Assistant Professor In the Department of Natural Resources At the University of Minnesota Crookston

Dr. Rob Golembiewski was hired as an Assistant Professor in the Department of Natural Resources at the University of Minnesota Crookston campus with the responsibility of teaching and directing the Golf & Turf Management program. Most recently he was owner and president of Paramount Landscape located in Phoenix, Arizona. Dr. Golembiewski received both his B.S. in Crop & Soil Science and M.S. in Botany & Plant Pathology from Michigan State



University, and his Ph.D. in Agronomy from The Ohio State University. Upon completion of his Ph.D., he was hired as an Assistant Professor of Horticulture at Montana State University where he spent two years establishing the turfgrass science program. Dr. Golembiewski then spent one and a half years as the Eastern U.S. Turf & Ornamental field development biologist for Dow AgroSciences evaluating pesticides and conducting field research.






- SALES AND LEASES
- NEW USED REBUILT
- SERVICE & REPAIRALL MAKES
- PARTS, BATTERIES & ACCESSORIES
- RENTALS AVAILABLE

www.versatilevehicles.com

Brainerd 218-824-3533 18188 Highway 371 North • Brainerd, MN 56401 NEW ADDRESS

Savage 952-894-1123

5345 West 125th Street • Savage, MN 55378




Dr. White, surrounded by his biggest support group – his family – at his Hall of Fame induction ceremony. Pictured below is Jeff Johnson, a former student of Dr. White's and now Superintendent at The Minkahda Club in Minneapolis.





Mike Sonnek, Spring Hill Golf Club, Erin McManus, Medina Golf & Country Club attended Dr. White's Hall of Fame ceremony.



An Adventure in Life

By Randy Witt, CGCS

Senior Superintendent, Hong Kong Golf Club

One of the changes within myself over the years has been the development of patience. The once young, hungry, demanding, and at times intolerant individual has developed into a person that has learned patience and a good sense of

humor are very important tools in the golf course business. Many aspects of this business we have no control over, no matter how hard we try to be in control. Of course weather is the single most important factor which we deal with and have no control over whatsoever. Drought, wetness, heat, humidity, storms, etc. We deal with these factors but have no control over. Over the years I've learned that the weather and Mother Nature are the eventual winners we just cope with both. Many times our ability to be patient and cope and not panic will determine if we are successful in our endeavors.

More importantly, our ability to be patient will many times dramatically affect our health and well being, which ultimately is the most important factor in the end when all is said and done.

Patience has become an important issue as I've progressed along my career path here in Hong Kong, along with the ability to laugh. Some of the problems and situations I've encountered require patience and the ability to laugh or one would probably not survive for very long here. For starters, the size of the staff, 100 plus individuals, lends itself to the law of averages of craziness to occur. Equipment getting buried in mud. Sand Pros starting on fire, delivery trucks taking a wrong turn and getting stuck in a rough area, everyday type occurrences. Without patience and seeing the humor in the situation, one would soon have a heart attaché or be overcome with stress. The

sun will rise another day.

In the case of the sand pro catching on fire, the American saying of "A Chinese Fire Drill" immediately came to mind. A Sand Pro had developed a carburetor problem; so two mechanics were transfinally put the fire out. This whole process covered at least 10 minutes. The end result was one barbecued and destroyed sand pro. Viewing this Chinese Fire Drill was so funny that any sense of frustration and anger never really came



Authentic Chinese fire drill.

porting the machine back to the shop facility. The machine stopped, and in the process of trying to get it restarted, a fire broke out. I came upon the scene as one mechanic was attempting to put out the fire with the use of his shirt. This only served to make a bigger fire. The next step was the Second Mechanic ripping of his shirt and soaking it in a nearby stream and trying to throw his wet shirt on the fire. This too proved useless. Step 3 was to run to a nearby bunker, get some sand, and try to bury the fire. Step 3 was also a failure. The final step was for a shirtless mechanic to drive back to the maintenance facility, get a fire extinguisher, return and

out. What good would it do. No one got hurt and it's over with. Just another tale to be told, another chapter for the book.

May through August is the rainy season in Hong Kong and many parts of Asia. Each month we have received in excess of 500 mm of rain, with 25 mm equaling 1 inch. So, 20 inches of rainfall in 30 days can make for some very wet conditions at times. Even though drainage is pretty good, many rough areas can be difficult to mow. The photo above shows what happens when the Chinese persist in trying to mow areas of turf with standing water in them.

(Continued on Page 21)