

A second study was set-up to examine the effect of mefluidide (Embark) and amidochlor (Limit) applied at different growing degrees on the seedhead and vegetative growth suppression of 'Kenblue' kentucky bluegrass. Seedhead suppression was excellent for applications made at 25, 50, 75, and 100 GDD. Seedhead control was lost at the 150 GDD date. Figure 1 shows the effect of application date on clipping yields. Suppression of clipping yield was a fairly constant percentage of the check. It would appear that applications at or before 100 GDD will provide excellent seedhead and vegetative growth suppression of annual bluegrass. This is unusual since ethofumesate is considered a preemergent herbicide. Rainfall within 24-48 hours after application appears to reduce its effectiveness, contrary to most preemergent herbicides which need rainfall to activate the herbicide. Tables 4 and 5 show the effect of ethofumesate on Penncross and Penneagle creeping bentgrasses. Injury to Penncross at the 1.5 lb/A rate was noticeable, however, not severe enough to cause loss of turf. The optimum rate would appear to lie between 1 and 1.5 lbs/A. We did observe a shift to more Kentucky bluegrass and creeping bentgrass and less annual bluegrass in plots treated with ethofumesate. This study will be expanded in 1986.

Variety Trial Evaluations

Our variety trial evaluation program is expanding and in 1985 fine fescue, Kentucky bluegrass and low maintenance variety trials were established. Results from the 1985 evaluations of Kentucky bluegrass, fine fescue and perennial ryegrass trials are shown in Tables 6, 7, and 8. In examining these tables, the LSD value at the bottom of each column should be used as guide in evaluating the cultivars. Cultivars whose means differ by less than LSD value are not considered statistically different in terms of performance.

Preemergent Herbicides and Cultivation

This study was designed to examine the effect of cultivation and time of cultivation on the performance of three commonly used preemergent turf herbicides. This was the second year of this study. Four herbicide treatments, benefin, bensulide, DCPA, and a control, were applied on May 1 of 1984 and 1985. Immediately after herbicide treatment, four cultivation treatments were applied to one half of the plot area. The cultivation treatments consisted of core cultivation one pass or three passes, vertical mowing, and a control. Four weeks after herbicide application, the cultivation treatments were applied to the other one half of the plot area. Results are shown in Tables 2 and 3. After two years of data, it appears that cultivation operations can be safely performed without significantly decreasing annual grass control.

Ethofumesate for annual bluegrass control

Ethofumesate (Prograss) is a herbicide used to control annual bluegrass in established Perennial Ryegrass. A study was established to determine its activity on annual bluegrass and tolerance on creeping bentgrass. Applications were made at rates of 0.75, 1.0, and 1.5 lbs/A. Most treatments were applied two to three times at one month intervals. Figure 2 shows injury patterns on annual bluegrass treated at four different rates. A 0.75 lb/A rate was applied three times on 9/6, 10/1, and 11/1. The other three treatments were applied 9/15 and 10/15 at rates of 0.75, 1.0, and 1.5 lbs/A on each application date. Injury to annual bluegrass showed a typical rate response with 1.5 lbs/A

Table 6. 1985 ratings for the USDA National Kentucky bluegrass tests.

<u>Entry</u>	<u>Quality Ratings</u>					<u>Grand Mean</u>
	<u>4-18</u>	<u>7-3</u>	<u>8-28</u>	<u>9-26</u>	<u>10-30</u>	
Majestic	4.7	7.0	6.3	7.3	7.7	7.1
RAM-1	3.3	6.7	6.0	7.3	7.7	6.9
Bristol	3.7	4.3	7.3	7.3	7.7	6.7
Bonnieblue	6.0	5.0	6.3	7.0	8.0	6.6
Glade	3.0	6.0	6.3	6.7	7.3	6.6
Trenton	6.0	4.7	6.3	7.0	8.0	6.5
Eclipse	3.3	7.3	6.0	6.3	6.3	6.5
Fylking	2.7	6.7	6.7	6.0	6.3	6.4
Vanessa	3.0	6.3	7.0	6.3	6.3	6.4
Rugby	5.0	4.0	6.3	7.7	7.3	6.3
Enmundi	3.3	6.0	6.0	6.3	6.7	6.2
Barblue	4.0	5.7	6.3	6.0	6.7	6.2
Sydsport	4.0	6.7	5.7	5.7	6.3	6.1
Columbia	4.7	4.7	6.0	6.7	6.7	6.0
Holiday	3.7	6.7	5.3	6.3	5.7	6.0
Kimono	3.0	6.3	6.0	5.7	5.7	5.9
Nugget	1.7	6.7	6.0	5.0	5.7	5.9
Victa	4.0	5.7	5.7	6.0	6.3	5.9
Mosa	2.3	5.7	5.7	6.3	5.7	5.9
Birka	4.3	5.3	6.0	6.3	6.0	5.9
Merion	4.3	5.7	5.0	6.3	6.0	5.8
Adelphi	5.7	5.3	4.7	6.3	6.7	5.8
Merit	3.7	5.7	6.0	5.3	6.3	5.8
Cheri	3.7	5.3	5.3	6.0	6.0	5.7
Touchdown	4.3	5.3	6.0	5.7	5.7	5.7
Baron	3.0	5.3	5.0	6.3	6.3	5.7
Plush	4.3	7.0	5.0	5.3	5.7	5.7
Parade	4.3	4.7	5.0	6.0	6.3	5.5
Bayside	5.3	3.7	5.7	6.0	6.3	5.4
Gello	4.0	4.0	5.3	6.3	6.0	5.4
Shasta	4.3	3.0	5.7	6.3	6.7	5.4
Geronimo	5.0	3.7	6.3	5.3	6.0	5.3
A-34	5.0	4.7	5.3	5.7	5.3	5.2
Welcome	2.3	6.0	5.7	4.3	5.0	5.2
Enoble	2.0	3.7	5.0	5.3	6.3	5.1
Harmony	2.0	5.7	5.0	4.7	5.0	5.1
Mystic	7.3	6.0	5.3	4.0	4.7	5.0
Wabash	7.3	4.7	4.0	5.3	5.7	4.9
Monopoly	3.7	4.3	4.7	5.0	4.7	4.7
America	4.3	3.3	5.0	4.7	4.7	4.4
Kenblue	4.7	3.0	4.0	4.3	5.7	4.3
Vantage	5.7	2.3	4.0	4.7	5.0	4.0

Table 7. 1985 ratings for the fine fescue variety trail.

<u>ENTRY</u>	<u>Evaluation Date</u>				<u>GRAND MEAN</u>
	<u>7-3</u>	<u>8-28</u>	<u>9-25</u>	<u>10-30</u>	
Valda	6.3	7.3	5.7	5.7	6.3
Biljart	7.0	7.0	6.3	5.0	6.3
Scaldis	6.3	7.3	5.3	6.0	6.2
ST-2	6.7	8.0	4.3	5.3	6.1
Aurora	6.3	7.3	5.0	5.3	6.0
Reliant	6.0	7.0	5.0	5.0	5.8
HF 9-3 (Weekend)	5.3	6.7	7.3	4.0	5.8
BAR FO 81-225	5.3	7.0	5.3	5.0	5.7
X-8	6.7	6.0	6.3	3.7	5.7
Banner	5.0	5.7	7.7	4.3	5.7
Mary	5.3	7.3	5.7	4.0	5.6
Beauty	5.7	7.0	5.3	4.0	5.5
Shadow	5.0	6.3	6.3	4.3	5.5
Enjoy	5.0	7.0	5.0	4.0	5.3
CF-2	4.7	6.7	6.3	3.7	5.3
Waldina	5.0	7.3	4.0	4.7	5.2
Longfellow	5.0	5.7	5.7	4.3	5.2
Magenta	5.3	6.0	5.7	3.7	5.2
Pennlawn	5.0	6.0	5.3	4.3	5.1
Jamestown	5.7	6.3	5.0	3.3	5.1
Tamara	5.0	6.3	5.0	4.0	5.1
Flyer	5.0	6.7	5.0	3.3	5.0
X-11	6.0	5.0	5.0	4.0	5.0
Wilma	4.0	6.7	5.0	3.3	4.8
430	6.0	5.7	4.0	3.3	4.8
Atlanta	5.0	6.0	4.7	3.7	4.8
Z-16	6.0	5.0	5.0	3.0	4.8
Estica	5.7	6.3	3.7	3.0	4.7
X-13	4.0	6.0	6.0	3.0	4.7
Highlight	5.0	6.0	4.0	3.7	4.7
Epsom	5.7	5.3	4.3	3.3	4.6
Waldorf	4.7	5.3	5.0	3.3	4.6
Tatjana	3.7	5.7	5.0	4.0	4.6
Checker	5.7	4.7	4.7	3.3	4.6
Center	3.7	5.3	5.3	3.7	4.5
X-9	5.0	5.7	4.3	3.0	4.5
Koket	3.3	6.0	4.7	3.7	4.4
Lovisa	5.7	4.7	4.7	2.7	4.4
Pernille	4.3	5.7	4.0	3.0	4.3
FRI-Frt 83-1	5.7	5.7	3.3	2.7	4.3
X-12	5.0	5.0	4.0	3.0	4.3
Tvalo	3.7	5.3	4.3	3.7	4.2
Ensylva	5.0	5.3	3.0	3.3	4.1
Geres	3.7	5.0	4.3	3.3	4.1
Unknown	4.0	4.7	3.7	4.0	4.1
Commodore	3.7	4.7	3.7	2.7	3.7
Boreal	4.3	4.3	4.0	2.3	3.7
Ruby	3.7	4.0	3.7	2.7	3.5
Wintergreen	3.7	4.7	3.0	2.7	3.5
Robot	3.0	4.0	3.3	2.0	3.1
LSD(P=0.05)	1.3	1.1	1.1	1.0	

Table 8. 1985 perennial ryegrass cultivar evaluations.

<u>Varieties</u>	<u>7-9-85</u>	<u>9-9-85</u>	<u>9-26-85</u>	<u>10-30-85</u>	<u>Grand Mean</u>
Bellatrix	6.0	5.0	7.0	6.7	6.2
Regal	4.0	6.3	6.3	7.0	5.9
Crown	4.3	5.7	6.7	7.0	5.9
Yorktown II	6.0	5.3	5.7	6.7	5.9
Ranger	5.0	5.7	6.0	7.0	5.9
Belle	4.7	5.7	6.3	6.3	5.8
Idole	5.3	5.7	6.3	6.0	5.8
Premier	4.0	6.3	6.0	7.0	5.8
Derby	4.3	5.7	6.3	7.0	5.8
Dasher	4.0	6.3	6.0	6.0	5.6
Blazer	5.3	5.0	6.0	6.3	5.6
Acclaim	4.3	5.3	6.0	6.3	5.5
Fiesta	4.3	5.7	5.7	6.3	5.5
Clipper	3.3	6.3	6.3	6.0	5.5
Fronto	4.3	5.7	6.0	6.0	5.5
Ensporta	6.0	5.0	5.0	5.7	5.4
Citation #153	3.7	22.7	6.0	6.0	5.4
Jackpot	5.3	5.7	5.0	5.7	5.4
CBS	4.0	5.0	5.7	6.3	5.3
Trimmer	5.0	4.7	4.7	6.3	5.2
Pennfine	4.3	6.0	5.0	5.7	5.2
Omega	3.7	5.7	5.7	5.7	5.2
Hunter	6.0	4.3	5.3	5.0	5.2
Diplomat	5.7	4.3	5.3	5.7	5.2
Delray	4.0	6.0	5.0	5.3	5.1
Princess	5.0	4.3	5.0	6.0	5.1
Pennant	4.3	5.0	5.0	6.0	5.1
Manhattan	5.0	4.3	5.3	5.7	5.1
Arno	4.3	5.0	5.3	5.3	5.0
Runner	4.7	4.7	5.0	5.0	4.8
Goalie	4.0	5.0	5.0	5.3	4.8
Player	6.0	4.0	4.0	5.0	4.8
Elka	5.3	3.7	4.7	5.0	4.7
Loretta	5.3	4.3	4.7	4.7	4.7
NK 200	3.7	4.3	5.0	5.0	4.5
Compas	4.7	3.7	4.3	4.7	4.4
Venlona	4.7	3.3	4.3	4.7	4.3
Norelea	3.0	2.7	3.3	3.3	3.1
LSD(P=0.05)	0.5	-	0.8	0.5	