

1988 Turf Weed Control, PGR, and Management Studies  
Bruce Branham  
Department of Crop and Soil Sciences  
Michigan State University

The drought of 1988 offered some unique opportunities to study grasses under these conditions. One ongoing study which yielded some valuable information was an evaluation of various cool-season turfgrass species and cultivars for their performance under low maintenance conditions. This trial consisted of 48 cultivars of Kentucky bluegrass (*Poa pratensis*), tall fescue (*Festuca Arundinacea*), fine fescues (*Festuca rubra*), and perennial ryegrass (*Lolium perenne*) evaluated under conditions of no irrigation and low nitrogen fertility (1 lb N/M/YR). Data in table 1 show the percent survival of each species. The two species which fared the best were tall fescue and perennial ryegrass. Surprisingly, the fine fescues did not survive the drought very well which was unexpected because the fine fescues have always been touted as low maintenance, drought hardy grasses. The Kentucky bluegrasses also did not do very well but some of their lack of survival can be attributed to the difficulty in establishing bluegrasses and the gradual deterioration seen with the improved, sod-quality bluegrass varieties under low maintenance. The two species that fared well in this test were tall fescue and perennial ryegrass. Tall fescue has long been noted for its ability to withstand droughts and has done so by being very deep rooted compared to other cool season turfgrasses. This trial has demonstrated that for medium to low maintenance turfs, tall fescue should be strongly considered for use in Michigan. Our trials have never shown any problems with winterkill of tall fescue in Michigan, however, this has long been a concern. Until we get more data, we still caution against the use of tall fescues on wet or poorly drained soils.

Data in tables 2-5 show the 1988 quality ratings for variety trials of Kentucky bluegrass, tall fescue, fine fescues, and perennial ryegrass. The tall fescue variety trial was established in May of 1988 so data was only available from August, September and October of 1988. Because of the number of varieties now available in all species, a good rule of thumb for selecting varieties is to select those varieties that are in the top 25% of the trial. This will assure you of selecting high quality varieties.

A study was conducted in 1988 to examine the ability of various iron sources to mask the injury seen from spring applications of Embark to control annual bluegrass seedheads. Embark was applied at 1/8 lb AI/A (8 oz/A) alone and in combination with various commercially available iron sources and the treatments were evaluated for percent seedhead control and turf quality. Data in table six describes the rate and sources of iron used in the study and shows the results in terms of turf quality and seedhead density and control. Most of the iron sources tested did provide some masking of Embark injury with Scott's Iron S granular source providing the best turf quality. Also providing good masking of Embark injury was Ferromec at 2 and 4 oz/M. The most interesting observation of the study was that many of the iron sources antagonized the efficacy of Embark. This was clearly shown in the percent seedhead control data. Only Scott's Iron-S and Agriflex provided the same level of control as Embark alone. Ferromec AC at 2 oz/M showed slight antagonism but was still effective. The Lesco Iron plus N and Ferromec AC at 6 oz/M caused a noticeable loss of seedhead control but they were not statistically different from the Embark alone treatment. All other rates and sources of iron caused a significant loss of seedhead control.

TABLE 1. Low Maintenance Variety Trial

Variety	Species	Quality Ratings			Grand Means	%	%
		4/25	5/20	7/1		Survival	Crab
5D2	Tall Fescue	4.8	6.3	3.3	4.8	83.3	3.3
Apache	Tall Fescue	5.2	6.3	2.7	4.7	86.7	3.7
Spartan	Hard Fescue	5.8	6.0	2.0	4.6	20.0	0.7
Bonanza	Tall Fescue	5.3	5.7	2.7	4.6	80.0	3.0
Scaldis	Hard Fescue	5.7	5.7	2.2	4.5	20.0	1.0
Shadow	Creeping Fescue	5.3	5.0	2.7	4.3	43.3	.3
Agram	Creeping Fescue	5.2	5.3	2.2	4.2	21.7	.3
Tempo	Tall Fescue	5.3	5.3	1.3	4.0	81.7	3.0
Bighorn	Sheep Fescue	4.0	6.0	2.0	4.0	18.3	.3
Mustang	Tall Fescue	4.8	5.3	1.7	3.9	88.3	3.3
Victory	Creeping Fescue	5.2	5.0	1.7	3.9	28.3	2.0
Calicote	Perennial Ryegrass	5.3	4.3	2.0	3.9	45.0	11.0
Kenblue	Kentucky Bluegrass	5.7	4.7	1.3	3.9	35.0	28.3
Parade	Kentucky Bluegrass	5.2	4.7	1.8	3.9	11.0	5.3
Galway	Tall Fescue	4.8	5.0	1.7	3.8	86.7	3.0
Touchdown	Kentucky Bluegrass	5.3	4.7	1.5	3.8	4.3	2.3
Houndog	Tall Fescue	4.5	5.7	1.3	3.8	91.7	1.7
Delray	Perennial Ryegrass	5.7	4.3	1.3	3.8	31.7	11.7
Tournament	Hard Fescue	4.8	5.0	1.5	3.8	12.0	1.0
Pennfire	Perennial Ryegrass	6.3	4.0	1.0	3.8	60.0	12.7
Trident	Tall Fescue	4.3	5.3	1.5	3.7	83.3	5.0
Arid	Tall Fescue	4.0	4.7	2.3	3.7	90.0	2.0
Park	Kentucky Bluegrass	5.5	4.3	1.2	3.7	27.7	11.0
Delta	Kentucky Bluegrass	5.7	3.0	2.2	3.6	13.3	28.3
Flyer	Creeping Fescue	5.2	3.7	2.0	3.6	13.3	3.0
Azay	Sheep Fescue	5.3	4.3	1.2	3.6	17.0	1.3
Jazz	Perennial Ryegrass	4.8	4.3	1.5	3.6	53.3	14.3
Goalie	Perennial Ryegrass	6.0	3.7	1.0	3.6	20.0	20.0
Aurora	Hard Fescue	4.7	4.7	1.2	3.5	10.7	2.3
Maverick	Tall Fescue	4.5	4.7	1.3	3.5	75.0	3.7
America	Kentucky Bluegrass	5.2	3.7	1.3	3.4	23.3	3.0
Nassua	Kentucky Bluegrass	5.5	3.0	1.7	3.4	13.3	38.3
Banff	Kentucky Bluegrass	5.2	4.0	1.0	3.4	10.0	2.7
All-Star	Perennial Ryegrass	4.3	4.3	1.0	3.2	55.0	7.0
Gator	Perennial Ryegrass	4.0	4.3	1.3	3.2	65.0	7.0
Georgetown	Kentucky Bluegrass	4.7	3.3	1.7	3.2	25.0	7.0
Dawson	Chewings Fescue	3.8	3.0	2.3	3.1	10.0	2.7
Ruby	Creeping Fescue	3.7	3.3	2.0	3.0	36.7	5.0
Fiesta	Perennial Ryegrass	4.5	3.3	1.0	2.9	63.3	8.0
Blazor	Perennial Ryegrass	4.3	3.3	1.0	2.9	53.3	5.3
Adelphi	Kentucky Bluegrass	4.5	3.0	1.2	2.9	21.7	10.7
NK 200	Perennial Ryegrass	4.0	3.3	1.0	2.8	46.7	16.0

TABLE 2. 1988 Quality Ratings for USDA Fine Fescue Variety Trial

Variety	Quality Ratings					Grand Means
	4/25	6/30	8/4	9/10	10/31	
Fri-Frt-83-1	6.7	7.0	7.2	7.0	7.7	7.1
Beauty	8.0	5.5	7.0	6.8	6.8	6.8
CF-2	7.7	6.2	6.0	6.7	7.3	6.8
Shadow	7.7	6.3	6.3	6.0	7.3	6.7
Aurora	7.7	6.2	7.0	6.8	5.8	6.7
Enjoy	7.2	6.8	6.0	6.2	7.2	6.7
Epsom	7.3	6.8	6.3	6.2	6.5	6.6
ST-2	7.7	6.3	7.0	6.8	5.0	6.6
HF 9-3	7.2	6.5	5.7	5.8	7.5	6.5
Atlanta	7.2	5.8	6.0	6.7	6.8	6.5
Mary	7.5	5.8	5.7	6.3	6.8	6.4
Banner	7.5	6.0	5.7	6.2	6.7	6.4
Reliant	6.7	6.2	7.3	6.3	5.3	6.4
Koket	7.7	5.2	5.7	6.3	6.3	6.2
Longfellow	7.7	5.7	6.3	5.5	5.8	6.2
Pennlawn	6.8	5.7	5.7	6.2	6.5	6.2
Tamara	6.5	6.0	6.7	5.0	6.5	6.1
Wilma	7.7	5.5	5.7	5.8	5.8	6.1
Valda	7.7	5.3	6.3	6.0	5.0	6.1
Jamestown	7.0	5.5	5.7	5.8	6.2	6.0
Center	6.8	5.3	6.0	5.3	6.5	6.0
Scaldis	6.7	5.7	6.3	5.5	5.7	6.0
Flyer	7.2	5.5	5.7	6.0	5.5	6.0
Waldorf	6.0	6.2	6.3	5.0	6.2	5.9
Checker	6.7	5.8	6.3	4.8	5.8	5.9
Boreal	6.3	4.7	6.0	6.8	5.7	5.9
430	6.7	5.0	6.0	5.0	6.8	5.9
Waldina	7.7	4.7	6.0	6.5	4.3	5.8
Estica	6.2	4.8	6.3	5.7	6.0	5.8
Tatjana	7.7	4.3	5.3	5.0	6.3	5.7
Bar FO 81-225	7.3	5.3	5.8	5.0	5.0	5.7
Magenta	6.8	5.2	5.3	5.3	5.7	5.7
Lousia	6.0	5.0	6.0	5.7	5.7	5.7
-	6.3	5.2	5.7	6.3	4.7	5.6
Penille	5.7	4.8	6.3	5.7	5.3	5.6
Unknown	6.5	4.5	5.3	6.2	5.3	5.6
Biljart	7.0	5.0	6.0	5.3	4.3	5.5
Ivalo	6.5	4.3	5.7	6.0	5.2	5.5
Highlight	6.2	4.7	6.0	4.3	6.3	5.5
Ruby	6.3	4.0	6.3	6.2	4.7	5.5
-	6.9	5.0	6.1	5.3	3.8	5.4
Ensylva	5.3	5.0	5.8	5.7	5.3	5.4
Wintergreen	6.3	4.7	5.7	5.3	5.0	5.4
Ceres	6.0	5.0	6.0	5.0	5.0	5.4
Commodore	6.7	4.7	5.5	5.3	4.7	5.4
-	6.5	5.0	5.0	5.0	4.7	5.2
Robot	6.3	4.3	5.0	5.3	4.3	5.1
-	6.9	5.0	4.1	5.3	3.8	5.0
-	7.9	4.0	4.1	4.3	3.8	4.8
-	6.9	4.0	3.1	4.3	3.8	4.4

TABLE 3. National Perennial Ryegrass Variety Trial

Variety	Quality Ratings					Grand Means	Red Thread Lesions/Plot
	5/20	6/30	8/2	9/9	10/27		
PST-2PM	7.0	6.5	6.8	7.3	6.7	6.9	2.0
PST-M2E	7.0	7.2	6.8	6.8	5.8	6.7	5.3
PST-2H7	6.7	7.2	6.5	7.0	6.3	6.7	5.7
SR-4100	7.3	6.0	7.3	6.8	6.0	6.7	0.7
Allaire	7.0	6.2	6.5	6.7	6.8	6.6	2.0
Repell	7.7	6.7	6.7	6.7	5.3	6.6	1.7
Tara	7.0	6.3	6.5	7.0	6.0	6.6	2.0
Manhattan II	7.0	6.5	6.5	6.7	5.7	6.5	3.0
ISI-851	7.0	6.7	6.7	6.7	5.0	6.4	0.7
Runaway	6.3	6.2	6.5	6.8	6.0	6.4	1.3
Pick 300	6.7	5.5	6.7	6.8	6.2	6.4	0.3
Pennant	6.3	6.2	6.7	7.0	5.7	6.4	1.0
Manhattan	7.0	6.5	6.0	6.7	5.3	6.3	1.7
Rodeo	6.3	7.3	6.0	6.2	5.7	6.3	1.7
SR 4000	7.0	7.2	5.8	6.3	4.8	6.2	0.3
Rival	7.0	6.5	5.5	6.8	5.3	6.2	0
PST-2HH	6.7	6.0	6.2	6.8	5.3	6.2	2.3
Bar Lp 410	6.7	6.0	6.5	7.2	4.7	6.2	1.3
Barry	7.0	6.3	6.0	7.0	4.7	6.2	4.7
PST-259	7.0	6.0	6.3	6.5	5.0	6.2	2.7
PST-2DD	6.3	5.3	6.5	6.8	5.8	6.2	1.7
PSU-222	6.7	5.7	6.5	6.3	5.7	6.2	2.7
PSU-333	6.0	6.3	6.7	6.3	5.3	6.1	5.7
Birdie II	6.0	5.8	6.5	6.7	5.7	6.1	0.3
PST-250	6.0	6.3	5.8	6.8	5.5	6.1	5.0
Pick-647	6.0	5.8	6.7	6.3	5.7	6.1	0
Citation II	7.0	5.5	6.3	6.0	5.3	6.0	2.3
Pick 233	6.3	5.8	6.3	6.3	5.3	6.0	5.3
Goalie	5.7	5.8	6.2	6.7	5.8	6.0	1.3
Pick 600	6.3	5.8	6.5	6.3	5.0	6.0	2.7
Palmer	5.7	5.8	6.2	6.8	5.5	6.0	2.3
Omega II	6.3	6.3	5.8	6.7	4.7	6.0	1.0
Bar LP 454	7.0	6.8	5.3	6.3	4.0	5.9	3.0
Belle	5.0	5.3	6.5	6.8	5.8	5.9	0.7
SR 4031	6.3	5.0	6.0	6.3	5.8	5.9	1.3
Yorktown II	6.3	6.7	6.2	6.0	4.3	5.9	2.0
Gator	7.0	5.8	6.3	5.7	4.7	5.9	1.0
NK 80389	6.3	7.3	5.3	5.3	5.0	5.9	2.0
Derby	6.3	4.7	6.3	6.3	5.7	5.9	0.7
Pick 715	6.3	5.0	5.8	6.8	5.2	5.8	2.3
Ovation	6.0	6.0	5.7	6.2	5.3	5.8	0.3
Prelude	6.7	5.0	6.5	6.2	4.7	5.8	1.3
J207	6.3	6.3	5.5	6.0	4.7	5.8	2.0
Regency	6.0	5.7	5.8	6.5	4.8	5.8	1.0
ISI-K2	6.3	6.7	5.3	6.0	4.5	5.8	1.7
Mom-Lp-763	6.0	5.7	5.8	6.3	4.7	5.7	0.7
Brenda	5.3	5.3	6.2	6.2	5.3	5.7	2.0
Pennfine	5.3	5.3	6.2	6.3	5.0	5.6	2.7
Sunrye	5.7	4.7	5.8	6.3	5.7	5.6	1.3
KWS-A1-2	6.0	7.0	5.7	5.5	4.0	5.6	2.3

TABLE 3 cont'. National Perennial Ryegrass Variety Trial

<u>Variety</u>	<u>Quality Ratings</u>					<u>Grand Means</u>	<u>Red Thread Lesions/Plot</u>
	<u>5/20</u>	<u>6/30</u>	<u>8/2</u>	<u>9/9</u>	<u>10/27</u>		
Diplomat	5.7	6.2	5.5	6.0	4.7	5.6	1.7
Vintage-2DF	6.0	5.3	6.2	5.3	5.0	5.6	3.7
Acrobat	7.0	4.8	5.0	5.8	5.0	5.5	0
Caliente	6.0	5.3	5.8	6.0	4.5	5.5	1.0
Ranger	6.7	5.8	5.2	5.7	4.3	5.5	5.7
Del 946	6.3	3.7	6.0	6.7	5.0	5.5	0
Patriot	6.0	6.0	5.3	5.7	4.3	5.5	1.3
Regal	5.7	4.0	5.3	6.2	6.0	5.4	1.3
Ronja	5.3	6.7	4.8	6.0	4.3	5.4	0.7
Sheriff	6.3	5.0	6.2	5.3	4.2	5.4	0.7
Cowboy	5.7	3.7	5.8	6.5	5.3	5.4	0.3
J208	5.7	6.2	5.0	5.5	4.5	5.4	5.0
Pavo	5.0	6.2	5.3	5.7	4.3	5.3	0.7
Delray	6.3	4.3	5.0	5.7	3.7	5.0	1.7
Linn	4.0	4.3	2.7	4.3	3.0	3.7	0.3

TABLE 4. 1987 Tall Fescue Variety Trial

Variety	Quality Ratings			% Cover	Grand Quality Means	Leaf Texture 1-9, 9 Finest
	8/4	9/6	10/29			
Jaguar II	7.7	8.3	6.3	83.3	7.4	6.0
Hubbard 87	7.7	7.5	6.2	85.0	7.1	5.7
PST-5HF	7.3	7.8	6.0	85.0	7.1	5.0
Pick DM	6.0	8.0	6.8	78.3	6.9	5.3
Rebel II	7.2	7.5	6.0	91.7	6.9	6.0
Monarch	7.2	7.7	5.8	86.7	6.9	5.7
Apache	7.8	7.0	5.7	86.7	6.8	6.3
Tribute	7.2	7.5	5.7	85.0	6.8	5.8
PST-5AG	6.7	7.7	5.8	83.3	6.7	6.0
Falcon	7.2	6.8	6.2	94.3	6.7	5.0
PST-5MW	6.7	7.2	6.3	81.7	6.7	5.7
Pick 127	7.5	7.2	5.3	81.7	6.7	5.0
Bonanza	7.2	7.2	5.7	83.3	6.7	5.3
Pick GH6	6.8	7.3	5.8	81.7	6.7	5.2
Trail Blazer	6.3	7.3	6.2	88.3	6.6	7.0
PST-5AP	6.8	7.2	5.7	85.0	6.6	5.0
Wrangler	6.7	6.8	6.0	95.0	6.5	6.0
Thoroughbred	6.8	7.0	5.7	93.3	6.5	5.8
Rebel	6.8	7.2	5.5	91.7	6.5	4.7
Pick DDF	6.3	7.3	5.7	73.3	6.4	5.7
PST-5OL	7.2	6.8	5.3	87.7	6.4	5.3
Mesa	7.0	6.8	5.5	90.0	6.4	5.0
Pick TF9	6.2	7.3	5.8	80.0	6.4	5.8
Legend	6.3	6.8	6.0	86.7	6.4	5.8
PST-DBC	7.0	6.8	5.3	86.7	6.4	5.0
Cimmaron	6.2	7.0	6.0	81.7	6.4	5.8
PST-5D1	6.7	7.0	5.3	90.0	6.3	5.7
Taurus	6.0	7.3	5.7	91.7	6.3	6.0
KWS-DUR	6.7	7.3	4.8	76.7	6.3	6.5
Arid	6.5	7.0	5.3	86.7	6.3	5.8
Trident	6.5	6.8	5.3	78.3	6.2	6.8
Chieftan	6.5	7.2	5.0	86.7	6.2	5.7
Sundance	6.0	7.0	5.7	86.7	6.2	6.0
Olympic	6.7	7.2	4.7	91.7	6.2	4.7
Adventure	6.7	6.7	5.2	86.0	6.2	5.7
Aztec	5.8	7.0	5.7	81.7	6.2	6.5
Richmond	7.2	6.3	5.0	88.3	6.2	4.7
PST-5BL	6.0	7.2	5.3	76.7	6.2	6.7
JB-2	6.3	6.3	5.7	88.3	6.1	5.5
Normac 99	6.3	6.8	5.2	76.7	6.1	6.0
Carefree	6.5	6.8	5.0	90.0	6.1	6.3
Titan	6.8	6.8	4.7	89.3	6.1	5.3
Finelawn I	6.7	6.7	5.0	96.0	6.1	5.0
Tip	7.2	6.0	5.0	91.7	6.1	4.7
PST-5FN	6.2	6.7	5.3	78.3	6.1	6.3
Pick SLD	5.7	6.7	5.8	81.7	6.1	6.7
Pacer	6.7	6.8	4.7	92.7	6.1	5.8
PST-5EN	6.7	6.7	4.7	88.3	6.0	6.0
Bel 86-1	6.7	6.8	4.5	80.0	6.0	5.3

TABLE 4 cont'. 1987 Tall Fescue Variety Trial

<u>Variety</u>	<u>Quality Ratings</u>			<u>% Cover</u>	<u>Grand Quality Means</u>	<u>Leaf Texture 1-9, 9 Finest</u>
	<u>8/4</u>	<u>9/6</u>	<u>10/29</u>			
Normarc 77	6.8	6.5	4.7	76.7	6.0	5.0
Pick 845 PN	6.3	6.7	5.0	76.7	6.0	6.2
Finelawn 5GL	6.5	6.7	4.7	93.3	5.9	6.0
Williamette	6.7	6.5	4.3	83.3	5.8	6.0
PST-5D7	6.0	6.6	4.9	78.3	5.8	7.1
PE-7	5.5	6.8	5.2	86.7	5.8	6.3
Bel 86-2	6.0	6.3	5.0	85.0	5.8	6.3
PST-5DM	6.0	6.5	4.8	85.0	5.8	6.2
Jaquar	6.7	6.0	4.4	81.4	5.7	4.9
Syn GA	6.2	5.7	4.7	86.7	5.5	5.3
Fatima	5.8	6.3	4.3	90.0	5.5	5.5
Ky-31	6.3	5.5	3.7	90.0	5.2	5.3
	0.0	0.0	0.0		0.0	

TABLE 5. 1986 National Kentucky Bluegrass Trial

Variety	Quality Ratings						Grand Means
	4/25	5/20	6/23	8/2	9/6	10/27	
P 104	6.5	7.3	8.3	8.7	8.5	8.5	8.0
Midnight	6.2	8.0	7.8	7.8	8.0	8.3	7.7
Bristol	7.2	8.2	7.2	8.0	7.7	7.8	7.7
Lofts 1757	7.2	7.8	7.3	7.5	7.5	7.7	7.5
Challenger	7.3	7.3	6.7	7.5	7.7	8.0	7.4
Mystic	6.3	6.7	8.7	7.2	7.8	7.7	7.4
Blacksburg	5.7	7.7	7.7	7.8	8.2	7.0	7.3
Aspen	7.0	7.0	7.0	7.8	7.2	7.3	7.2
Glade	6.8	7.7	7.5	7.0	7.3	6.8	7.2
Destiny	6.7	7.2	7.7	6.8	7.2	7.5	7.2
Ba 73-540	7.0	7.7	8.0	6.3	7.0	6.8	7.1
Eclipse	6.7	6.7	7.7	7.2	7.2	7.2	7.1
Ram-1	6.7	5.8	7.7	7.7	7.3	7.3	7.1
Able I	6.7	6.8	8.3	6.8	7.0	6.8	7.1
Ba 72-500	7.0	7.2	8.0	7.3	7.0	5.8	7.1
Liberty	7.0	6.7	7.5	7.0	6.8	7.0	7.0
Georgetown	7.0	7.5	7.0	6.0	7.2	7.2	7.0
Classic	7.0	6.8	7.2	7.0	7.5	6.3	7.0
Dawn	6.8	7.2	6.7	7.2	6.8	7.2	7.0
Haga	7.3	7.2	7.5	6.5	6.8	6.5	7.0
Adelphi	7.0	6.7	6.2	6.8	7.5	7.5	6.9
PST-CB1	7.0	7.2	6.0	7.0	6.8	7.5	6.9
Asset	7.3	6.8	7.3	6.2	6.2	7.5	6.9
America	6.2	6.5	7.3	7.0	7.0	7.2	6.9
Ba 69-82	7.0	7.3	7.2	6.2	6.8	6.7	6.9
Tendos	6.2	6.0	7.3	7.2	7.2	7.3	6.9
Sydsport	6.5	6.7	7.0	6.7	7.5	6.5	6.8
Nassau	6.3	6.2	6.3	6.8	7.5	7.5	6.8
Ba 72-441	6.7	7.5	7.8	6.7	6.5	5.5	6.8
K3-178	6.7	7.2	5.5	6.7	6.8	7.5	6.7
Touchdown	6.5	6.2	6.0	7.2	7.5	7.0	6.7
Trenton	6.7	6.8	7.2	6.8	7.3	5.5	6.7
Ba 70-139	6.8	7.0	7.2	7.3	6.7	5.3	6.7
WW Ag 491	6.7	7.0	7.2	7.0	6.2	6.2	6.7
F-1872	7.0	6.7	6.0	6.2	7.3	7.0	6.7
Merion	6.8	7.0	6.8	7.2	6.5	5.7	6.7
Parade	6.0	6.7	6.3	7.0	7.2	6.8	6.7
Cheri	6.7	7.0	7.5	6.7	6.8	5.3	6.7
NE 80-88	6.8	6.2	6.0	6.8	7.0	7.0	6.6
Rugby	7.0	6.8	6.8	6.5	6.8	5.8	6.6
239	6.7	7.5	6.2	6.3	6.8	6.2	6.6
K1-152	7.2	7.0	6.0	6.2	6.8	6.5	6.6
Gnome	7.0	6.2	7.0	6.7	6.0	6.7	6.6
Aquila	6.7	6.3	6.0	7.0	7.0	6.2	6.5
Somerset	6.3	6.3	7.0	6.7	6.8	6.0	6.5
Ba 70-242	6.5	7.7	6.5	6.3	6.7	5.3	6.5
Julia	6.8	7.2	7.0	6.7	5.3	6.0	6.5
Ba 72-492	6.7	6.8	7.2	7.0	6.7	4.7	6.5
Parade	6.3	6.5	6.0	6.0	7.3	6.8	6.5
Victa	6.3	6.5	7.5	6.3	6.3	5.5	6.4



TABLE 5 cont'. 1986 National Kentucky Bluegrass Trial

<u>Variety</u>	<u>Quality Ratings</u>						<u>Grand Means</u>
	<u>4/25</u>	<u>5/20</u>	<u>6/23</u>	<u>8/2</u>	<u>9/6</u>	<u>10/27</u>	
Merit	6.3	7.5	6.8	6.0	6.2	5.5	6.4
WW Ag 496	6.7	6.0	6.8	6.7	5.3	6.3	6.3
Monopoly	6.3	6.8	6.7	6.2	5.3	6.3	6.3
WW Ag 468	5.7	7.0	6.3	6.7	5.8	5.8	6.2
Ikone	7.0	6.7	5.7	6.7	6.0	5.0	6.2
Barzan	6.3	5.0	6.8	6.2	6.2	6.5	6.2
Welcome	6.0	6.7	6.8	7.0	5.3	4.7	6.1
Compact	5.8	6.2	6.7	6.3	5.7	5.8	6.1
Baron	6.0	6.0	6.7	6.2	6.5	5.0	6.1
Harmony	5.8	6.8	6.3	6.0	6.0	5.3	6.1
Connie	5.8	6.0	6.8	6.7	5.3	5.3	6.0
Wabash	6.3	4.7	5.7	6.3	6.0	7.0	6.0
WW Ag 495	5.8	6.0	5.3	6.5	6.5	5.8	6.0
Huntsville	6.8	5.0	5.3	6.3	6.2	6.3	6.0
Majestic	6.3	4.8	5.8	6.7	6.0	6.3	6.0
HV 97	6.0	6.8	6.3	5.8	4.7	6.2	6.0
Ba 73-626	6.5	6.0	6.3	6.7	6.0	4.3	6.0
Bar VB 534	5.8	5.0	5.7	6.7	7.0	5.5	5.9
Cynthia	5.3	5.0	6.0	6.7	7.2	5.3	5.9
A-34	6.0	6.0	6.3	6.3	5.3	5.3	5.9
Amazon	5.7	5.5	6.5	6.5	5.2	5.3	5.8
Kenblue	5.3	4.0	5.0	6.2	5.0	6.0	5.3
Joy	5.8	3.3	5.7	5.7	5.3	5.7	5.3
Annika	5.5	4.7	6.0	5.3	5.0	3.0	4.9
S. Dakota Certified	5.7	3.0	3.7	6.0	5.0	5.3	4.8

TABLE 6. Embark plus Iron sources. Effects on turf quality and seedhead control.

<u>Embark @ 1/8 lb/A plus</u>	Rate of Iron Carrier	<u>Quality Ratings</u>								<u>% Seedhead Density</u>	<u>% Seedhead Control</u>
		<u>% Fe</u>	<u>5/12</u>	<u>5/17</u>	<u>5/20</u>	<u>5/24</u>	<u>5/27</u>	<u>5/31</u>	<u>6/3</u>		
Scott's Iron S	1.5 lbs/M	5	9.0	8.5	9.0	9.0	9.0	9.0	9.0	8.3	90.6
Chipco Microgreen	2 oz/M	4	8.3	8.3	8.3	8.5	8.7	8.5	8.7	41.6	52.9
Ferromec	2 oz/M	6	8.8	8.5	8.5	8.7	8.7	8.5	8.3	40.0	54.7
	4 oz/M	6	8.8	8.5	8.5	8.7	9.0	8.2	7.8	66.7	24.5
	6 oz/M	6	8.8	8.3	8.5	8.5	8.7	8.3	9.0	60.0	32.0
Ferromec AC	2 oz/M	6	8.3	8.2	8.0	8.8	8.3	8.5	9.0	13.7	84.6
	4 oz/M	6	8.5	8.3	8.5	8.7	8.7	8.8	8.6	36.7	58.6
	6 oz/M	6	8.5	8.3	8.2	8.7	8.7	8.8	8.5	25.0	71.7
Lesco + Iron	3 oz/M	5	8.5	8.0	8.5	8.5	8.5	8.7	8.8	26.7	70.0
Agriplex	0.5 oz/M	5	8.7	8.0	7.8	8.7	8.3	8.8	9.0	8.3	90.6
	1.5 oz/M	5	8.5	8.2	7.6	8.5	8.5	9	8.8	3.7	96.0
FeSO <sub>4</sub>	3.6 oz/M	20	8.5	8.3	8.2	8.5	8.3	8.3	8.5	41.7	52.9
Embark alone		-	8.0	7.8	7.5	8.0	7.8	8.2	8.3	5.3	94.0
Control		-	<u>8.5</u>	<u>8.3</u>	<u>8.2</u>	<u>8.7</u>	<u>8.2</u>	<u>7.7</u>	<u>7.0</u>	<u>88.3</u>	<u>0</u>
LSD (P=0.05)			0.8	0.6	0.6	0.6	0.6	0.7	1.2	30.9	

For highest quality turf with effective seedhead control, we currently recommend Scott's Iron S plus Embark. Agriplex does not antagonize the seedhead control from Embark but does not give as good of a masking effect as does the Scott's product.

Data in table 7 shows the results of a study of Prograss and plant growth regulators to control annual bluegrass in fairway turf. This study is being conducted at six golf courses in Michigan: Blythefield Country Club and Kent Country Club, both in Grand Rapids; Walnut Hills Country Club in East Lansing; Orchard Lake Country Club in Orchard Lake; Bloomfield Hills Country Club in Bloomfield Hills; and Barton Hills Country Club in Ann Arbor. The study was initiated in August of 1987 and will be conducted for at least one more year. The PGR treatments (Cutless and Scott's TGR) are applied twice a year (April and August) while Prograss is applied twice in the fall (September and October). Examining the data shows that none of the treatments have caused much change in the amount of annual bluegrass when compared to the control. However, differences between golf courses is significant implying that management factors may be more important than the currently available chemicals. It will be very interesting to follow this study over the next year.

Many superintendents are trying to convert their fairways from annual bluegrass to creeping bentgrass. The quickest and most popular method to convert fairways is use Roundup and then reseed with creeping bentgrass. However, significant amounts of annual bluegrass germinate along with the bentgrass with the resulting fairways containing only 40 to 80 percent bentgrass. Prograss, a selective annual bluegrass herbicide, has been used after seeding the bentgrass to control the germinating annual bluegrass. Rates and timing of the Prograss application are extremely important because the seedling bentgrass can be severely injured by Prograss. We investigated three Prograss rates applied at either 4 + 8 weeks after bentgrass germination (WAG) or at 6 + 10 WAG. The Prograss was applied to plots that were seeded with either 'Penncross' creeping bentgrass or annual bluegrass. Data in table 8 shows the result for creeping bentgrass and annual bluegrass, respectively. If Prograss applications are delayed to 6 + 10 WAG then the bentgrass is less injured and establishes a little more quickly. However, the annual bluegrass is also less affected with the net result being more annual bluegrass. Also, the 3/8 plus 3/4 lb AI/A rate seems to be the best since it does not injure the bentgrass as severely as the other rates tested. The 4 + 8 WAG applications are more effective for controlling annual bluegrass and should be used in situations where less than ideal playing conditions can be tolerated the following spring. Where maximum grass cover is desired the 6 + 10 WAG treatment schedule will give the best results although more annual bluegrass will be present. Thus, a trade off between turf quality and annual bluegrass quantity must be decided. The least annual bluegrass will be found where early (4 + 8 WAG) applications are made but the turf will be more injured and it will take a longer period of time for the bentgrass to establish and give uniform cover. Faster establishment will occur by waiting longer after germination to apply the Prograss (6 + 10 WAG) but more annual bluegrass will remain in the turf.