

Effect of Mercury Compounds and Arsenate of Lead on Soil Nitrification

By J. W. WHITE, *Professor of Soils*

Pennsylvania Agricultural Experiment Station

Read at the 5th Annual Educational Conference of the National Association of Greenkeepers of America, held at Columbus, O., February 3-6.

A STUDY of the bulletins of the United States Golf Association Green Section concerning recommendations for the development and maintenance of turf grasses brings out some interesting facts worthy of consideration at this time. The purpose of my paper, therefore, is to give you my impressions of certain recommendations found in those bulletins in addition to a discussion of the topic assigned to me. I trust that what I may say will not be taken as destructive criticism, but rather as helpful suggestions that may lead to a better understanding of the fundamental principles underlying the art of greenkeeping and fairway management.

After a study of the bulletins in preparation for our first Greenkeepers' Conference at Penn State in 1929, I resolved to face the issues squarely even though my ideas may be entirely contrary to those previously presented. With your permission, therefore, I wish to review in general the recommendations given at our Penn State Conferences in 1929 and 1930.

The exacting demands of the green committees concerning the development of a perfect turf, free from weeds, lead to recommendations entirely contrary to the fundamental principles of soil fertility and these have led to disaster in many instances, especially in our Eastern states. The continuous and excessive use of sulphate of ammonia and the ban on lime and phosphorus have been the cause of most of your trials and tribulations. The evils of an extremely acid soil are too well known to the soil chemist to warrant its development as a means of weed control for a soil too acid to grow weeds cannot support a normal growth of turf grasses.

CONTROL WEEDS THROUGH VIGOROUS TURF

THE most logical means of weed control is through the development of a vigorous turf which leaves no room for weed invasion. The use of some form of phosphatic fertilizer and lime in moderation should under proper management attain this end. There is no logical reason to believe that golf grasses require fundamentally different treatment than is accorded those found in lawns and well-kept pastures. Grasses grown under very acid conditions and abnormally stimulated in growth by too frequent applications of ammonium sulphate are more susceptible to fungus disease and other injuries than is true of grasses grown under normal conditions. By that I mean soil conditions where the plant has the opportunity to assimilate plant food in accordance with the fundamental principles of plant nutrition.

It is our recommendation in Pennsylvania that the soil reaction should be maintained approximately between ph. 5.8 and 6.5 and that superphosphate should be applied each year and also that if necessary limestone should be applied at intervals to maintain the proper control of excessive soil acidity.

The amounts of phosphorus, limestone and nitrogen to be applied naturally vary with soil conditions. An annual application of seven pounds of superphosphate per 1,000 square feet should be sufficient under average conditions.

If sulphate of ammonia is used as the source of nitrogen, limestone should be applied at the rate of 75 pounds for each 100 pounds of ammonium sulphate applied. In other words, an annual application of 11.5 pounds of ammonium sulphate per 1,000 square feet requires the use of 8.6 pounds of limestone.



PROFESSOR J. W. WHITE
Penn State College

TOXIC SOIL CAUSES GRASS FAILURE


AT the time of the Penn State Conference it was pointed out that in many instances the failure of turf grasses during the summer months was due to the toxic effect of soil acidity and not to the invasion of fungus diseases. This was proven by the fact that mercury compounds even after repeated applications failed to restore the turf to normal color and vigor. However, an application of lime restored the grasses to normal condition. The evils of soil acidity brought about by the excessive use of ammonium sulphate were first noted on poorly-drained greens where there occurred an accumulation of salts.

Where it becomes necessary to use lime during the growing season it is recommended that limestone be used rather than hydrated lime since the latter has a tendency to cake and discolor the green while limestone particles being heavier should work in between the grasses and entirely disappear from view.

The statement that certain grasses have the ability to utilize nitrogen in the form of ammonia has not been entirely accepted in this country. In accordance with our modern conception of plant nutrition, most of our economic grasses depend upon soil nitrates as their source of nitrogen. The ability of certain grasses such as those of the agrostis family, which includes the bents, to grow under very acid soil conditions is attributed to their ability to utilize ammonia instead of nitrates. The speaker believes that the relative resistance to acidity of such grasses as those included in the agrostis family is due primarily to the fact that they are strong feeders of calcium. That is, they have the ability to utilize small amounts of calcium in forms and amounts unavailable to other grasses such as the Poa family which includes Kentucky blue grass.

UNDER sold conditions such as found on greens, fairways, pastures and lawns there exists at all times during the growing season a scarcity of soil nitrates; in other words, nitrates become a limiting factor in plant growth. This is due to several causes:

(1) Under sod there exists keen competition between the grasses and soil microorganisms for the available nitrogen present. The soil microorganisms use as their source of food or energy the old roots of the grasses which are high in carbon. These soil organisms require nitrogen in a definite propor-



Thousands of Dollars Invested in Greens . . . A Few Dollars Will Save Them From Destruction by Deadly Brown Patch

A carpet of green velvet—a perfect putting surface—that is what makes players proud of their golf club. It takes time and money to get turf into such perfect condition.

Then comes the deadly Brown Patch and works havoc. A very small investment will save all this.

Barbak 211—the perfect disinfectant will keep your greens free from Brown Patch. Tests of treated and untreated areas prove that Brown Patch will travel over an untreated area and come to a dead stop on the line precisely where Barbak 211 has been used.

As a preventative, Barbak 211 protects against Brown Patch for a longer period than ordinary disinfectants. One well-known greenskeeper places this at 42 days. As a cure, it checks the fungus immediately and quickly brings back a normal stand of turf.

Barbak 211 may be applied in solution or dry, mixed with a top dressing, sand or fertilizer and watered in. Write for further facts.



AMERICAN CYANAMID SALES CO., INC.
535 Fifth Avenue New York City

Barbak

TWO • 211 • ELEVEN

TURF DISINFECTANT

“Milorganize”

Your Course
with



Analysis:

NITROGEN (equivalent to ammonia)	6.0 - 6.5%
PHOSPHORIC ACID (P_2O_5)	2-5 - 3-0%
POTASH (K_2O)	0-25 - 0.5%
MOISTURE	less than 5.0%

The Nitrogen in Milorganite
is

**ALL ORGANIC
WATER INSOLUBLE
HIGHLY AVAILABLE**

Perfect Mechanical Condition
Best Carrier for Lead Arsenate

OUR SERVICE BUREAU and
LABORATORIES are at your Service.

*Our trade-mark on each bag is your
guarantee of high-grade material.*

**Milwaukee Sewerage
Commission**

Milwaukee Wisconsin

tion to the carbon assimilated in order to balance the composition of their bodies which, like the higher plants, contain definite ratios of carbon to nitrogen.

(2) Under condition of sod, especially in case of greens frequently rolled, nitrification is reduced to poor aeration; that is, the supply of oxygen in the soil atmosphere is insufficient to meet the demands of the nitrifying organisms.

(3) In case of extreme soil acidity the activity of the nitrifying organisms is considerably reduced. The optimum reaction for soil nitrification is between ph. 6.0 and 7.5.

(4) The presence of an excess of ammonium salts brought about as the result of poor drainage where large amounts of ammonium sulphate have been applied has been found to reduce nitrification. There are many other soil factors which may control nitrification such as excess of water, low temperature, and the presence of an excess of soluble organic matter and certain mineral salts. Of the heavier metals mercury and silver have the most pronounced effect in reducing the activity of nitrifying organisms. The effect of these metals is to precipitate the protoplasts or destroy the cell structure of soil microorganisms.

TREATMENT OF SOILS WITH MERCURY COMPOUNDS

IN order to gain further information concerning the controlling influence of mercury compounds on the activity of nitrifying organisms, the speaker applied to both an acid and neutral soil, mercury compounds commonly used for the control of brown patch and other fungus diseases of turf grasses. The soil used was taken from the ammonium sulphate plot of the Pennsylvania field experiments which had received a total of 2740 pounds of ammonium sulphate distributed uniformly over a period of 47 years. The soil has a ph. of 4.77. An equal portion of the same soil was treated with limestone in amounts sufficient to produce a neutral soil of ph. 7.0.

The field plot soil in addition to ammonium sulphate had received liberal applications of superphosphate and muriate of potash during the period of the field experiment. In addition to the mercury compounds, lead arsenate used for the control of grubs, was also included in the nitrification studies. One-half of each soil was treated with ammonium sulphate at the rate of 226 pounds per acre or 5.2

EFFECT OF MERCURY COMPOUNDS AND ARSENATE OF LEAD ON SOIL
NITRIFICATION

Results expressed as parts per million of nitric nitrogen produced in 14 days.

Materials used and rate per 1,000 sq. ft. of soil	Unlimed Soil pH 4.77		Unlimed Soil pH 7.02		Totals	
	No Amm.		No Amm.		No	
	Sul.	Sul.	Sul.	Sul.	Lime	Limed
Untreated Soil	55	43	58	41	98	99
1/2 lb. Bichloride of Mercury	41	25	61	38	66	99
2 lbs. Bichloride of Mercury	44	26	61	41	70	102
1/2 lb. Calomel	46	30	52	38	76	90
2 lbs. Calomel	46	24	58	44	70	102
1 lb. Semesan	55	34	56	37	89	93
3 lbs. Semesan	45	28	60	38	73	98
11 1/2 lbs. Arsenate of Lead	59	45	62	40	104	102
46 lbs. Arsenate of Lead	94	4	66	38	139	104

pounds per 1,000 square feet of soil. The experiment was conducted for a period of two weeks. The accompanying summary shows the rate of application of mercury and arsenic compounds and the total nitric nitrogen produced in case of each treatment.

A study of the results shown in the accompanying table brings out the fact that on the unlimed soil, parallel to the condition found on many greens and fairways of our eastern golf courses, mercury has considerably reduced the activity of the nitrifying organisms. Arsenate of lead, on the other hand,

GET BUSY! Spring is Here


Let America's *lowest-priced* golf course tractor do *all* your work—quickly—economically. The Staude Utility Golf Course Tractor is speedy—light—powerful—easy to operate—and its repairs amount to almost nothing.

COMPLETE

with
Model A
Ford
Chassis

\$655

F. O. B. ST. PAUL



COMPLETE

with
Model T
Ford
Chassis

\$475

F. O. B. ST. PAUL

Staude Utility Golf Course Tractor

Manufactured Continuously Since 1916

Nearly 2000 golf clubs are using the Staude to tremendous advantage. This fact is its greatest recommendation. Its service is completely out of proportion to its low cost.

The Staude is thoroughly dependable. Long life and satisfactory service are built right into it. It will unfailingly answer all your requirements. Investigate the Staude before deciding upon any type of power equipment.

Sold on a Money-Back Guarantee through your local Ford dealer, golf supply house or direct from us

Write for Full Particulars and Easy Payment Plan.

E. G. STAUDE MAK-A-TRACTOR CO.

2630 University Ave., St. Paul, Minn.

THE ROYER COMPOST MIXER

*Saves in
Labor Costs*



**Read what the Chief of Parks of
the National Capital Says About
His Two ROYERS :**

PUBLIC BUILDINGS AND PUBLIC PARKS
OF THE NATIONAL CAPITAL

Washington, D. C.
February 9, 1931.

Dear Sirs:

Your communication of January 28 in reference to the two Royer Mixers which this office purchased from your company has just been received.

We first purchased a small model which proved so satisfactory that we decided to purchase one of the larger type mixers. The small mixer is being used permanently by the propagating gardens and greenhouses and the larger machine in the preparation of compost for lawn areas throughout the park system.

We have found that the Royer Mixers have *saved us considerable in labor costs* as it has eliminated all of the old hand-mixing methods.

F. T. GARTSIDE,
Chief, Park Division.

Write us for full particulars

Sold through leading golf equipment dealers

Royer Foundry & Machine Co.
158 Pringle St. Kingston Station Wilkes-Barre, Pa.

has stimulated the production of soil nitrates in the cases of each soil. The addition of lime, however, has eliminated the injurious effect of the mercury compounds as shown from the fact that on the unlimed soil the three mercury compounds reduced nitrification on an average of 25 per cent compared to only 1.5 per cent on the soil which had received limestone.

The reduction of nitrification in case of each of the three mercury compounds used was as follows: Bichloride 31 per cent, calomel 26 per cent, and Semesan 17 per cent. Arsenate of lead increased nitrification on the unlimed soil 40 per cent and on the limed soil 5 per cent.

CONTROL OF FUNGUS DISEASES MUST BE CAREFULLY HANDLED

IN addition to the factors already mentioned which may reduce the supply of available nitrates in the soil, we now learn that mercury compounds used for the control of fungus diseases also produce an injurious effect on nitrifying organisms in acid soils. The addition of lime, however, has been shown to overcome this toxic effect. The results of this experiment are presented to you with no idea in mind of discouraging the use of compounds of mercury necessary to combat fungus diseases of turf grasses, but rather to call to your attention the fact that such materials may reduce the available nitrogen of the soil especially in case of those soils which are in need of lime.

The slow recovery of grasses, often noted, following repeated application of fungicides, may be due to the reduction in the supply of available nitrogen in the soil as the result of injury to nitrifying organisms.

In conclusion may I say that today we are facing a new era in the scientific management of greens and fairways. The greenkeepers are turning to their state agricultural experiment stations for advice, gained as the result of many years of careful research. The soil chemist, the pathologist and the plant breeder are joining hands in an effort to aid those responsible for the development and maintenance of our American golf greens and fairways.

The results of various turf grass experiments initiated by the United States Department of Agriculture are beginning to bear fruit. Art and science, theory and practice are going forward hand in hand and there should be no conflict between.