

A Bentgrass Breeding Consortium to Support the Golf Industry

Michael Casler
University of Wisconsin

Andrew Hamblin
University of Illinois

Suleiman Bughrara
Michigan State University

Objectives:

1. To develop elite clones of creeping bentgrass with multiple pest resistances and stress tolerances that can be delivered to the seed industry for use in synthesizing new creeping bentgrass cultivars broadly adapted to a range of ecological and environmental conditions including reduced pesticide application.

Start Date: 2003

Project Duration: three years

Total Funding: \$170,817
(three institutions)

Creeping bentgrass (*Agrostis palustris*) is the premier grass for golf course putting greens and is one of the most desirable grasses for fairways and tee boxes for many of the nation's golf courses. Recent breeding advances demonstrated that genetic variation exists within creeping bentgrass for a range of pest resistances and stress tolerances. Many of these traits allow bentgrass to be grown in environments and under conditions that were impossible just a few years ago.

For many golf courses, maintenance of high quality turf requires frequent, varied, and intensive pesticide applications. Pesticide costs can consume up to 10% of the total budget for a high-management golf course. Intensive management (including frequent and low mowing, irrigation) and heavy play serves to enhance and/or spread the development of pest problems, particularly fungal diseases.

Genetic resistance to disease and other pests is a widespread phenomenon in agricultural and horticultural plants. Disease resistance has been used to protect economically important plants for over 90



Creeping bentgrass plants immediately after transplanting into a ryegrass fairway sod at Madison, WI. These plants will fill in the empty circles by early summer, 2004 and be inoculated with *Pythium* spp..



The "chamber of death" that will be used to provide an optimum environment for expression of *Pythium* spp.

years. While there has been much research on genetics and breeding for individual pests of creeping bentgrass, there has been no concerted effort to develop multiple pest-resistant germplasm.

Three populations have been developed for this study. At the University of Wisconsin, crosses have been made between creeping bentgrass clones selected for resistance to grey and speckled snow mold fungi. At Michigan State University and the University of Illinois, crosses have been made among clones selected for turf quality from a large number of golf courses. All clones were selected from old golf courses where there have been many years of natural selection for survival under close-mowing conditions.

From crosses made at each location, 350 plants are being chosen for this study, for a total of 1,050 creeping bentgrass plants. Each plant is being vegetatively cloned in greenhouses. Researchers met in Chicago in February, 2003, and exchanged clonal material, each contributing 200 clones from our respective collection and breeding programs. They each increased the 600 clones using vegetative propagation techniques in the greenhouse during the remainder of the winter and

early summer.

Five field experiments were planted with four replicates each: one at Urbana, IL; one at East Lansing, MI; two at Madison, WI; and one at Land 'O Lakes, WI. These experiments will be inoculated and scored beginning in 2004. Plants were transplanted into perennial ryegrass fairways and topdressed for rapid establishment and sod formation.

Summary Points

- Researchers met in Chicago in February, 2003, and exchanged clonal material, each contributing 200 clones from their respective collection and breeding programs. Researchers each increased the 600 clones using vegetative propagation techniques in the greenhouse during the remainder of the winter and early summer.

- Five field experiments were planted with four replicates each: one at Urbana, IL; one at East Lansing, MI; two at Madison, WI; and one at Land 'O Lakes, WI. These experiments will be inoculated and scored beginning in 2004. Plants were transplanted into perennial ryegrass fairways and topdressed for rapid establishment and sod formation.