

Don't forget the obvious in that spreader patterns are also important when applying fertilizer. Be sure to calibrate the spreader and your overlap – with a rotary spreader is 50% where the edge of your pattern should touch the spreader tracks from the previous pattern. Be careful with potassium as it is overrated – only apply if a soil test indicates you need it.

Soccer fields with considerable wear can handle three fall applications: after the heat in August, and 1 lb. in September and October. The risk with this fertilizing strategy is that it will increase snow mold, but this risk is normally justified on areas under severe wear pressure. Slow release should be used for the first two applications and fast release for the last one.

Reduce your fertilization rates and you have to reduce your expectations. Your best bet is to let the grass plants dictate fertilization. If you see clover, dollar spot, red thread or rust in your fields, you have too little nitrogen. If you have perennial brown patch, pythium, snow mold and/or *Poa annua* problems, your fertility might be too high. You can't make up for missed fall applications in spring. Furthermore, leaching is not as much of a problem in fall because plants take up the fertilizer immediately – but that doesn't give us license to over-apply or apply carelessly.

Fall Seeding

Though Kentucky bluegrass is the preferred grass in lawns and athletic fields, it's not very competitive as a seedling. Perennial ryegrass is easier to overseed because of its tremendous competitiveness as a seedling, but it is more sensitive to diseases and does not spread like Kentucky bluegrass. When overseeding thinned out grasses, Kentucky bluegrass will work fine if it is a very thin turf and the area will not be trafficked for six growing weeks or more. But if the turf is "in play" and is needed constantly, use perennial ryegrass. Though many people use sod for a quick fix during a playing season, newly laid sod doesn't handle traffic well and overseeding may be a better choice.

The best time to seed is August, but don't seed any later than mid-September. If time restraints are a problem, then do it

in September. A rule of thumb is if the field is seeded the first of September under normal conditions, it will be available for play by the first week of June. Dormant seeding works very well combined with aggressive aerification at the end of the year.

Seeding at the wrong time of year is always a problem with weed control. Most herbicides have some kind of restriction when it comes to seedlings. Whether you spray over the top or prior to seeding, make sure you read the label carefully because every herbicide differs in its restrictions with use around seedlings. The best way to control weeds in seedlings is through mowing and most people don't mow a new stand soon enough. In order to maximize establishment, good seed to soil contact is required. The application of high phosphorous fertilizers prior to

seeding will also help establishment.

Finally, Dr. Reicher does not recommend perennial rye when establishing new fields. He recommends 100% Kentucky bluegrass blends instead. He maintains recommended seeding rates are too high. Regardless of the seeding rate, a turf stand can only support a given number of seedlings.

In terms of fall broadleaf weed control, Dr. Reicher referred to a Michigan State University study showing you can still get great weed control in the late fall as long as the target weeds are green and photosynthesizing. They will take up the herbicide but will show little effect only to die during the winter. Where aerification is concerned, you cannot kill turf or over aerify provided you are not doing it at 90F. Lastly, he recommended keeping mowing heights the same year round. ■

Industry News • STA New Members

Executive Director Appointed

Ken Cousineau, CAE, has been selected to succeed Vince Gillis, CAE, as Executive Director of the Canadian Golf Superintendents Association. Mr. Cousineau, who assumed his new duties February 11th, brings to the CGSA strong association management experience, both in managing internal administrative functions and external relationships with government and stakeholders.

Breeding and Evaluation of Turfgrasses Technical Conference

May 1-2, 2002. Organized by the Sports Turf Research



Institute, the conference, being held in West Yorkshire, will interest all turfgrass breeders and evaluators, turf scientists and other turf professionals.

A panel of 8 speakers will present 10 conference sessions. Further details can be obtained from Anne Wilson, Head of External Affairs, STRI,

telephone 01274 565131, email info@stri.co.uk.

TPI Confirms England for 2003 Winter Conference

For the first time in its 35-year history, the Mid-Winter Conference and Exhibition of Turfgrass Producers International will be held outside of North America. The 2003 meeting will occur from February 26-28 in Birmingham, England. Open to everyone in the green industry, anyone interested in receiving information about exhibiting or attending should contact TPI at 800-405-8873 or 847-705-9898, email can be sent to info@TurfGrassSod.org.

Welcome New STA Members

- Jim Nelson, City of Kitchener, ON
- Tim Ernst, City of Kitchener, ON
- Jorg Hermanns, Hermanns Contracting Ltd., Schomberg, ON
- Laurie Weatherall, City of Ottawa, ON
- Bruce Warren, Parc Downsview Park, Toronto, ON