

1. Location & Growing Period

Geographic location determines the number of months for active growth and aids in the selection of a suitable turfgrass species. Growing months for turf can differ by several months between various locations. Therefore, the total amounts of nitrogen to be applied per year must be adjusted for location. For instance, a bermudagrass sports field in Tennessee may only need 6-9 lb of nitrogen (per 1,000 ft²) per year compared to the exact same field in Florida that needs more than 9 lb per year, with the difference being the length of the growing season.

2. Root Zone Composition

The make up of the root zone also affects the amounts of nitrogen to be applied. Fields consisting of high silt and clay contents require different application procedures versus a sand-based field. Root zones consisting predominantly of silt and clay have lower percolation rates, helping to prevent the loss of nitrogen through leaching. Therefore, applications of nitrogen can be limited to a monthly basis using a fertilizer consisting of both fast and slow release nitrogen sources.

Sand-based root zones require applications to be applied more often with less total nitrogen per application. These root zones promote drainage, which reduces the holding capacity for nutrients like nitrogen. To ensure the availability of nitrogen for the plant, it should be applied every 10 to 14 days at half the normal rate depending on irrigation and precipitation levels. Sand-based fields have other nutrient retention problems as well. The lack of cation exchange capacity of sand allows other nutrients such as potassium, which

Table 1. Nitrogen applications (lb/1,000 ft²) for bermudagrass fields, southern US. Note that May through October are actively growing turf months and Jan-April and Nov-Dec are overseeding periods.

*Total amount of nitrogen/year

Month	Sand-Based Fields		Native Soil Fields	
	Slow Release	Fast Release	Slow Release	Fast Release
January	-	-	-	-
February	-	-	-	-
March	1	0.5-1	-	0.5
April	1	1	1	1
May	2	1	2	1
June	-	1	-	1
July	2	1	2	1
August	-	1	-	1
September	2	1	1	1
October	-	1	-	-
November	1	1	1	0.5
December	-	-	-	-
Total* (no overseed)	6	6	5	5
Total* (overseed)	9	9	7	7

normally is found at acceptable levels in native soils, to potentially leach out (Turgeon, 1996). Therefore, sand-based athletic fields should receive applications of potassium and phosphorus, as well as other micronutrients, more frequently than native soil athletic fields. As stated earlier, soil testing will also help determine these nutrient requirements.

3. Turf Quality

Another way to determine nitrogen needs is by assessing overall turf quality. Turfgrass growing under low nitrogen levels will exhibit chlorosis. Chlorotic plants appear yellowish-green to yellow (Emmons,

1995). However, yellowing turf does not necessarily mean that levels of nitrogen are inadequate, but proper nitrogen fertility will correct any deficiencies. Other environmental stresses can produce the same effects. Density is another turfgrass quality that can be used as an indicator for nitrogen deficiencies. Often, turfgrass areas infested with weeds can indicate a lack of nitrogen available to the plant (Emmons, 1995). Weeds are not the cause of bad turf. Rather, weeds are caused by bad turf. Low nitrogen fertility reduces the competitive nature of the turf, which allows invasive weeds to take over.

4. Species Selection

Turfgrass species selection also affects the amounts of nitrogen needed. Bermudagrass, Kentucky bluegrass and perennial ryegrass generally require more input of nitrogen per year than any other turfgrasses being used on athletic fields. These grasses are vigorous and aggressively growing plants that require high nitrogen fertility. Increased rates of nitrogen must be applied in order to keep the plant healthy and able to recuperate from wear. Bermudagrass can receive rates of nitrogen per 1,000 ft² of 6-15 lb per year depending on geographic location and field usage.



Nitrogen differences on Kentucky bluegrass.

Table 2. Nitrogen applications (lb/1,000 ft²) for Kentucky bluegrass/perennial ryegrass athletic fields. Note that June-July applications are optional, slow release fertilizer recommended.

Month	Sand-Based Fields		Native Soil Fields	
	Slow Release	Fast Release	Slow Release	Fast Release
January	-	-	-	-
February	-	-	-	-
March	-	-	-	-
April	0.5-1	1	1	1
May	1	0.5-1	-	0.5
June	1	0.5	1	0.5
July	-	0.5	0.5	-
August	0.5	0.5	-	0.5-1
September	1	1	1-1.5	1
October	-	-	-	-
November	-	-	-	-
December	-	-	-	-
Total N per year	4-4.5	4-4.5	3.5-4	3.5-4

Kentucky bluegrass can receive rates ranging from 3-6 lb per year.

5. Type of Venue

Quality requirements for sports fields differ between little league parks and professional stadiums. Typically, city operated fields will not be mowed as many times or as low as in professional stadiums. This difference changes nitrogen needs. Lower mowing frequency and higher mowing heights requires less nitrogen input. Lastly, fields that collect clippings will need to have more nitrogen applied than fields that mulch clippings.

General Rules

Despite all of the conditions described above, there are some generalized rules for producing a fertility program that is right for you. First, nitrogen should be applied at least one time per active growing month. Amounts of nitrogen will differ, but applications should be made every growing month to ensure sufficient amounts. Figures 1 and 2 describe application timing of fast and slow release fertilizers on Kentucky bluegrass, perennial ryegrass and bermudagrass. Highly used fields should receive one pound of nitrogen per

month of active growth while low use fields will only need as little as half a pound of nitrogen.

The more applications made per month the better. Try splitting applications in half every 14 days. Applying fertilizers more frequently aids in keeping nitrogen available to the plant at all times (Calhoun *et al.*, 2002). Use fertilizers with both fast release and slow release nitrogen forms. One type of fertilizer is not sufficient for an entire season. Fertilizers with different nitrogen forms and percentages should be used to maximize growth (Puhalla *et al.*, 1999).

Recommendations of applying phosphorous and potassium by a soil analysis report should be followed. However, some turfgrass managers apply potassium at a one-to-one rate with nitrogen (Emmons, 1995). This is significant to managers with sand-based root zones. Potassium aids in stress tolerance of the plant, but is readily leached from sand root zones (Turgeon, 1996). Finally, applications of nitrogen and potassium should be given at the end of each growing season when shoot growth slows. During this time, the plant is storing carbohydrates, rebuilding damaged roots, and preparing for harsh environmental conditions (Emmons, 1995). For some sports field managers, nitrogen applications do not end with the induction of dormancy at the end of the growing season, but continue with the overseeding of ryegrass for play in the winter season. Fields overseeded with ryegrass need to be fertilized continually throughout the cool season growing months.

Sports Turf Association's
Sports Turf Management & Maintenance Course

1/3 required courses for the Parks & Open Space Professional Training Program (Level 1)

May 2-5, 2011, University of Guelph
For more information & registration: 519-763-9431
www.sportsturfassociation.com/POSA/Prof Training Program



To learn more about STA advertising and promotional opportunities, contact Lee at the office.




99 John St. North, Box 171
Harriston, Ontario
N0G 1Z0
Phone (519) 338-3840
Fax (519) 338-2510
Email spearese@wightman.ca

SUPPLIERS OF PREMIUM
TURF SEED PRODUCTS

Tables 1 and 2 describe examples of fertility programs for sports fields. These examples are meant to be modified and adapted to fit the needs of individual fields. Table 1 describes nitrogen applications for bermudagrass fields in the southern United States. Amounts of nitrogen in pounds per 1,000 ft² are given in terms of slow release and fast release fertilizers in either native soil or sand-based athletic fields. In addition, Table 1 describes a continuance of the fertility program for overseeded turf. Table 2 describes applications for fields with Kentucky bluegrass and perennial ryegrass. Nitrogen amounts are also given in pounds per 1,000 ft². Application


Figure 1. Cool season fertility fast and slow release timing.



EDITOR'S NOTE

The rates of application of nutrients provided are somewhat higher than those for Canadian conditions, however the principles remain the same. Sports turf managers should be aware of these differences in the rates of nutrient application to different species in the regions of North America.

Figure 2. Warm season fertility fast and slow release timing.



amounts are given for both fast and slow release, as well as for native soil versus sand-based athletic fields.

Guidelines for creating a fertility program are useful, yet, keep in mind that they are only guidelines. Each individual field requires its own specific fertility program based upon the needs of the sports turf managers, players, owners and others who enjoy the field. Finding what works for you is not an easy task, so be patient and do not be afraid to try new things.

References

- 1) Calhoun, Ronald, Lisa Sorochan, John Sorochan, John Rogers III and James Crum. 2002. "Optimizing Cultural Practices to Improve Athletic Field Performance." Bulletin E18TURF, Michigan State University Extension.
- 2) Emmons, Robert. Turfgrass Science and Management. Delmar: 1995, 176-194.
- 3) Puhalla, Jim, Jeff Krans, and Mike Goatley. Sports Fields. Ann Arbor: 1999, 39-52.
- 4) Turgeon, A.J. Turfgrass Management. Prentice: 1996, 164-184.



Turf Specialist in Cultural Practices & Custom Grows
 Drill & Fill • Root Pruning • Aerification • Overseeding • DeepTine • Verticutting

For more information, contact us at
877-727-2100 www.zandersod.com

(Or contact our Turf Specialist, Alexander Dickie directly at 905-505-5014)



and get it **RIGHT!**



AERASEEDER



...for an easy way to denser turf.

100 Union Street, Elmira, Ontario, Canada N3B 2Z2
1-877-467-3478 • www.reistindustries.com