

TurfGrass TRENDS




Volume 5, Issue 8

August 1996

Field Testing of Biological Pesticides

by David J. Shetlar
Ohio State University

Over the last two decades, there has been a steadily increasing outcry for alternatives to standard, synthetic pesticides. Rachael Carson's "Silent Spring" was the first major alarm sounded pointing out that synthetic pesticides can often have widespread and undesired effects on animals and the environment. In the 1970s and 1980s, environmental groups, politicians and celebrities continued to decry the use of pesticides. Eventually, whether founded in fact or fiction, many people began to question the use of pesticides and sometimes attempted



1. Northern masked chafer grubs. The one on the right is normal, the one on the left is infected with a milky disease. Notice the drop of blood at the end of the snapped leg. It is clear in the normal grub and milky in the infected grub.

All photos provided courtesy of Dr. David J. Shetlar

IN THIS ISSUE

- **Field Testing of Biological Pesticides** 1
 - How are biological pesticides discovered?
 - Field testing - small plots
 - Scaling up
 - Training the user
 - A case study
 - Bacteria
 - Fungi
 - Entomopathogenic nematodes
- **The Basics of Turfgrass Fungicides**
 - Part Four: Handling and Applying Fungicides** 10
 - Measuring, weighing and mixing fungicides
 - Fungicides compatibilities
 - Tank storage time and pH affect fungicide efficacy
 - Proper equipment calibration
 - Fungicide placement
 - Post-application irrigation and fungicide efficacy
 - Monitoring the results of fungicide applications
- **In Future Issues** 16

TurfGrass TRENDS

Maria L. Haber

Publisher

Dr. Richard J. Hull

Editor

Dan Dinelli

Field Editor

Dr. Richard J. Hull

Science Advisor

Anna Schotanus

Circulation Manager

THE DESIGN GROUP

Layout & Production

TurfGrass TRENDS

1775 T Street NW

Washington, DC 20009-7124

Phone: 202-483-TURF

Fax: 202-483-5797

76517.2451 @ CompuServe.com

Abstracts: 800-466-8443 TGIF

Reprint Permission: 202-483-TURF

TurfGrass TRENDS is published monthly. ISSN 1076-7207. The annual subscription price is \$180.00.

Copyright 1996 *TurfGrass TRENDS*. All Rights Reserved. Copy permission will be granted to turf management schools.

Information herein has been obtained by *TurfGrass TRENDS* from sources deemed reliable. However, because of the possibility of human or mechanical error on their or our part, *TurfGrass TRENDS* or its writers do not guarantee accuracy, adequacy, or completeness of any information and are not responsible for errors or omissions or for the results obtained from the use of such information.

to get local regulations passed that would ban their use in domestic landscapes and on school grounds. Many home owners now are requesting that pesticides not be used unless absolutely necessary and many more are seeking alternatives to standard pesticides.

Entomologists have long been aware that most insect and mite pests have many predators, parasites and pathogens (lethal diseases) that can keep their populations below damaging levels. However, with the development of synthetic insecticides, much of the work on these biological controls was abandoned until the last two decades. This renewed interest in biological controls is in response, not only to the public outcry for alternatives, but because continued use of pesticides has resulted in the emergence of pests resistant to many pesticides and the uprising of "secondary pests." Secondary pests arise when their normal natural controls are killed by pesticides. Without these biological controls there is nothing to stop the increase of these pests to damaging numbers.

Because predators and parasites of insect pests are often difficult to rear and manage, many entomologists have turned to their pathogens - fungi, bacteria, virus and nematodes. These biological controls may be mass reared and are often applied as if they were regular pesticide sprays. This has given rise to the term "biological pesticide."

One of the confounding problems found, when developing a biological pesticide, is the field testing of these living organisms as if they were non-living, chemical pesticides. The process of developing a new biological pesticide is difficult and expensive. In order to bring a biological pesticide to market, a

company has to discover the pathogen, learn how to produce it, test it in small plots, test it in larger production settings, and market their new product.

How are biological pesticides discovered?

Location and identification

Many university and industry scientists are busy searching the globe for any new species or strains of diseases that attack harmful insects. These searches take several forms. Looking at native populations of the target pest may uncover significant natural controls, especially if the pest is an import to the US (i.e., Japanese beetle populations in Japan and China). Often, insects that are closely related to the target pest yield diseases with control potential (i.e., looking at sugar cane grubs for diseases that may attack turfgrass infesting grubs). Other scientists look for new forms of known pathogens (i.e., simply by taking soil and dust samples from around the world, we now have over 5,000 strains of *Bacillus thuringiensis*, a known insect pathogen).

In many cases, an infected insect is found and collected. The specimen may be used to expose additional insects so as to get more infected insects and increase the amount of the pathogen for further work. In other cases, especially if the pathogen looks like a genus of a known pathogen, the disease may be cultured on an artificial medium in order to get a larger sample. These samples are then "characterized" by standard microscopic examination (e.g., spore size and shape) or by molecular methods (e.g., protein characteri-



2. Japanese beetle grubs infected with *Beauveria* fungus.

zation, genetic makeup, etc.). Tests have to be performed to determine if the pathogen is, indeed, lethal to a target pest and if the new pathogen is an improvement over known pathogens. If the pathogen is sufficiently different from known examples and is an improvement, further development takes place and the pathogen is often "patented" in order to protect any future economic benefits.

Screening problems

Standard screening for new pesticides is usually targeted against the damaging stage of a pest while biological controls may be active only on specific stages or ages of pests. Japanese beetle adults can dine happily on one of the new BT strains while the larvae, especially young ones, are killed rather quickly. This brings up another problem with screening - using standard ages of the target. Many laboratory tests are performed on newly hatched insects. These tiny insects, all of the same age, serve as ideal experimental animals, but in nature, insect populations occur in mixed ages. As an example, several BTs will kill first instar sod webworms but little or no effect is obtained against the fourth, fifth and sixth instars. In cool-season turf in late June, there may be first through sixth instar bluegrass sod webworms present in any patch of turf. Therefore, standard screens using a uniform age-class of insects does not mimic field conditions.

Field testing - small plots

Figuring dose

Standard pesticide screens usually involve a range of concentration so that a dosage rating, usually the LD50, can be determined. However, with many biological pesticides a single spore or nematode can potentially kill the insect. Therefore, when varying concentrations of a pathogen yields no LD50, how does one determine the amount to use in the field? In many cases, simple guesses are made!

Benefits of small plot tests

Small test plots, usually in the range of ten by ten feet or less, are useful because small amounts of the pathogen can be used, the pest populations can be measured easily and are probably more uniformly present than in larger areas. Applications can also be determined more precisely by using highly calibrated equipment, and special environmental needs can be met, such as immediate watering. In these small plots, extreme ranges of the biological pesticide can be applied in order to better determine what the actual dosage has to be to perform adequately.

Small plot problems

A turfgrass stand is a complicated habitat. This habitat consists of the turf plants (leaves, stems,

roots), thatch, soil of varying textures and chemical makeup, changing moisture levels, other microbes, insects and animals. This is obviously very different from the laboratory petri dish in which a target insect and pathogen have been placed together. Therefore, most initial small plot tests involve a bit of "just tossing it out" (the pathogen, that is) experimentation. If this general toss doesn't work, then further tests are needed to try and hold some of the turf habitat traits constant. Sites with and without thatch may be needed, varying soil pH and moisture may be needed and other, possibly competing, organisms will have to be measured or eliminated.

Measuring efficacy

In standard chemical insecticide tests, the chemical is applied and after a short period, usually a week to a month, the insect "kill" is measured. This is usually based on the number of live insects remaining in the treated plots compared to "control" or "check" plots that were not treated. Many biological pesticides take considerably longer to act or they may do unexpected things to the insect. When white grubs become infected with the milky disease bacterium, *Bacillus popilliae*, they usually stop feeding immediately but they may remain mobile for several weeks to months before actually dying. Likewise, when caterpillars pick up certain strains of BT, they don't die within minutes or days, but may take one to two weeks before they expire.

Another problem with evaluating biological pesticides is choosing a "standard" for comparison. In chemical tests, this standard is often the top selling insecticide or a known insecticide within the same general chemical category. In many cases, biological pesticides are compared to these same standard pesticides. Standard pesticides have immediate effects while biological pesticides, especially ones involving living microbes, may be progressively lethal over time. Insects that became infected this week may not die until next season. However, they were eventually eliminated from contributing to the next generation.

Scaling up

Production

Once a biological pesticide has been successfully isolated, laboratory tested and small plot tested, the next major hurdle is to produce sufficient quantities of the pathogen to perform large plot or commercial sized applications. For many pathogens, this means moving from "counter top" production (production of small quantities in petri dishes or flasks), to medium fermentor production (perhaps ten to 100 gallons at a time). At this stage, many biological pesticides suddenly run into problems. In the larger production setting, the pathogen may lose its virulence or activity. Many bacteria and fungi seem to become "lazy" in the larger fermentor setting. They may lose their toxins, or the amount of toxin produced may be reduced. They may lose their viability or survivability. Therefore, constant testing for quality control must be performed in order to ensure that the cultured pathogen remains as active as the original organism.



3. A northern masked chafer grub infected with *Metarhizium* fungus in the process of producing its greenish spores (the darker patches within the white mycelia).



4. A Japanese beetle grub broken open to show the infection with *Steinernema carpocapsae* nematodes. The larger white curled nematodes are the adults while the "halo" around the body consists of hundreds of the new infective juveniles.

Formulation and packaging

In small plots, it is fairly easy to deal with unusual small containers of liquids containing a biological pesticide. However, if a gallon of the original material only covers a thousand square feet, then about 44 gallons will be needed to cover an acre. When compared to standard insecticides that may require one or less gallons to cover an acre, the weight incurred in shipping becomes a real, and expensive, problem. Many biological pesticides have limited shelf life. Most entomopathogenic nematodes can be kept viable for six months if not exposed to extreme heat. Many bacteria form resistant spores that can remain active for several seasons.

Commercial testing: making a fit

Once the small plot tests have yielded the specifics about application that allow the biological pesticide to perform at its best, fitting the new control product into the existing cultural system can be a real difficulty. Turfgrass is managed with a variety of fertilizers, herbicides, fungicides and insecticides. In some cases, these chemicals may be lethal to the biological pesticide, especially in tank mixes. Therefore, if a broadleaf herbicide is lethal to a bacterium being applied at the same time to kill black turfgrass ataenius adults, the applicator will often

opt for a standard insecticide that will not require two separate applications.

Training the user

Expectations

Based on past experience with standard insecticides, most people making their first application of a biological pesticide expect the same things to happen - rapid, and often visual, kill of the target pest. Golf course superintendents "expect" to see a "body count" of cutworms within hours after applying a standard insecticide. However, if a nematode or spinosad (a pesticide derived from a bacterium) is used, no cutworms appear on the surface, within hours or days.

A case study

Entomopathogenic nematodes

Attempts to use two groups of insect killing nematodes have occurred since the 1930's. These are now in the genera, *Steinernema* and *Heterorhabditis*. As previously mentioned, applied field work on these organisms was abandoned when modern synthetic pesticides were discovered.

In the late 1970's, interest in these nematodes was renewed in academic circles and a small "biotech" firm was established in California (now Biosys, Columbia, MD). With some infusion of venture capital, this firm spent considerable time learning how to produce moderate numbers of nematodes, *in vivo* (growing them in living insects), then large numbers of nematodes, *in vitro* (growing on artificial media), and finally massive numbers in thousand gallon fermentors.

At first, only one nematode, *S. carpocapsae*, appeared to be "cooperative" and readily adapted to *in vitro* production. Numerous tests were performed in the laboratory, in petri dishes and small containers containing target insects. There appeared to be few insects that *S. carpocapsae* could not kill in this manner. However, when university entomologists were given this nematode, they soon observed that field applications were not working or the nematodes only worked once in a while.

My evaluations of this nematode began in 1986. Tests in the laboratory demonstrated that this nematode could easily kill sod webworms, black cutworms, Japanese beetle grubs and northern masked chafer grubs. However, in small plot tests on a golf course, the grub control was sporadic and marginal. Realizing that these nematodes were applied in the microscopic, infective juvenile stage, we suspected that the standard application techniques may be killing the nematodes if immediate irrigation did not follow the application. Sure enough, if the nematodes were immediately watered into the turf with a minimum of $\frac{1}{4}$ inch of water, efficacy greatly increased (Shetlar et al. 1988). This has been reconfirmed through work by Downing (1994) and Yeh and Alm (1995).

Subsequent to this finding, we began larger scale treatments of entire lawns. Again, even with irrigation, the nematodes seemed to fail. Fortunately, we had saved some of the nematode material that was used. Under the microscope these nematodes appeared alive and healthy. However, when given a chance to kill insects in petri dish tests, nothing happened! Apparently, the nematodes had lost their ability to kill the insects. Was this a case of a

"lazy" pathogen, created by the *in vitro* process? Some rapid investigations by the Biosys scientists found that the active nematodes had non-pathogenic bacteria in their storage organs. The nematodes don't actually kill their host directly but they regurgitate a lethal bacterium in the insect's body cavity. The bacterium quickly kills the insect and the nematodes begin to reproduce while feeding on these bacteria. In our case, the bacterium had lost its virulence in culture. The nematodes didn't know the difference between lethal bacteria and nonlethal bacteria. The result, the nematodes were "shooting blanks"!

Armed with this new information, Biosys and many university researchers performed bioassays with the nematodes before using them in the field. This is now a standard procedure during the "quality control" process of nematode production. Finally, armed with nematodes that worked and the knowledge that irrigation and avoidance of direct sunlight improved nematode survival, Biosys wanted to begin selling their nematodes on a commercial basis. While working cooperatively with university researchers, several larger scale applications of nematodes were used by golf course superintendents for management of their black cutworms on greens and tees. It soon became evident that, while effective when used according to directions, the nematodes were not easy to mix and apply, when compared to standard insecticides. The nematodes arrived in jars containing a screen or sponge and these had to be thoroughly rinsed out in clean water. Many superintendents balked at the prospect of rinsing and washing five to 20 of these containers. Golf course superintendents also like to apply fungicides at the same time that they apply insect control. Many of these fungicides, herbicides and previously applied insecticides can be lethal to the nematodes.

The end result, golf course superintendents only used the nematodes if they had no choice. Some superintendents felt that being able to say that they had eliminated the use of insecticides to manage one or two pests was worth the extra effort. Most, however, wanted a biological control but something less difficult to use.

Subsequently, Biosys pioneered a new formulation, a water dispersible granule. The only drawback was that the granule had a shorter shelf life (less than one year) and fewer nematodes per unit volume could be contained. Through some inventive marketing, this product ended up being best suited for the home owner trade. In these markets, home owners only want small amounts to treat special problem areas of their landscapes. When flea larvae and pupae were found to be susceptible to these nematodes, the market increased dramatically.

Also, during the development of *S. carpocapsae* for cutworm and sod webworm management, different nematodes were being discovered and tested for other insect targets. University of Florida researchers soon found a nematode, now *S. scapterisci*, that appeared to be superior to any other at locating and killing mole crickets. This nematode was eventually licensed to a firm in Florida under the trade name of Proactant-Ss™. Soon thereafter, *S. riobravos* was described and this species also was good at attacking mole crickets. It is sold under the trade name Vector-MC™. Both products have enjoyed an increase in sales as golf course superintendents become more comfortable with the idea that these nematodes need immediate irrigation, applications should be made in the late afternoon to avoid direct sunlight, and dead mole crickets do not appear on the surface the next morning.

On other fronts, species of *Steinernema* that appear to be more suitable for white grub control have been reared in quantities sufficient to make them marketable (Selvan et al. 1994). Likewise, the *Heterorhabditis* species, which have been difficult to rear in mass, are now also being grown in commercial quantities. *H. bacteriophora* and related species have always been among the best agents for control of white grubs.

In retrospect, field evaluation and development of biological pesticides requires constant fine tuning of handling, mixing and application techniques. After becoming complacent to the sameness of using standard pesticides, we have had to find all the weak links in delivering a biological pesticide

and find solutions. For the end users, this will also require rethinking their ways of applying and utilizing control products.

References:

- Cranshaw, W.S. and M.G. Klein. 1994. Chapter 45. Microbial control of insect pests of landscape plants, pp. 503-520. In: A.R. Leslie (ed.), Integrated Pest Management for Turf and Ornamentals, Lewis Publishers, Boca Raton, FL.
- Downing, A.S. 1994. Effect of irrigation and spray volume on efficacy of entomopathogenic nematodes (Rhabditida: Heterorhabditidae) against white grubs (Coleoptera: Scarabaeidae). J. Econ. Entomol. 87: 643-646.
- Georgis, R. and G.O. Poinar, Jr. 1994. Chapter 43. Nematodes as bioinsecticides in turf and ornamentals, pp. 477-489. In: A.R. Leslie (ed.), Integrated Pest Management for Turf and Ornamentals, Lewis Publishers, Boca Raton, FL.
- Selvan, S., P.S. Grewal, R. Gaugler and M. Tomalak. 1994. Evaluation of steinernematid nematodes against *Popillia japonica* (Coleoptera: Scarabaeidae) larvae; species, strains, and rinse after application. J. Econ. Entomol. 87: 605-609.
- Shetlar, D.J., P.E. Suleman and R. Georgis. 1988. Irrigation and use of entomogenous nematodes *Neoplectana* spp. and *Heterorhabditis heliothidis* (Rhabditida: Steinernematidae and Heterorhabditidae) for control of Japanese beetle (Coleoptera: Scarabaeidae) grubs in turfgrass. J. Econ. Entomol. 81: 1318-1322.
- Yeh, T. and S.R. Alm. 1995. Evaluation of *Steinernema glaseri* (Nematoda: Steinernematidae) for biological control of Japanese and Oriental beetles (Coleoptera: Scarabaeidae). J. Econ. Entomol. 88: 1251-1255.

Bacteria

Several bacteria produce toxins that affect insects or cause an infection that kills the pests. Most bacteria produce a spore that can survive harsh environmental conditions and many bacteria can be grown in artificial media, therefore reducing the cost of their production.

Bacillus thuringiensis (commonly called "BT") has numerous strains that produce a toxin that affects the gut lining of specific insect groups. Affected insects stop feeding and die within a few days.

BT variety *kurstaki* - is registered under several trade names (Dipel Dust™, Sod Webworm Attack™, Bactosphene™, etc.) and is registered for sod webworm. Laboratory tests indicate good efficacy against first and second instar larvae but poor activity against larger larvae. The BT var. *kurstaki*, strain 'Spodoptera' (Javelin™) has shown good activity against the tropical sod webworm in Gulf States.

BT variety *israelensis* - is registered under several trade names (VectoBac™, Bactimos™, etc.) and is effective against mosquito larvae in water as well as black fly larval control in streams.

BT variety *japonensis*, strain 'buibui' - has recently been tested for control of white grubs with good success. Registered products are expected within a couple of years.

Bacillus popilliae is called the milky disease of white grubs. The bacterium causes infected insects to stop feeding and their body fluids to turn a characteristic white color. Infected insects may take weeks or months to die, even though they have stopped feeding. Numerous strains have been identified that attack certain species of white grubs. Only the Japanese beetle strain is in commercial production under several trade names (Milky Spore™, Doom™, Japademic™) by two firms: Fairfax Labs in New York (914)266-3705, and St. Gabriel Laboratories in Gainesville, VA (800)801-0061. Field studies in New England States have yielded 30 to 50% infection. Tests in Ohio and Kentucky have resulted in 20% infection or less.

Serratia entomophila is called the amber disease of white grubs. The bacterium causes infected insects to stop feeding and their body fluids to turn a honey-amber color. Affected insects turn flaccid within a few weeks and soon rot. A commercial product, Invade™, is being used in New Zealand for white grub management in pastures but no products are currently registered in the United States.

Fungi

In general, fungi usually require high moisture and are relatively intolerant of sunlight. Though they can often be cultured on artificial media, creating the right conditions for spore formation is usually the major problem in commercial production.

Beauveria bassiana is called the white fungus of insects. Infected insects become sluggish and eventually stop all activity. Within a few days or weeks the fungus sporulates by forming a dense white, cottony mass over the insect exterior. Chinch bugs and billbug adults are commonly attacked during periods of rainy, warm weather. A recent product,

Natuallis-T™, has been registered for use against a variety of agricultural pests as well as turf infesting mole crickets and chinch bugs. Sufficient replicated field tests of this material have not been performed against these turf pests.

Metarhizium anisopliae is called the green fungus of insects. Infected insects become sluggish and stop all activity. Fungal sporulation begins as a white coating but the blue green spores soon coat the exterior of infected insects. Though several strains are being developed by foreign and U.S. companies for management of white grubs, no commercial products are yet available.

Entomopathogenic Nematodes

These nematodes are specialized roundworms that carry a bacterium which is lethal to insects. The juvenile nematodes usually enter insects through the mouth, anus or breathing pores though some species may be able to penetrate through the insect cuticle. Once inside the insect, the nematode regurgitates its specific bacterium. The bacteria multiply, killing the insect and preventing other bacteria from colonizing the cadaver. The nematodes feed on the bacteria, mature and reproduce. These nematodes are not harmful to animals other than insects and they can not enter plant tissues.

Steinernema nematodes are commercially available under several trade names (Biosafe™, Vector™, Savior™, Scanmask™, etc.). *S. carpocapsae* is the most commonly produced species because of the ease with which juveniles can be grown in large fermentation tanks. *S. carpocapsae* is most useful for management of cutworms, sod webworms, billbugs and fleas. However, nematodes are very susceptible to desiccation, can not tolerate direct sunlight, and they may be killed by other insecticides or fungicides commonly applied to turf. *S. feltiae* and *S. glaseri* are also marketed for surface insect and white grub management. Steinernematid nematodes, in general, have not performed well for management of white grubs. *S. riobravos* (Vector-MC™) and *S. scapterisci* (Proactant-Ss™) are species registered for control of mole crickets and properly made applications have produced satisfactory control.

Heterorhabditis bacteriophora nematodes are commercially available but generally from smaller suppliers. Recently, Ecogen has begun larger scale production of this nematode under the name of Curiser™. This nematode has generally been the best performing species for control of white grubs.

Terms to know:

LD50 - the lethal dose of a pesticide or chemical required to kill 50% of a group of exposed plants or animals.

Entomopathogenic - literally insect killing disease. Entomopathogenic nematodes are microscopic nematodes that enter insects and release an insect killing bacterium.

In vitro - outside the body, to grow something in an artificial medium.

In vivo - inside the body, to grow something within a living organism.

Dr. David J. Shetlar is an Associate Professor of Landscape Entomology with The Ohio State University. He grew up in Oklahoma, obtained his BS and MS in Zoology from the University of Oklahoma and his Ph.D. in Entomology from Penn State. After working as a turfgrass research entomologist for Chemlawn R&D for six years, he joined OSU. His research interests center around development of biological and biorational controls of turfgrass pests, use of pest resistant turfgrasses and improving sampling and monitoring tools.

Erratum

In our most recent survey of golf course superintendents, under the question: "Which other publications do you read?" we erroneously included *Golf and Environment* in the list of alternatives. We should not have done that as *Golf and Environment* is a video magazine.

The results published were not accurate because respondents were unsure of whether *Golf & Environment* referred to the popular video magazine, or a print publication they never heard of.

We regret any confusion this may have caused and apologize for the error.

The Basics of Turfgrass Fungicides

Part Four: Handling and Applying Fungicides

*By Eric B. Nelson
Cornell University*

Acquiring knowledge of fungicide properties and their behavior in soils and plants is only half the job of implementing an effective and environmentally responsible fungicide program. Undoubtedly the most important part of this process is making sure that you are delivering the proper amounts of the correct fungicide to the appropriate place at the right time. To assure this, routine monitoring of your application procedures and equipment is necessary.

Studies have shown that the vast majority of turfgrass managers do not actually apply what they think they are applying. Nearly all make mistakes in mixing, loading, configuring equipment, and calibrating delivery devices. National losses due to these mistakes have been estimated to be in the billions of dollars. Additional losses have occurred because of reduced fungicide efficacy resulting from improper measuring and calibration. It is important, therefore, that care be taken in measuring, mixing, and loading fungicides and in routinely calibrating and maintaining equipment. Further precautions should be taken to assure proper timing and placement of fungicide applications.

Measuring, weighing and mixing fungicides

It is important that the proper protective clothing, including chemical-resistant gloves, goggles, and a respirator be worn when handling any fungicide since the concentrated forms of the fungicides can be particularly dangerous if splashed onto your skin or in your eyes. Also, some fungicide formulations such as wettable powders may be quite dusty during handling and may easily be inhaled. It is important to avoid smoking, eating, or drinking during fungicide handling operations since you could easily carry the fungicide to your mouth with contami-

nated hands or food. In general, utmost cleanliness and hygiene should be practiced during any and all fungicide handling operations.

Nearly all fungicides commonly used for turfgrass disease control are purchased as concentrated formulations and require some sort of measuring and mixing to dilute the fungicide prior to application. The amount of mixing and handling depends to a large extent on the type of formulation. Many granular formulations come packaged in bags in sufficient quantity to cover a designated area. Similarly, water soluble packets contain prepackaged fungicide formulations that are mixed with water and used to treat a designated area. In both of these cases, minimal measuring and weighing are required. However, for formulations such as wettable powders (WP), water dispersible granules (WDG), emulsifiable concentrates (EC), and flowables (F or FLO), a certain degree of measuring, weighing, and mixing are necessary for proper application.

It should be obvious that measuring out the correct amount of fungicide is critical for optimum fungicide efficacy. Too little may result in inadequate control and too much may result in phytotoxicity or other undesirable side effects. Both liquids and wettable powders/WDG's are mixed with water in basically the same manner. A given volume or weight of formulations is added to a measured volume of water. The amount of fungicide and water are determined from the desired rates of application and the output of the sprayer.

Fungicide compatibilities

When mixing fungicides together with other pesticides, growth regulators, or fertilizers, the compat-

ibility of the mixture can be a serious consideration in determining fungicide efficacy. In some cases, combinations resulting in enhanced levels of fungicidal activity have been identified. These include combinations of sterol inhibiting systemic fungicides and chlorothalonil for the control of a number of turfgrass pathogens, combinations of metalaxyl/mancozeb, fosetyl Al/mancozeb, chloroneb/thiram, and etridiazole/PCNB for the control of *Pythium* diseases, and anilazine/Zn (or Cu) for the control of anthracnose. However, in many cases, combinations of other chemicals with fungicides can reduce the efficacy of the fungicide. The physical and chemical compatibilities of the spray partners are of the most concern.

The physical compatibility of the materials should first be tested to be sure that no unwanted oily films and layers, foams, flakes, gels, or precipitates are formed. Additionally, wettable powders should be checked for lumps when mixed with some materials whereas some liquid formulations may settle into layers when mixed with other chemicals. Physical compatibilities can be tested easily by preparing the appropriate concentrations of tank-mixed components each in a small container. Add each component one by one to the fungicide suspension, shaking between each addition. When all of the components have been mixed together, gently shake the container and examine the contents immediately after shaking to see if there is any excessive foaming, and after 30 minutes to 1 hr to check for any precipitates. If the mixture does not look uniform, it should not be used as a tank mix.

The chemical compatibility of the tank mix partners should also be considered. Don't mix anything that will lead to a highly alkaline or highly acid condition, since this will lead to the degradation of some fungicides. Don't use adjuvants unless you know they are safe. If you are unsure of the phytotoxicity of a mixture, perform a test on a small area of turf before mixing on a large scale. Phytotoxicity can be affected by the air temperature, plant stress, plant genotype, etc. Finally, do not mix materials targeted for both foliar and root problems unless each material in the mixture behave similarly in the plant (e.g., they are each

contact materials, each localized penetrants or each upwardly-moving systemic fungicides). Otherwise, less than optimal control will result for one of the diseases in the complex. Similarly, do not mix fungicides with essentially the same mode of action. This can lead to phytotoxicity.

Fungicide formulations are more effectively mixed with other chemicals of similar formulation. For example, liquids can be mixed more effectively with other liquids and wettable powders or water dispersable granules can be mixed with other wettable powders or water dispersable granules. However, it is also common to mix fungicides with other materials having different formulations.

When mixing liquids and solids in the same spray tank, it is important that they be added in the correct order to insure proper dispersion and uniformity. A convenient way to remember the proper order is to use the sequence W-A-L-E where W stands for wettable powders and water-dispersable granules, A stands for agitation, L stands for liquids, and E stands for emulsifiable concentrates. The proper procedure is as follows:

1. Add wettable powders and water dispersable granules first to a tank half full of water.
2. Agitate until these formulation are uniformly dispersed while adding water until the tank is 90% full.
3. Add all flowable liquids and other water soluble formulations.
4. Finally, add emulsifiable concentrates.
5. Top off the tank and continue agitation.

The materials are now properly mixed.

As always, the tank contents should be properly and continuously agitated during spray operations since many formulations form suspensions and not true solutions. And finally, always consult the label for compatibility information. Most fungicide labels will list compatible or incompatible combinations when they are known and have been tested.

Tank storage time and pH affect fungicide efficacy

Fungicides should, whenever possible, be mixed and sprayed as soon after mixing as possible. However, in cases where fungicide mixtures are placed in the spray tank in advance of the application, special precautions must be taken to avoid chemical decomposition of the fungicide as it sits in the tank. One of the primary factors contributing to the instability of a fungicide is the pH of the water.

Most of the water used to prepare fungicide sprays in the United States is quite alkaline (high pH). Studies have shown that under these alkaline conditions, a number of commonly-used fungicides can break down and lose their effectiveness (Table 1). For example, anilazene, chlorothalonil, thiophanates, and thiram are all hydrolyzed at pH values greater than 9.0. A few fungicides such as fosetyl Al and benomyl are unstable at pH levels below 5.0. Fungicides such as iprodione, vinclozolin, propiconazole, and triadimefon are insensitive to pH and remain stable even after storage in the spray tank for 24 hr.

Even though many fungicides are relatively stable at extremes in pH, storage in the tank for prolonged periods of time will accelerate their decomposition and the loss of their effectiveness. For example, even though fenarimol is relatively stable when initially mixed, it is unstable at acid pH values when stored for 24 hours or more.

Because of the critical role of pH in fungicide efficacy, the water used for spray applications should be checked on a weekly basis and the pH adjusted if necessary. More importantly, the pH of the fungicide mixture should be determined and adjusted if necessary. A number of commercially available buffering agents are useful for such pH adjustments. A pH range of 5-6 is most desirable.

Proper equipment calibration

Considerable effort goes into the determination of proper fungicide application rates described on the

Table 1. pH stability and photostability of turfgrass fungicides

Fungicide	Comment
Chloroneb	Stable
Cyproconazole	Stable
Etridiazole	Stable
Flutolanil	Stable
Metalaxyl	Stable
Propamocarb	Stable
Propiconazole	Stable
Triadimefon	Stable
Benomyl	Unstable at pH<4
Fosetyl Al	Unstable in acidic (pH<2) and alkaline (pH>9) conditions
Anilazene	Unstable at pH>9
Chlorothalonil	Unstable at pH>9
Mancozeb	Unstable at pH>7
Quintozone	Unstable at pH>9
Thiophanate methyl	Unstable at pH>9
Thiram	Unstable at pH>9
Vinclozolin	Unstable at pH>9
Fenarimol	Photodecomposes rapidly
Iprodione	Unstable at pH>7, Photodecomposes in aqueous suspensions

Compiled from: Tomlin, C., 1994. The Pesticide Manual, 10th Edition, Crop Protection Publications, British Crop Protection Council, UK, 1341 pp.

label. It is important, therefore, that the proper amount of fungicide is delivered to the area to be treated. The effectiveness of any fungicide will depend on the proper application and placement of the material. Proper calibration insures that your application equipment is delivering the correct amount of fungicide to the area being treated. Even if you have meticulously weighed and mixed the fungicide, improper delivery of the spray will result in less than desirable disease control.

It has been estimated that 60% of all sprayers have calibration errors greater than 10%. Nearly 45% of all sprayers have more than a 10% variation in discharge from individual nozzles. In addition to these problems, many sprayers are used at inaccurate travel speeds and improper boom height for the type of nozzle and spacing, have pressure gauges that read too low, and have an inadequate match between hose size and nozzle type.

Determining the output of a sprayer

The output of a sprayer is one simple estimate of overall sprayer performance. It is the amount of spray material delivered per unit area. The output

can be measured by first marking off an area 100 ft by 100 ft or any area equivalent to 10,000 sq. ft. Fill the spray tank with water and spray the entire area as if you were applying the fungicide. When you have finished, measure the amount of water needed to refill the tank. Divide this amount by 10; this represents the delivery rate per 1000 sq. ft. Also the amount of spray delivered per 1000 ft² multiplied by 43.5 equals the amount applied per acre. Alternatively, you can determine the time it takes to cover the desired treated area. With the sprayer motionless, you can then collect the spray delivered in the predetermined time period and measure its volume. While it is useful to perform this test from time to time through the season to monitor sprayer performance, it will not reveal problems with unequal delivery among nozzles. These should be examined separately.

Calibration of nozzle output on boom sprayers

The following steps are recommended for the calibration of boom sprayers:

1. Make sure all nozzles are of the desired type and that the pressure at the nozzle is appropriate for the nozzle being used. Flat fan and swirl chamber nozzles often perform best at pressures of 30-60 psi.
2. Clean nozzles and screens to remove any material that could potentially clog the nozzle or impede delivery.
3. Check to see that the spray pattern from each nozzle is uniform and that the spray patterns overlap by 30-50%.
4. Measure the delivery volume of each nozzle. This can be done by placing the same-sized containers under each nozzle. If all containers fill at the same rate, your nozzles are OK. Replace nozzles that deliver more or less volume than the average nozzle output.
5. Select your operating speed (usually 3-5 mph). Be sure to use the same speed during calibration as that used during spray applications.
6. Determine the delivery rate as described above.

Calibration of granular applicators

Granular application equipment comes in a variety of sizes and consists of drop types and rotary types. In either case, calibration involves determining the

weight of material applied per unit area. In all cases, the granular material to be applied should also be used in the calibration since different granule sizes and shapes flow at different rates. Speed is usually not a critical factor but should be chosen such that it allows the material to flow freely.

It is important to realize that once your equipment is properly calibrated, it needs to be recalibrated and the delivery checked on a regular basis. Fungicide delivery may change with equipment wear, gauge error, nozzle wear, wheel slippage, speedometer error, and friction loss. It is important, therefore to monitor your equipment continuously and recalibrate regularly. This includes cleaning or replacing nozzles and checking nozzle pressure, checking nozzle spacing, boom height, and sprayer output. Proper calibration will insure that you are not wasting material or sacrificing fungicide efficacy.

Timing of fungicide applications

The timing of fungicide applications is another critical aspect of maximizing fungicide performance. Application timing is more complicated than it appears at first glance. Of obvious importance is the timing of an application relative to the active stages of the pathogen. However, other timing considerations include the time of day, temperature/humidity relationships, wind patterns, and practical considerations of traffic and public perceptions.

For optimum disease control, fungicide applications must be timed to coincide with periods when the target pathogen is in an active growth stage. This is the stage most susceptible to fungicide treatment. Most often these periods of pathogen activity correspond with symptom development in the turfgrass plant. Therefore, most fungicide applications are best made as a curative application after a correct diagnosis has been made. However, with some diseases, the period of maximum pathogen activity precedes the development of symptoms, sometimes by several months. This is the main reason why fungicides used for summer patch control must be applied in the late spring even though summer patch symptoms typically appear in mid to late summer.

Pathogens in a dormant stage are generally not susceptible to fungicides.

Another important timing consideration is the time of day, particularly as it relates to temperature and humidity relationships. Both temperature and humidity can affect fungicide drift. The higher the temperature and lower the relative humidity, the greater the opportunity for fungicide evaporation or volatilization. Under these conditions, small spray droplets may evaporate completely, leaving volatilized fungicide residues in the air where they may travel up to several miles from the spray site. This can be avoided by applying early in the morning when temperatures are lower and relative humidities are higher than is normally the case during the middle parts of the day.

In addition to the reduced drift hazard from fungicide volatilization early in the morning, drift may also be minimized in the morning hours because of calmer winds and lower convective air turbulence. As the turf surface heats up and solar radiation becomes stronger during the day, a greater temperature differential occurs between the turfgrass surface and the air. This creates upward air currents that can carry spray droplets away from the target site.

Another important timing consideration is the impact of spray applications on public exposure. With the exception of some golfers, most people are less likely to frequent turfgrass sites early in the morning or late in the evening than at other times of the day. Therefore, these times are ideal for avoiding potential public exposure to fungicides and for minimizing the opportunities for the public to become concerned over a pesticide application and to question the environmental responsibility of the pesticide application and of the applicator.

Fungicide placement

Fungicide placement is one of the more important factors affecting fungicide performance. Generally, if the fungicide does not come in contact with the pathogen, the disease will not be controlled. The nature of the disease to be controlled, the amount of

thatch, and some of the inherent properties of the fungicide being used all determine where the fungicide should be placed. For example, if the disease to be controlled is caused by a pathogen that infects and survives in the foliage, placement of the fungicide is generally not a problem. The fungicide can simply be applied as a spray. However, if the disease to be controlled is caused by a root-infecting pathogen, placement of the fungicide becomes more problematic.

The main difficulty in placing the fungicide in contact with root pathogens is getting the fungicide through the thatch layer. Generally, the thicker the thatch layer, the more impenetrable it is to fungicide movement. Since many of the fungicides used for turfgrass disease control are adsorbed quite readily to thatch, other techniques must be used to get the fungicide into the root zone. This can be accomplished either by aerification prior to the fungicide application, or by applying excessive amounts of water to leach the fungicide into the root zone.

Another consideration in fungicide placement is making sure that you avoid skips and overlaps when making applications. Skips leave untreated areas where disease symptoms may develop whereas overlaps may lead to phytotoxicity. There are various ways of monitoring your spray patterns. The most common method involves the use of dyes that color the turf slightly so that the actual spray pattern can be visualized. As with other tank mixed materials, however, care should be taken to assure that dye materials are compatible with the fungicides being applied.

Post-application irrigation and fungicide efficacy

Often, for the control of root diseases on turfgrasses, it is recommended that the fungicide be watered-in. This is because most fungicides are not taken up and translocated inside turfgrass plants to turfgrass roots and therefore must be moved into the soil profile to contact pathogens. On the other hand, if they are absorbed and only translocated upward in the plant, some action must be taken to place the fungicide in the root zone and allow the

fungicide to reach its intended target. Moving the fungicide through the turf/soil profile with water is usually the method of choice.

No firm recommendations are usually made regarding the amount of water required for optimum fungicide activity. This is because the water status of the soil, the soil type, and the chemical nature of the fungicide all affect how much watering-in should be done. Apply too much water, and you leach the fungicide from the root zone or dilute it to the point where it loses efficacy. Use too little water and the fungicide never reaches its intended target.

Little research has been conducted to establish optimum post-irrigation schedules for turfgrass fungicides. However, some general guidelines might be helpful. First, never water in fungicides used for foliar disease control. Studies have shown that if a fungicide applied for foliar disease control is not allowed to dry on the leaf surfaces, there is a significant reduction in its efficacy. For products such as sterol inhibiting fungicides applied for root disease control, the amount of water used to move the fungicide into the root zone should be sufficient also to wet the upper root zone. If the soil is dry to begin with, movement of the water front can be monitored to determine the depth of water penetration. If the soil is already moist, post-spray irrigation should not exceed 1 inch of water. On sandier root zones, this should be reduced to $\frac{1}{2}$ inch.

Often times, fungicides applied for root disease control may be applied in excess of 5 gallons of water per 1000 ft². In these cases, a minimal amount of post-irrigation watering is necessary. In all cases, the irrigation should be applied before the fungicide has dried on the foliage.

Monitoring the results of fungicide applications

Many times following a fungicide application, little effort is made to monitor the results of the fungicide application other than by observing that, after a few days, the disease problem does not seem to be getting worse or, alternatively, the fungicide application appears not to have worked. Often, detailed monitoring of the results of a fungicide application can shed light on the nature of the problem, point to potential equipment or application failures, effectively assess fungicide efficacy, and provide a means of adjusting fungicide timing or placement for more effective future disease control.

Dr. Eric B. Nelson is an Associate Professor at Cornell University, where he is affiliated with the Department of Plant Pathology. He has degrees in botany, from Indiana University, and plant pathology, from Ohio State University. Dr. Nelson is active in research on the ecology and control of soil-borne plant pathogens, concentrating on biological control of plant diseases. He also conducts outreach programs in turfgrass pathology. *TurfGrass TRENDS* is presently publishing his extensive series on disease control with fungicide applications.



Please return the
form and your
payment to:

TurfGrass TRENDS,
1775 T Street, NW,
Washington, D.C.
20009-7124

ORDER

TurfGrass TRENDS

ORDER

Name

Title

☐ **YES,**

SEND THE

TurfGrass TRENDS

SUBSCRIPTION THAT I

HAVE MARKED.

(12 ISSUES PER YEAR)

Business

Address

City

State

Zip

Phone

Fax

PAYMENT ENCLOSED:

☐ 6 MONTHS @ \$96.00

☐ 1 YEAR @ \$180.00

8/96

In Future Issues

- What Insecticides Can and Cannot Do
- Field Application of Trichoderma
- Index of Fungicides Registered for Turfgrass Disease Control
- Managing Turf for Minimum Water Use

TurfGrass TRENDS Editorial Board

Dr. Richard J. Hull
University of Rhode Island

Dr. Eric B. Nelson
Cornell University

Dr. Tim Murphy
University of Georgia

Dr. Michael G. Villani
Cornell University

TurfGrass TRENDS Advisory Board

J. Douglas Barberry, Turf Producers International

Richard Bator, Kirtland Country Club

F. Dan Dinelli, North Shore Country Club

Merrill J. Frank, Columbia Country Club

Michael Heacock, American Golf Corp.

Vince Hendersen, River's Bend Country Club

Paul Latshaw, Merion Golf Club

Kevin Morris, National Test Evaluation Program

Sean Remington, Chevy Chase Club

Tom Schlick, Marriott Golf

Jeff Spangler, Troon North Golf Club

Mark A. Tamn, NaturaLawn of America

Paul Zwaska, Baltimore Orioles

Use of TGT articles

Permission will be granted on request for TGT articles as course material and for reprints in publications.
For course material: We can group articles by subject for you.

Please send request to:

TurfGrass TRENDS

1775 T Street, NW, Washington, DC 20009-7124

Phone: 202-483-TURF Fax: 202-483-5797 E-mail: 76517.2451@CompuServe.com

Index and abstracts are available electronically through: Michigan State University, TGIF 800-466-8443; PLCAA, at <http://www.plcaa.org>.
TurfNet at <http://www.turfnet.com>

TurfGrass TRENDS

1775 T Street NW

Washington, DC 20009-7124

ADDRESS CORRECTION REQUESTED

FIRST CLASS

US POSTAGE

PAID

WASHINGTON, DC

PERMIT NO. 8796