



TURFAX™

of the International Sports Turf Institute, Inc.

The International Newsletter about Current Developments in Turfgrass
©1996 International Sports Turf Institute, Inc., - All rights reserved.

Volume IV Number 6

November-December 1996

CONTENTS:

- JB Comments — Sports Turf Surfaces -
 Diving and Surface Profile Water Content.
- Upcoming JB Visitations.
- Upcoming International Event.
- New Publications Available.
- JB Visitations:
 - Cultivar-herbicide phytotoxicity.
 - University turfgrass education accreditation.
- Putting Green Rolling Concepts.

JB COMMENTS - SPORTS SURFACES

Completely modified root zones are now becoming the norm on most college and professional sports fields across the United States and certain other parts of the world. A sufficient number of various types of complete root zone profile constructions are now in place on sports fields around the country, such that comparative observations are being drawn by interested observers.

Most of these complete root zone modifications involve some kind of high-sand root zone profile construction. An observation being mentioned is that more turf tearing and divoting are occurring on fields constructed above an impermeable polyethylene barrier, regardless of whether any supplemental drainage technique also is employed. Is this due to an excessive water content in the upper surface portion of the root zone profile that is associated with or may be attributed to the impermeable polyethylene barrier?

It has been the personal experience of this author that divoting is much more severe when the turf and/or upper portion of the soil profile is more wet. This is easily observed on horse race tracks where multiple races are run on a daily basis under varying conditions. Irrigating during the morning of race day results in increased divoting even on tracks with a perched-hydration zone, rather than a polyethylene barrier.

Obviously, rain just prior to or during a competition on a sports field will increase the extent of turf tearing and divoting. Fields that already have a high moisture content in the upper portion of the soil profile prior to the rain are likely to experience even more severe turf damage. It would be interesting to hear from the TURFAX™ readership as to just what their observations are on this question. Obviously, there is a need for definitive research to address this question on a sound basis rather than just general observations.

UPCOMING JB VISITATIONS:

Provided for Institute Affiliates who might wish to request a visitation when I'm nearby:

- Jan. 6 to 9 - Reno, Nevada.
- Jan. 18 to 20 - Orlando, Florida.
- Jan. 21 to 23 - Indianapolis, Indiana.
- Feb. 7 to 13 - Las Vegas, Nevada.
- March 1 to 3 - Montreal, Canada.
- March 11 to 12 - Columbus, Ohio.

NEW PUBLICATIONS AVAILABLE:

Conference Proceedings of the 67th International Golf Course Conference and Show. 121 pages. GCSAA. (1996).

The proceedings contains 1 to 3 page abstracts of 51 papers presented at the 1996 Conference in Orlando, Florida. The topic headings include: *Poa*: wanted dead or ALIVE, Grow-in pains, Identity crises, Innovative superintendent, *Poa*: wanted DEAD or alive, Northern exposure, Southern living, Golf course decision - makers forum, Audubon cooperative sanctuary program certification, Employment security forum, and Public golf forum.

Contact: Golf Course Superintendents
Association of America
1421 Research Park Drive
Lawrence, Kansas, 66049-3859, USA
Phone: (913) 842-2240
Fax: (913) 832-4449

COLOR ATLAS OF TURFGRASS DISEASES - by Dr.'s Toshikazu Tani and James B Beard. Ann Arbor Press, Chelsea, Michigan. 240 pages, 1997.

This book that focuses on diagnosis of over fifty major diseases of turfgrass, plus identification of the causal pathogen, with balanced emphasis on both the warm- and cool-season grass hosts. The 450+ large color photos illustrate multiple stages in the development of each disease, rather than just a single photo as is typical in most books.. Suggested cultural and chemical controls are given for each disease-causal pathogen. Step-by-step color photos of laboratory diagnostic techniques also are presented. A unique, one-of-a-kind book that is a must for turfgrass practitioners. The book price is US \$79.95 plus postage, and is available from Beard Books.

UPCOMING INTERNATIONAL EVENT:**World Scientific Congress of Golf Announced**

The Third World Scientific Congress of Golf will be held at the University of St. Andrews, Scotland, from July 20 to 24, 1998.

In 1994 there were over 300 delegates from 32 countries in attendance. Those attending will be from diverse backgrounds such as coaches, teachers, golf manufacturers, golf course designers, architects, turf agronomists, sport psychologists, and sports medicine specialists. The 1998 conference program will focus on the (a) golfer, (b) golf equipment and playing the game, (c) golf course design, (d) golf turf maintenance and management, and (e) social and ecological impacts. Those interested in presenting papers and/or attending should contact:

World Scientific Congress of Golf
Department of Physical Education
University of St. Andrews
St. Andrews, Fife, KY16 9DY, Scotland, UK.

ISTI Chief Scientist: James B Beard
TURFAX™ Production Editor: Harriet J. Beard

The goal of the six issue per year TURFAX™ newsletter is to provide international turf specialists with a network for current information about turf. This newsletter is faxed to all Institute Affiliates that use the ISTI technical assistance services on an annual basis. Faxing is more costly, but ensures quick delivery to those outside the United States.

For non-affiliates, a TURFAX™ subscription is available by annual payment of U.S. \$60.00. Payment may be made by sending a check to the address given below. Foreign orders please send a check or money order on a U.S. bank.

Direct inquiries to:
International Sports Turf Institute, Inc.
1812 Shadowood Drive
College Station, Texas 77840 USA
Telephone: (409) 693-4066
Fax: (409) 693-4878

JB VISITATIONS:

October - Japan

Conducted a technical assistance advisory visitation in Japan involving turfgrass research and educational programs for golf courses, horse race tracks and sports fields.

At one research facility there is an interesting field simulation model of a sports stadium with the perimeter heights varying in relative scale, plus positioning relative to sun that is consistent with the planned stadium. Turfgrasses are being grown inside, along with monitoring instrumentation in multiple locations to assess the total daily levels of radiant energy in the photosynthetically active range. It will be interesting to follow the findings of this innovative new study.

In terms of turfgrass cultivar assessments, El Toro zoysiagrass (*Zoysia japonica*) has now been introduced into Japan. Its performance in research plots has been very good compared to the existing grass selections used in Japan, none of which are formally named and maintained as pure cultivars. El Toro has superior low temperature color retention, root development, and turfgrass quality.

In the creeping bentgrass (*Agrostis stolonifera* var. *stolonifera*) cultivar studies for putting greens, Penn A-2 is showing superior performance after two years of assessments.

Also visited turfgrass research plots where the herbicide selectivities of three *Zoysia japonica* cultivars were being studied. Each exhibited different phytotoxicities ranging from none to significant injury. These results emphasize the need and responsibility for the developer of a cultivar to evaluate the pesticide selectivity characteristics by the time of release, rather than just assuming all cultivars within a species are similar in their tolerance to pesticides. Otherwise, the end-user may suffer the consequences.

November - Indianapolis, Indiana

Participated in the Annual Meeting of the American Society of Agronomy. Among the eight divisions of the Crop Science Society of America,

only the C-5 Turfgrass Division is growing in number of members. The other seven agricultural crops divisions are declining in membership.

This same trend is occurring in terms of undergraduate enrollments at university Agronomy-Crops-Soils Departments around the United States. In an effort to cushion this decline in enrollment, most of these departments are placing great emphasis on developing a turfgrass science undergraduate education program. Some universities are putting significant amounts of money toward this goal, while others are only using promotional techniques without any significant investment in a sound undergraduate turfgrass science education program. In the case of certain universities, they are actually hiring a new faculty member whose primary responsibility is to develop or expand an undergraduate turfgrass teaching program. This is occurring even at universities which have exhibited minimal to no interest in turfgrass science in the past.

There are obvious questions resulting from this increased emphasis on turfgrass science education at universities. What are the actual employment needs in terms of the number of graduates trained in the turfgrass science area? Are we headed toward a saturated job market for formally trained BSc turfgrass graduates, especially in the golf course superintendent sector?

There also is the question of how an aspiring turfgrass student can sort through all the promotional hype from the various universities? How do they determine which universities actually are investing in a significant, comprehensive set of specific turfgrass courses, plus the needed laboratory instructional equipment, turfgrass library acquisitions, and summer work-study training programs that are needed? In view of the potential problems in this developing situation, strong arguments could be made for the development of an **undergraduate turfgrass accreditation assessment program** sponsored by a national turfgrass organization. It has been accomplished very effectively in such areas as forestry and veterinary science. Why not a similar program in the turfgrass area to avoid some of the potential problems?

PUTTING GREEN ROLLING CONCEPTS

by

Dr. James B Beard

The golfer commonly refers to putting green surfaces in terms of being fast or slow, which refers to the speed or velocity of ball roll. By definition these terms would refer to the distance traveled per unit of time. In actuality, research techniques and practitioner assessments, such as the stimpmeter, measure the distance of ball roll. Thus terminology used in this paper will be the more correct ball roll distance (BRD).

BALL ROLL RESISTANCE

The BRD as a result of a given impact force is affected by the sum total of the resistances (R_t) caused by the grass shoot/leaf characteristics (r_g), surface irregularities (r_s), and atmospheric (r_a) dimensions. The greater the total resistance, the shorter the BRD. The total ball roll resistance can be expressed by the following equation:

$$R_t = r_g + r_s + r_a$$

To fully understand and manage the putting green surface quality in terms of the desired ball roll distance, one must recognize the components contributing to the three resistances to ball roll, which are summarized in Table 1.

Atmospheric resistances (r_a) encompass the wind velocity and direction, and to a much lesser extent the density of the atmosphere as related to altitude. Individuals fail to recognize the major impact of wind velocity and direction on ball roll. Wind influence is much greater than the individual turfgrass cultural practices. Research involving ball roll distance assessments conducted without a wind barrier surrounding the measurement area are highly questionable.

There are interactions among the various components of turfgrass shoot-leaf resistances (r_g) which can either have positive or negative effects. A higher shoot density is being promoted as giving a greater ball roll distance. However, this is not necessarily the case. For example, the leaf

density of a fine-leaf fescue putting green is quite high and the leaf is very narrow. However, the ball roll distance will be much shorter than for a creeping bentgrass green with a lower shoot density and wider leaf. The reason is the distinct leaf stiffness of the fine-leaf fescue caused by extensive sclerified tissue in comparison to a creeping bentgrass leaf. Leaf growth extension rate can strongly influence the resistance to ball roll, which in turn is controlled primarily by the cultural practices and microenvironment that cause either an increase or slowing of leaf growth.

Table 1. Components of the three resistances to ball roll.

Turfgrass Characteristics (r_g)	Surface Irregularities (r_s)	Atmospheric (r_a)
<p><u>Leaf Stiffness:</u> sclerified tissues lignin content potassium level species/cultivar</p> <p><u>Leaf Extension Rate:</u> cutting height nitrogen level temperature water content species/cultivar</p> <p><u>Leaf Orientation:</u> mowing pattern species/cultivar</p> <p><u>Leaf Density:</u> cutting height nitrogen level species/cultivar</p> <p><u>Leaf/Stem Ratio:</u> cutting height nitrogen level species/cultivar</p>	<p><u>Turfs Altered By:</u> coring slicing spiking vertical cutting footprinting ball marks spike marks</p> <p><u>Impediments:</u> dew exudate water topdressing weeds seed heads</p> <p><u>Contours</u></p>	<p><u>Wind:</u> velocity direction</p> <p><u>Air Density:</u> water vapor content atmospheric pressure (altitude)</p>

The surface resistances (r_s) influencing ball roll distance are a much discussed concern that can be controlled to a great extent by cultural practices. When considering the range of components affecting the resistances to ball roll, there are two cultural practices that have the greatest overall effect in modifying the surface resistance. These components are (a) the mowing practices, including cutting height, mowing frequency, mowing pattern, and allied grooming operations and (b) turf rolling which effects not only the ball roll distance, but the smoothness of the surface. The remainder of this paper will address turf rolling practices that influence the surface resistances to BRD.

RENEWED INTEREST IN TURF ROLLING

The option of turf rolling reentered the cultural program with the frequent use of high-sand root zones in the construction of modern putting greens, and is being driven by the golfer's desire for fast putting greens. Use of high-sand root zones of the proper particle size distribution, such as the USGA perched hydration method, results in minimal susceptibility to soil compaction. Such root zones can be rolled without imparting serious compaction effects; thereby offering the potential for improved smoothness and uniformity of ball roll. It is particularly attractive in that ball roll distance can be enhanced via turf rolling, which reduces the need to use excessively close mowing heights that result in turf thinning and the subsequent development of moss and algae problems.

Two alternatives to turf rolling in terms of achieving an increased ball roll distance on putting greens are extraordinarily close mowing and frequent topdressing. However, the very close mowing required eventually introduces problems in terms of a weakened turf, with resultant thinning that provides openings for sunlight to reach the soil surface, thereby facilitating the invasion of moss and algae. Topdressing is more expensive and disruptive of play, although it is an essential cultural practice that offers other beneficial responses, such as enhanced biodegradation of potential thatch accumulations and filling of certain types of depressional areas that can not be accomplished via rolling.

This author first observed a newly developed mobile, mechanically-powered turf roller for putting greens over ten years ago in Melbourne, Australia. This led to authorship of a turf rolling article in the January 1986 issue of *Grounds Maintenance*. Six years later, the interest in turf rolling of high-sand root zone putting greens started to increase. Turf equipment manufacturers accelerated the development of powered mechanical models of turf rollers specifically designed for putting greens.

TURF ROLLING INVESTIGATIONS

Reports from five turf rolling studies have been published. Each addresses specific dimensions of turf rolling on either a short-term or long-term basis. The BRD was assessed in all five studies by the stimpmeter method.

- ISTI - International Sports Turf Institute (1)
- OSU - Ohio State University (2)
- PSU - Pennsylvania State University (3)
- NCS - North Carolina State University (4)
- MSU - Michigan State University (5)

The findings from these five investigation sites are summarized as follows:

Basic Rolling Effect. Based on the studies conducted to date, the first morning response from turf rolling may typically be a ball roll distance increase in the order of 10%.

Rolling Effect Duration. These studies suggest that at least a 2-day effect on BRD can be achieved from a single turf rolling in many situations.

Turf Roller Weight. ISTI studies revealed no increase in ball roll distance when the single roller pressure as one event was increased from 4.8 to 11.9 pounds per lateral inch. Note that while roller weight may not significantly affect BRD, it may be an important factor affecting the smoothing benefits of turf rolling, with higher weights being more beneficial.

Diurnal Rolling Response. Following a single early morning rolling, the ball roll distance typically declined throughout the initial day after rolling. This was probably caused by increased ball roll resistance due to vertical leaf extension. There was a major reversal to a higher BRD after mowing on the 2nd day, even though there was no subsequent second-day turf rolling. A similar diurnal response of lower magnitude was observed in some cases on the 3rd day.

Single Versus Multiple Roller Effects. In two ISTI studies, a single event rolling in early morning involving 1, 2, 3, and 4 single roller passes were evaluated. The most significant response was achieved from 4 consecutive single rollings in one early morning event, with the BRD increased an impressive 20% and persisting more than 76 hours.

Rolling Interval Effects. The difference in ball roll distance between the rolled and unrolled turf plots increased throughout the 2- and 19-week duration of two studies. Also, there was an increase in BRD as the frequency of turf rolling was increased from 1 to 4 to 7 times per week.

Roller Operational Effects. ISTI studies assessed single roller operating speeds of 2.5, 4.0, 4.4, and 4.8 feet per second. There was no observed effect on the BRD. Similarly, there was no effect on ball roll distance when turf rolling following early morning mowing was in the same versus the opposite direction as the mowing operation. MSU studies of rolling before early morning mowing had a minimal effect on BRD, when compared to rolling immediately after mowing or rolling at midday to increase the afternoon BRD.

Turf Responses. The OSU study of 2-weeks duration revealed that turf rolling had negative effects on the visual turf quality, including a negative color response and some wear stress.

A 10-week turf rolling duration on a clayey soil in the NCS study resulted in a reduction in turf quality, but no effect on the root mass. Turf rolling at a once-per-week frequency in the NCS study resulted in no adverse affects on turfgrass quality, thatch accumulation, or root biomass.

Disease and Moss Responses. MSU studies showed a turf rolling schedule of 3 times per week significantly reduced the severity of dollar spot disease. In contrast, Microdochium patch (pink snow mold) was significantly increased as a result of rolling 3 times per week. There also was a substantial reduction in the extent of moss invasion on the rolled treatments compared to the adjacent untreated checks. The rolling effects also resulted in a decrease in localized dry spots.

Root Zone Effects. Both the soil bulk density in the surface 1-inch (25 mm) of root zone and the saturated infiltration rate were not affected by turf rolling on a USGA specification high-sand root zone. Rolling on the clayey root zone resulted in increased bulk density.

A high bulk density is associated with a more compacted soil, poorer aeration, and slower infiltration rate. Turf rolling had no significant effect on bulk density of two high-sand root zones of the proper particle size distribution. The increase in bulk density of the fine textured soil is expected and a concern in turf rolling on such soils. There also is the question of how much turf cultivation will be needed to correct the compaction effects of turf rolling on these soils.

Enhanced Surface Smoothness. All these studies resulted in visual improvements in surface smoothness that includes both reduced lateral deflection and reduced vertical bumpiness in ball roll action. Due to difficulties in quantitatively assessing these effects, no specific quantitative data are available at this time.

Note: The findings just summarized relate only to creeping bentgrass putting greens. To date there have been no turf rolling investigations conducted on hybrid bermudagrass putting greens.

REFERENCES

1. Beard, J.B., S. Hammon, M. Morris, J. Holmes, and D. Kendziorski. 1992. Turf rolling effects on ball roll distance of bentgrass greens. International Sports Turf Institute Research Report No. 302.
2. Danneberger, K., E. McCoy, and T. Parobek. 1993. Rolling greens - Help or Hindrance? Turf Central. 4(11):7-10.
3. Hamilton, Jr., G.W., D.W. Livingston and A.E. Gover. 1994. The effects of light-weight rolling on putting greens. Science and Golf II: Proceedings of the World Scientific Congress of Golf. 1994. St. Andrews, Scotland. pp.423-430.
4. Hartwiger, C.E. 1995. Light weight roller use on bentgrass putting greens. MSc Thesis, North Carolina State University. 35 pp.
5. Nikolai, T., B.E. Leach, M.A. Smucker, D. Gaffner, P.E. Rieke, and J.N. Rogers III. 1996. Three soils and a lightweight greens roller Part II. 1996 Michigan Turfgrass Field Day, Michigan State University Extension. 1996. East Lansing, Michigan. p. 18.