

Monthly Newsletter

PROUD SPONSOR

Garden State Sports Turf

Maximizing Athletic Field Safety & Performance

- Complete Field Renovation Company
- Chemical applications
 - Field Maintenance

201-998-3293

John Salisbury

SFMANJ BUSINESS

Next Board meeting-October 15, 3:30 pm at Rutgers University. Horticulture Farm 2, Geiger Conference Room, Ryders Lane, New Brunswick.

During our September meeting nominations were received for 2002 Board of Directors. We are looking for two representatives in each category:

Private & public schools, University's & colleges, Education/Extension, Parks & Recreation, Professional facility, Commercial/contractor.

Any SFMANJ member may be nominated. All officers will be elected from the Board of Director's. Please send nominations to the above address or e-mail by Oct. 12.

WATCH for your membership renewal forms in October. Interested in becoming a member? Contact Steve Ratto at <u>equiturf@aol.com</u> or 201-837-2140. Members receive discounts to Rutgers classes and Floyd Perry seminars. Spring and fall field days are planned. Join us at the Turf Grass Expo in Atlantic City for our annual business meeting.

MONTHLY FIELD TIPS

Pregermination is a nifty little trick-of-the-trade that has been modified by many sports turf managers. The following information was published in SportsTurf written by Mike Andresen, sports turf manager for the Iowa State University Athletic Department. Use this modified version for soccer goalmouths, football field player's areas or lacrosse goal areas.

Preparation

Begin the pregermination process four or five days before you plan to apply seed to the field.

Put 5-pounds of seed into a 5-gallon water-tight bucket. Keep four 5-gallon plastic buckets on hand and mark them 'for pregermination only'. The woven-plastic seed bags that seed typically comes in are perfect for pregermination. To sponsor the SFMANJ

newsletter

Write to PO Box 370, Annandale, NJ 08801 – call 201-837-2140 Cost: Members \$150 Non-members \$175

Fill the container with water so that the seed is completely immersed. Keep at room temperature (70F).

Recommendations specify a 12-hour soak cycle, followed by a 12hour drain cycle, followed by another 12-hour soak cycle. For convenience, remove the bags each morning and allow them to drain during the eight-hour workday. Before the end of the work day, set up another soak cycle for the night.

If you forget about the bags for two or three days you still will obtain a good germaination.

Application

Net-Ster Michael

The second s

062

Pregerminated seed can be applied to a field by hand or with conventional seeding equipment. It's important to remember that root growth of pregerminated seed has already begun. You must use the seed within about five days of the start of the soaking process.

The seed must be dried before it's fed into spreaders or seeding equipment. Open the bags and spread the seed on a concrete floor with rakes. Calcined clay can be used to speed the drying process and make it easier to seed. Remember, seed left in equipment will continue to grow and will clog the system. The tubes in drill seeders are particularly vulnerable.

Repair

Seed can also be combined with topsoil, a combination of mushroom compost and topsoil (60/40) or sand mix to make a mix that can be placed in a specific area of the field by hand. Here are a few pregerminated mix combinations using a 5-gallon bucket to measure your topsoil.

Kentucky bluegrass: 0.2 lb. Pregerminated seed per bucket

Perennial ryegrass: 1.0 lb. Pregerminated seed per bucket.

Bluegrass/ryegrass mix (30/70% by weight): 0.5 lb. Pregerminated seed per bucket.

Dump a bucket of topsoil mix or sand mix on a concrete surface and spread a layer one inch thick. Spread the appropriate amount of seed over the topsoil or sand by hand. Use rakes and flat shovels to roll the pile together and mix in the seed. If the seed is too wet, add some calcined clay to soak up the water and make it easier to handle.

Simply fill the areas with the topsoil/seed mixture and pack it down with your foot or a roller. Spread a ¼ inch layer of the mix over thin areas. Water

Volume 4 •• October 2001