

“Protecting Turf from Winter Injury”

by John M. Roberts

The short days and cool temperatures of fall signal the start of winter dormancy for turfgrasses. Considering the heavy play most sports fields receive in early spring, it's critical that turf survives the winter. Damaged fields rarely have enough time fully to recover before the words “play ball” will be heard.

“Winter kill” is a generic term used to describe any loss or injury of turf during wintertime. From a prevention standpoint, it's helpful to break down the major causes of winter injury into more specific categories, which include the following: direct low temperature, traffic, winter desiccation, crown hydration damage (alternating freezing and thawing temperatures in wet soils), and low-temperature diseases. Ice covers are generally considered to be “indirectly” responsible for turf injury by forming a gas impermeable lens that creates an unhealthy environment for the turf below.

In a nutshell, the following preventive measures will help turf survive the winter.

High Fall Potassium Levels

Research continues to demonstrate potassium's benefit to turfgrass by improving its tolerance to various environmental and biological stresses, including drought, wear, heat and winter damage.

Potassium is highly water-soluble and easily leaches from plant tissues and sandy soils having a low cation exchange capacity. As a result, unexpected potassium deficiencies can occur. The use of slow-release potassium sources or more frequent, light applications throughout the year helps prevent this loss.

While potassium levels should remain high throughout the growing season, late fall is an especially important period for winter survival. In effect, potassium acts like antifreeze within a turf, enhancing its winter hardiness. Applying low nitrogen, high potassium fertilizers (1:2 to 1:5 ratios) in late fall continues to be popular among grounds managers to provide fields with potassium for the winter.

Low Nitrogen during Hardening

Approximately 30 to 40 days before winter dormancy, known as the winter hardening period, nitrogen (especially fast-release sources) should be used sparingly. Other practices that encourage active growth during this period are also discouraged. Unlike potassium, nitrogen during the hardening period increases tissue hydration levels and stimulates new growth, producing tissues that have thin cell walls. The net result of this over stimulated growth is turf more susceptible to freezing stress and winter diseases.

Drainage, Drainage and More Drainage

One of the key principles in reducing winter damage on sports fields is to provide rapid soil drainage. Poorly drained fields are highly vulnerable to an array of winter injuries. Unless the drainage is improved, it's usually just a question of time before large sections of turf are lost.

Both the subsoil and the surface need to drain freely. Installing drain lines, constructing fields with coarsely textured soils, and aerifying to relieve compaction help improve the water infiltration and percolation rates. To reduce standing water and accelerated surface runoff, fields in the northern Sports Field Managers Association of New Jersey

Established 1925

Fertl Soil

Specializing in Topdressing and Construction Soils
For Golf Courses and Athletic Fields

548 Rosedale Road

Kennett Square, PA 19348

610-444-0496

states are often crowned (14 to 18 inches) to compensate for the impervious nature of frozen soils (even sandy textures).

Avoid Over Watering in Late Fall

Late fall irrigations should be either avoided (preferably) or, if necessary to prevent drought stress, lightly applied. A grass plant prepares for winter by under going a number of physiological changes, including a dehydration of its tissues. This “drying out” condition in late fall is necessary for turf stands to achieve their maximum levels of winter hardiness. Otherwise, wet or saturated tissues are especially susceptible to direct low temperature kill, winter diseases and crown hydration injury.

Increase Mowing Heights

If feasible, skip the last mowing or raise the mowing height by ½ inch in mid fall. This allows turf to increase its carbohydrate reserves, which are vital for winter survival. Investigators have shown turfgrasses to be particularly vulnerable to winter injury during the late winter and early spring when carbohydrate levels are at their lowest. Warning! Turfgrasses are generally more susceptible to snow molds at higher mowing heights.

Continued on next page

SFMANJ “Update” Advertisers, it's Time to Renew !!!!!!!

Receive a 10% discount if you renew or extend your ad for the entire 2003 season. This offer is good only if confirmation is received prior to or at the NJ Turfgrass Expo in Atlantic City, December 10-12. See us at our booth or call Jim at 908-236-9118.

2003 ADVERTISING RATES

	SFMANJ Members rates:	Non-members rates:
1/8 page -	\$50	\$60
¼ page -	\$75	\$90
½ page -	\$120	\$144
Full page -	\$150	\$216

Add \$25 for set up fee for each ad (this includes changing ads)

Purchase 5 consecutive issues and receive the 6th issue free.

Sponsor rates (last page of the “Update” Newsletter) \$180/ 1 yr

SFMANJ RESOURCE GUIDE WILL BE AVAILABLE TO MEMBERS IN MARCH 2003.

To advertise in the resource guide a vendor must be a commercial/contractor SFMANJ member. This guide will provide all SFMANJ members with member information, participating service and product providers and periodic updates.

CALL FOR MORE INFORMATION (908) 236-9118

Continued from page 5 "protecting Turf from Winter Injury"

Remove Excess Thatch

Thick thatch layers tend to dry out quickly and serve as a harboring place for snow mold fungi. Winter disease and desiccation damage will be reduced if coring or thatch removal is practiced during the year. Late fall aeration may lead to desiccation around the coring holds during winters when there is not snow cover.

Traffic Control

The brittle tissues of frozen turf during the winter are prone to injury by traffic. The most severe damage seems to occur when bare or slush-covered ground exists. Snow (especially dry snow) acts like an insulator, protecting the turf below from traffic and direct low temperature injury.

Disease Prevention

Two of the most common and destructive low-temperature fungi are the two snow molds, pink and gray. Like Typhula blight (gray), pink snow mold can occur under snow, or is often observed in the absence of snow cover during cool (less than 50 degrees F), wet weather in fall, winter of spring. Gray snow mold is common in northern regions that receive more than 90 days of snow cover. It is particularly severe when snow covers partially or completely unfrozen ground.

A combination of fungicides and cultural practices is needed to provide acceptable levels of control in locations where disease pressure is high. Cultural practices that improve drainage, reduce thatch and maintain a balance fertility program (moderate nitrogen levels) help reduce both diseases. In general, contact-type fungicides are used for the prevention of gray snow mold and should be applied within a few days of snowfall.

Protective Blankets, Topdressing or Straw

Protection from low temperature injury, earlier spring green-up, and reduced desiccation are just a few of the benefits synthetic covers, topdressing, straw or the selective placement of snow fences can provide. Geotextile covers are also used to protect young seedlings and speed up germination or regrowth between hash marks and around the goal mouths of soccer fields.

Unfortunately, winter covers are not a panacea and will not solve all winter problems, including ice-related damage and crown hydration injury. Unless treated, cool-weather diseases are also more damaging under covers.

Summary

A better understanding, innovative ideas, genetic breakthroughs and even small miracles might be necessary to eliminate all forms of winter injury. However, using today's "best management practices" that promote rapid soil drainage and encourage healthy, winter-hardened turf going into winter is a grounds manager's best line of defense. ▲

John Roberts is an extension turf specialist at the University of New Hampshire.

This article was originally published in the October, 1996 issue of Sports Turf

"Are you Putting your Field to Bed or Putting it Out to Pasture?"

by Dave Minner and Gary Peterson, Extension Turfgrass Specialist, Iowa State University

If you are putting your field to bed this fall and tucking it in nicely with some love and care than you can expect it to wake up next spring ready to go and ahead of the game. If you are darn glad the fall season is over and you don't want to see that field until next year, then expect it to look like a pasture because you are treating it like one. If you plan to use the field in early spring or summer than it will be in the same condition that you left it at the end of the autumn. Instead, trying to do all of the coring, seeding, and field preparation in the spring may leave too much to chance with wet or cold weather. Get a jump on next year by putting your field to bed so it will wake up ready to go. Here are a few end-of-the-season tips that will help you prepare your baseball/softball fields for next year.

Baseball/softball

The fall practice schedule for baseball and softball is usually *Continue on page 9.....*

YOUR "ONE-STOP SOURCE"

FOR AMERICA'S LEADING
BASEBALL SURFACES & SUPPLIES



THE PROFESSIONAL'S CHOICE
... SINCE 1922

USED BY OVER 100 PRO TEAMS,
OVER 600 COLLEGES, PLUS THOUSANDS
OF TOWNS & SCHOOLS WORLDWIDE.
SPECIAL MIXES FOR INFIELDS,
PITCHER'S MOUNDS & HOME PLATE AREAS.

**REGIONAL INFIELD MIXES
AND RED WARNING TRACKS
FOR EVERY STATE & CLIMATE!**

PLUS INFIELD CONDITIONERS
TO IMPROVE EXISTING INFIELDS:

IF TOO HARD AND POORLY DRAINING!

Pro's Choice

THE REDDER, LESS DUSTY, MORE UNIFORM
INFIELD CONDITIONER & DRYING AGENT
"RED" OR "GREY" FOR INFIELDS

"GREEN" FOR TURF

FOR CONSISTENT INFIELD CUSHION
IN WET OR DRY WEATHER!

IF TOO SOFT & DUSTY!

STABILIZER®

FOR FIRM, YET RESILIENT, PLAYING SURFACES

TO QUICKLY DRY INFIELDS!

The Original & Most Absorbent is Now

NEW & IMPROVED GRANULAR

DIAMOND-DRY.



OVER 200 OTHER INFIELD PRODUCTS!

Now in 6 Colors!

FENCEGUARD
Protective Safety Covers
for Chain Link Fence

WALL PADDING • WINDSCREEN • RAIL PADDING
TAMPERS • DRAG MATS • RAKES
HOLLYWOOD® BASES • FIELD MARKING MACHINES
BATTING PRACTICE COVERS • RAIN COVERS
PERMA-MOUND™ PADS • MOUND BRICKS
SAFE "T" MAT™ BATTER'S BOX PADS
TYPAR® & TERRA-BOND® GEOTEXTILES
ON-DECK CIRCLES WITH TEAM LOGOS
PERMANENT FOUL LINES & MUCH MORE!

**FREE INSTRUCTIONAL BROCHURES
DISTRIBUTION CENTERS NATIONWIDE!**

800-247-BEAM

908-637-4191 / FAX 908-637-8421

PARTAC PEAT CORPORATION
KELSEY PARK, GREAT MEADOWS, NJ 07838

"The best infield mix I've ever used."

— GEORGE TOMA