

mapping and surveying is the thing to do. Using graph paper, prepare a general map of the turf area including landmarks. Mark sampling spots at 6-10 feet (lawns) or 10-20 feet (sports fields) apart in a grid pattern. At each spot take a sample and record number, species, and stage of grubs found (also record 0s!). Experienced samplers can process about 20 samples per hour.

To determine whether treatment is required, transform the grub numbers into 'per ft²'-values and compare to damage thresholds. Most published damage thresholds lie in the range of 6-10 (Japanese beetle, oriental beetle, European chafer) and 15-20 (Asiatic garden beetle) grubs per ft². However, damage thresholds vary considerably with grass species, management type, and climatic conditions. In well-maintained tall fescue plots I have repeatedly observed grub densities in the range of 30 to 60 grubs per ft² without any signs of turf damage. With experience, turf managers should develop their own range of thresholds for the various turf areas they are responsible for.

Preventative white grub control

The arrival of new insecticide chemistry with long residual activity in the soil in the last few years has added preventative applications as an option in the management of white grubs. The 2 insecticides presently on the market, Merit, a neonicotinoid, and Mach2, an insect growth regulator, can be applied as early as late May and June, respectively, to provide season-long white grub control. If applied that early, various other insect pest can also be controlled (Merit and Mach2: billbugs, annual bluegrass weevil, greenbugs; Mach2: cutworms, sod webworms) or at least suppressed (Merit: chinch bugs). If white grubs are the primary targets, the optimal application time for Merit and Mach2 is June/July when the female beetles are laying eggs. At this time, control efficacy against many white grub species is typically in excess of 90%. As the larvae hatch and go through their 3 larval stages, they become less susceptible to these insecticides (and other insecticides). Applications against the 3rd larval stage in September are not recommended.

Oriental beetle is very susceptible to Merit but MACH2 has only provided 50-60% control on average and should be applied right around peak egg-laying activity. Japanese beetle is very susceptible to Merit and Mach2, and even mid-September applications can still provide around 70% control. Applications after mid-August, however, may not kill the grubs quickly enough to avoid impending damage. Masked chafers are less susceptible to Merit, and where this species prevails, applications should be done during egg laying (June-July) and at the highest label rate. The European chafer appears to be less susceptible to Mach2 and Merit, and applications should be done during the egg laying period (June) and at the highest label rate. The Asiatic garden beetle appears to be immune to Mach2 and Merit.

The obvious disadvantage of preventative applications is that they have to be done before white grub populations can be estimated through soil sampling. Thus, preventative applications are often applied to areas that would need only

partial or no control at all. This increases the cost of grub management, resistance development, and may in the long-term dramatically reduce populations of natural enemies by depriving them of prey or hosts. Smart turfgrass manager will restrict preventative applications to high-risk areas, i.e., areas with extremely low damage threshold and tolerance, areas with a history of white grub infestations, and areas with high beetle activity (egg-laying) in June-July.

Curative white grub control

If soil sampling has revealed white grub populations, areas with densities above treatment thresholds or ongoing damage may need to be treated. This curative control approach works best if applied while the grubs are still smaller (i.e., mid August to early September). Monitoring and sampling helps optimize application timing and restrict treatments to areas that actually have high grub populations. Once the grubs have reached the 3rd instar, they are much harder to control. Spring applications (late April through May) are generally the least effective and rarely justified because the grass can outgrow most grub populations. Only extremely high grub populations, unduly stressed turf, or digging grub predators can cause damage at this time. Any necessary treatments need to be applied before the grubs pupate. Due to the annual life cycle of the grubs, areas that had no damaging infestation or were successfully treated in the previous late summer/fall, will not need treatment in the following spring.

Continued on next page...

“Total Control”^{Inc.} Athletic Field Management

*Your Fields require it
Your leagues deserve it*

- **On site staff training**
 - Program development
 - Equipment calibration
 - Application technique
 - Infield maintenance
- **Quality applications**
 - Fertilizer
 - Pest control
 - Core aeration
 - Slice seeding

Call for complete service description
Licensed **Jim Hermann**
Insured **(908) 236-9118**

Athletic Field


Machinery Specialists

Serving Central and Southern New Jersey

FINCH
SERVICES, INC.



JOHN DEERE



419 Industrial Drive - North Wales, PA 19454 - 215-661-0390 - www.finchinc.com

For best results with any insecticide, mow the turf and rake out dead grass and thatch before treatment. This will reduce the amount of insecticide bound up by surface debris. Irrigate with 0.5-1" water immediately after treatment (or timely rainfall) to leach the insecticide into the root zone where the grubs are feeding. Irrigation also increases insecticide contact by drawing the grubs closer to the surface. If the soil is very dry, pre-treatment irrigation 1 day before treatment will also increase efficacy by bringing grubs closer to surface and reducing thatch binding and evaporation of liquid treatments. However, do not apply soil insecticides to saturated soil. Also, granular formulations need to be applied to dry grass to allow the granules to sift down into the thatch. Liquid and granular applications are usually equally effective, however, granular formulations may be more forgiving if post-treatment irrigation is delayed.

Successful treatments typically kill 75-90% of the grubs but product performance varies with soil type, thatch thickness, and grub species. Therefore, evaluate treatments and keep record of product performance. While speed of kills varies with insecticides, soil insecticide applications never work overnight. Affected grubs usually turn yellow or brown within a week of treatment. Wait at least 1-2 weeks before evaluating. But don't wait longer than 3 weeks to allow for a follow-up treatment if the 1st treatment was ineffective. In the latter case, don't apply the same product again at a rate exceeding the label rate. Rather try a different compound. While development of

grub resistance to insecticides is unlikely with the presently used short-residual insecticides, some grub control failures can be caused by enhanced microbial degradation of the insecticide, especially after repeated insecticide use. Avoid unnecessary applications and alternate insecticides.

The range of insecticides available for curative white grub control has already and will continue being effected by the implementation of the Food Quality Protection Act of 1996. Among the organophosphates, only trichlorfon (Dylox) and diazinon (Diazinon) are still available. However, Diazinon production and retail sale is supposed to stop in August 2003. Diazinon may also not be very effective against Japanese beetle grubs. Of the carbamates, only carbaryl (Sevin) is still available but generally does not seem to provide good white grub control. Presently available nematode products for grub control contain the species *Heterorhabditis bacteriophora*, *Heterorhabditis megidis*, or *Steinernema glaseri*. These nematodes can be very effective against Japanese beetle and masked chafer grubs, but are not effective against grubs of oriental beetle, Asiatic garden beetle, or European chafer. While these nematode products have to be handled and stored with more care than chemical insecticides (you are dealing with living organisms!), they have the advantage of no reentry interval due to their non-toxicity. Use of the halofenozide (Mach2) and imidacloprid (Merit) is generally not recommended for curative control. While they

Continued on page 15....



A Division of Pennington Seed, Inc.

Scott Rushe
Account Manager

LOFTS SEED
9327 US Route 1
Suite J
Laurel, MD 20723

Office: 800-732-3332 Ext. 418
Home Office: 814-234-7930
Fax: 814-234-1914
Cell: 814-571-1039
E-Mail: srushe@penningtonseed.com

Where Great Grass Begins™

"Monthly Field Tip"

Scout for insect pests before turf damage occurs

by Jim Hermann

1. **Visual inspection:** The most obvious and simple technique is to get down on your knees and separate the turf with your hands and inspect for insect populations and also damage to grass blades and stems.
2. **Disclosing solution:** If you don't see any pests they may be hiding in the thatch layer or run too fast for you to catch. Mix two tablespoons of liquid detergent with a gallon of water and sprinkle it over a square yard of turf. Within five minutes, cutworms, webworms, beetles, and even earthworms will come to the surface if they are present.
3. **Flotation:** Insects such as chinch bugs can be discovered by using a two-pound coffee can. Cut the top, bottom, and lower rim off the can and drive the sharp edge of the can into the ground being careful not to cut yourself. Fill the can with water and stir up the turf. Within minutes you can easily count the insects, which have floated to the surface. Refill when necessary and count three minutes. *Cont pg 15...*



Shearon Sports

A division of

Shearon Environmental Design Co.

Specializing In:

Turf Management Programs

Athletic Field Renovations

New Field Design & Construction

Agronomic Consultations

Irrigation Design & Installation

Drainage Design & Installation

337 Route 31 · Hopewell, NJ · (609) 466-0666
902 Cooper Road · Voorhees, NJ · (856) 424-1727
5160 Militia Hill Road · Plymouth Meeting, PA · (610) 828-5488
www.shearonsports.com



THE PRO'S CHOICE

TERRE
TURF SUPPLIES
SINCE 1925
The Pro's Choice

Grass Seed
Fertilizers
Insecticides
Nursery Stock
Fungicides
Herbicides
Mulches
Tools
Stone
Soil
Ties

The TERRE Co

206 DELAWANNA AVE CLIFTON NJ 07014

Hardgoods - 973-473-3393

fax # 973 473 4402

cell # 201-390-6025

John Salisbury Sports Field rep.

Keep Your Ball fields safe and looking great!! TERRE has the products from grass seed and fertilizer to infield clays and turfage products
Purchase 1 pallet of turfage get free field evaluation done.



This is where we draw the line.

Sherwin-Williams is your one source for high-quality coatings, equipment and advice. You get fast order turnaround, competitive prices and easy credit terms, plus all of the innovations that make your job easier. We'll help you with on-site delivery, spec assistance, custom color matching and technical support. With over 2,500 locations, coast to coast or across town, you'll benefit from the powerful distribution and local service that can only come from Sherwin-Williams. When it comes to our customers' needs, on the field is the only place we draw the line.

To learn more, visit the Sherwin-Williams location nearest you, contact your Sherwin-Williams representative or call 1-800-524-5979 to have a representative contact you.



The Pros Know. Ask Sherwin-Williams.™

“Ask The Grub Oracle”

by Dr. Albrecht M. Koppenhöfer

Dr. Albrecht Kippenhofer is an Assistant Extension Specialist in Turfgrass Entomology, Cook College/Rutgers University

Ask Dr. Koppenhöfer questions concerning your insect problems by e-mailing him at HQ@sfmanj.org

Question: Is there any factor other than historical data that would cause a turf site to be considered a high-risk area for grub infestation?

Answer: Grubs prefer soil in the range from sandy loam to loam. Female beetles have a preference for nice lush looking turf and moist soil when searching for egg-laying sites and generally seem to prefer sunny areas. The eggs have to absorb water from the soil in order to develop. They actually almost double in size in the process. Similarly, the first instar grubs are much more prone to desiccation than the bigger ones. So, overall sites that are moist during egg-laying periods are more prone to get higher grub populations.

For species in which the adult beetles do a lot of feeding, the presence of preferred food plants also increases the risk of infestations because the females will often lay eggs soon after having fed. Adult Japanese beetle feed on > 300 plant species but lindens, purple leaf plums, roses, Japanese and Norway maples, bracken, elder, grape, Indian mallow, multiflora rose, sassafras, smartweed, and Virginia creeper are among the most attractive food plants. Adult Asiatic garden beetle feed on > 100 plant species but have a preference for box elder, butterfly bush, cherry, peach, rose, strawberry, sumac, viburnum, and various flowers such as aster, chrysanthemum, dahlia, goldenrod. In addition, Asiatic garden beetle like to seek shelter in orange hawkweed and sorrel.

Masked chafer grubs are fond of decaying organic matter and prefer moist soils high in organic matter. Green June beetle grubs feed mainly on decomposing organic matter and infestations are more likely where manure fertilizer or compost has been applied. ▲

Continued from page 10... "White Grub Management"

may still provide good overall curative control depending on grub species (see above), their speed of kill is too slow to prevent impending turf damage. ▲

¹*Assistant Extension Specialist in Turfgrass Entomology, New Jersey Agricultural Experiment Station, Cook College, Rutgers, The State University of New Jersey, New Brunswick, NJ 08901.*


Continued from page 13.... "Monthly Field Tip"

- 4. Turf/soil plug samples:** See the description given on page 10, under the heading "Early detection, sampling and monitoring, damage thresholds". In Dr. Koppenhofer's article on **White Grub Management in Athletic field Turf**.

(Turfgrass Insect & Mite Manual, David Shetlar et. al, PhD, 1990)
The preceding scouting techniques are only a few of the many techniques available to the turf manager for the identification and diagnosis of turf insect problems. Always identify, quantify and justify the need for any pesticide application.

▲

Sports Field Managers Association of New Jersey



Athletic Field Paint

White or Colors

Bruning

Benjamin Moore

Coronado

Till's Own

Field Marking Equipment

Fox Valley Systems

Newstripe

Jaydee Equipment



David Fitzgerald Sr.
Till Paint Company
691 S. Evergreen Ave
Woodbury Hts. NJ
1-800-845-5929
Fax 856-848-1084



NATIONAL SEED PROFESSIONAL TURF PRODUCTS

Specializing in Quality Grass Seed
To Meet All Your
Turf Performance Standards

Call For a Catalog

1-800-828-5856

Carrying a full line of quality mixtures
especially formulated for:

**SPORTS & ATHLETIC FIELDS
LOW MAINTENANCE AREAS
GENERAL GROUNDS
GOLF, LAWN, RECLAMATION**

Technical Agronomic Support
And Custom Blending Available