

March/April 2004 <u>Vol. 4, No. 2</u>

e-mail: hq@sfmanj.org www.sfmanj.org

Don't Miss – SFMANJ Spring Field Day

Where: Drum Point Sports Complex, Brick NJ When: April 20, 8am-8:30 registration Till: 3pm Box lunch included • Door prizes • No vendor booths

"When it's Built it's too Late." This day promises more than education. Come see a beautiful newly built 60 acre facility in different phases of construction. This is a wonderful park with the same construction problems we face everyday. Learn from Ken Mathis, Parks Director, how to avoid some of these problems and how to build a successful facility. You have the opportunity to see the final changes needed before opening day.

The facility includes 4 softball, 12 soccer and 2 football fields including the beginnings of a skateboard park and bike trail. We will show you how the maintenance building was constructed, how Ken worked with the engineer to get what he wanted. While you are there check out the pesticide storage area. See how to install an internal soil drainage system to combat drainage problems on athletic fields. In between the tour stop at each field and learn how the irrigation system and well work. (Every field is irrigated). We will explain some construction problems and how to develop a playing surface with the existing turf, each field is in a different stage of development. Find out how Ken amended his soil and see the difference between the fields and common areas. Check out the weeds and find what they are and out how to get rid of them. Stop by the skinned infields, see the problems, learn how to correct them. Learn how to analyze the mix and compare it to the ASTM standards as we whip the field into playable quality. Stick around afterwards for door prizes and questions. Watch for the flier with registration coming in the mail soon. Be the first to fax the enclosed puzzle with the correct answers and win a free admission. +

SFMANJ Business

Next Board of Directors Meeting – March 11, Thurs. 4pm. At Storr Tractor Co., Rt. 22, Somerville.

If you work for a professional facility and are interested in serving on the Board of Directors of SFMANJ fax a resume to 907-730-7770. You must be a member in good standing. •

Inside This Issue

Board of Directors	3
SFMANJ New & Renewed Members	3
Membership Form	3
Calendar of Events	. 4
Head Groundskeeper	. 5
Did You Know	8
Rutgers Corner	. 9
Puzzle Contest	.11
Movement of Nitrogen	12
Ask the CFSM	14
Field Tip	14

This newsletter is the official bi-monthly publication of the Sports Field Managers Association of New Jersey. For information regarding this newsletter, contact: SFMANJ at 730-7770

> *Co-editors:* Jim Hermann, CSFM and Eleanora Murfitt-Hermann, CRS

SFMANJ does not necessarily support the opinions of those reflected in the following articles.

Lip Service, It's All Part of the Game

By Jim Hermann, CSFM

The elevations of the turf perimeter at the point of intersect between the infield skin and the turf is the reference point from which all infield elevations are calculated and or maintained. By maintaining the turf perimeter of the infield at a specific plane and minimizing the accumulation of material known affectionately by sports field managers as the "lip", you help to maintain the integrity of your surface drainage plan. A properly maintained perimeter will always provide for a smooth transition from the infield into the turf area. As this battle goes, so goes the war.

continued on page 6



THE CHOICE OF THE PROFESSIONALS

Athletic Field Construction & Drainage Specialists

> Site Evaluations Design/Build Laser Grading Renovations

Specializing in Sand/Slit Drainage Systems

235 County Line Road Amityville, NY 11701 631-691-2381 Fax: 631-691-2392

Welcome New & Renewed SFMANJ Members

Currently we have 330 members. If you did not see your name in last months or this newsletter please call (908)730-7770 to see if you renewed your membership. This year you can not renew with the National STMA chapter if you did please call us. Take advantage of the Spring Field Day discount by renewing today.

Charlie Ralph Albanir James Alberti Bill Almendinger Wayne Atkinson Victor Barone Tom Barton Eugene Bellusci Vincent Benenati Robert Beutel Nancy Bigos Vincent Bono Louis Bosco James Bradbury Paul Brandon Bob Buono Jack Campbell Frank Cannella Jr. John Carbone Jr.

...

Mulch Pro Inc. Middlesex County, Dept. Parks Holmdel Township Board of Ed. Piscataway Twp. Board of Ed. Cranford Board of Education Weehawken Township Berkeley Heights Township Ridgewood Village Tenafly Board of Education Tenafly Borough Ridgewood, Village of College of Staten Island Philadelphia Turf Co. Manalapan-Englishtown BOE Finch Turf Equipment Tri State Athletic Field Edison Board of Education Eatontown Borough Morris & Bergen Cnty Irrigation

SFMANJ Annual Membership Registration Form

* receive update information by email

Title
Employer
Address
City
State Zip
County
Phone Fax
*E-mail
Signature
Individual\$35 Associate\$35 Organization/Institution\$35 Additional member from facility\$20
Commercial/Contractor/Vendor/Supplier \$85 Additional member from company \$25
Student \$10
Send with Check or voucher to: SFMANJ P.O. Box 370 Annandale, NJ 08801

2004

SFMANJ Board of Directors

President Eleanora Murfitt-Hermann, CRS Washington Twp Parks & Recreation

Vice PresidentJim Hermann CSFM Total Control Inc

Secretary Fred Castenschiold Storr Tractor Company

Treasurer Dean Marzocca Dean's Lawn & Landscape Co.

Directors

Dr. James Murphy - Rutgers University

Jeff Cramer - Plainsboro Township DPW

Clare Liptak - Rutgers University

Brad Park - Rutgers University

George McCarthy - Spring Irrigation Co., Inc.

Robin VanDerlyn, Chatham School District

Advisor.....Dr. Henry Indyk TurfCon GSI Consultants, Inc.

Mission Statement

Committed to enhancing the professionalism of athletic field managers in New Jersey by improving the safety, playability and appearance of athletic fields at all levels through seminars, field days, publications and *networking* with those in the sports turf industry.

Contact us at:

P.O. Box 370 Annandale, NJ 08801 Web Site – www.sfmanj.org E-mail – hq@sfmanj.org Ph/Fax – 908-730-7770

National Organization Sports Turf Managers Association

www.sportsturfmanager.com e-mail: SportsTMgr@aol.com Phone: 1-800-366-0391

Robert Carfagno Fred Castenschiold Dr. Richard Caton James Chimento Dr. Bruce Clarke Dave Coleman Matthew Conti Jeff Cramer Timothy Cronin Kevin Crossley Mario Cunha Robert Czumbil Dan Dandrea Gregory DeBuck Dennis DeSanctis Sr. John DeVries Jr.

Cranford Board of Education Storr Tractor Co., Inc. NJ Turfgrass Association JC Landscape Constr/Mangmt. Cook College Pathology Raritan Township Public Works Princeton University Plainsboro Township Ridgewood, Village of Ho-Ho-Kus, Borough of Cranford Board of Education **Ewing Township** Buena Regional School District DeBuck's Sod Farm of NY, Inc. Aer-Core, Inc. Hillside Gardens Inc.

continued on page 4

Calendar of Events

Rutgers University Athletic Turf Classes

- * February 24-26 Athletic Field Construction and Maintenance course
- * March 9th The Importance of Understanding Athletic Field Soil
- * March 16th The Importance of Understanding Athletic Field Turfgrass
- * March 23rd Understanding Athletic Field Construction Procedures For information call 732-932-9271 10% off for SFMANJ members

Annual SFMANJ Spring Field Day

April 20, 8am-3:00pm Drum Point Sports Complex, Brick NJ Box lunch included, Door prizes No vendor booths \$35 members, \$45 non-members For more info call 908-730-7770 Fliers will be sent soon.

New Jersey Recreation & Parks Assoc. Annual Convention & Trade Show

March 14-17 at Ballys in Atlantic City For more information call NJRPA at 732-568-1270

continued from page 3

Anthony Diaforli Joy Dobrosky Robert Druzsba Thomas Elder Ralph Ellis Salvatore Fama Will Fanner Thomas Fisher Dave Fitzgerald Sr. Donald Frederick George Frey Peter Galosi Scott Geier Sean Goodwin Joe Gourniak Dr. John Grande Todd Gritschke Michael Haener Erik Hammerdahl Rob Haynes John Helton Ken Henshaw Robert Hickey Luis Hidalgo Christopher Holenstein Barry Imboden Dr. Henry Indyk Kenneth Jenks

Princeton Regional Schools Better Materials Corp. Woodbridge Township, Parks Ewing Township Berkeley Township Sherwin Williams Co. Old Bridge Township Complete Lawn Service, Inc. Till Paint Co. Inc. Cranford Board of Education Fair Lawn Parks & Recreation Cinnaminson Board of Ed. Hawthorne Borough DPW Enviroscapes, Inc. Crop Production Services Rutgers Univ., Snyder Research Rahway Public Works Michael B. Haener, CID Chatham School District Union Township Cinnaminson Board of Ed. Atlantic County Vocational School North Colonie Central Schools Ramapo Indian Hills B. of Ed. Summit Board of Recreation Hunterdon Central High School Turfcon Storr Tractor Co.

Van Kampen William Koonz Tom LaRowe **TJ** Lawson Diane Leon Frank LoSasso Gary R. Lucks Robert Manning Thomas Martin Paul Martino Dean Marzocca Ron Matakitis William Mateyka Ken Mathis Larry Mayerowitz George McCarthy Brian McCormick Craig McCoy Brian McGuirt Bill Menagh Thomas Miller Chris Monohan Mike Moore John Mujica Dennis Murphy Art Neff Thomas Newbery Nat Nuovo **Richard Oates Bradley Park** J. Casey Parker, CPWM Tony Pavelec Jr. Wes Perrine Harold Pierce Ray Poerio Steve Polakowski Judy Policastro Rich Resavy Melissa Ripa Steve Riviello Rich Romanik Charles Romano James Saxton Michael Shannon Kevin Shipman Frederic Sibley Ed Sinclair Brent Sliker, CPWM Thomas Stokes James Stryker Mark Tindall Robert Tirserio Craig Tolley Steven Toth Robert Tranquilli Suz Trusty Walter Tucker Dawn Tuttle Scott Van Demark Barry Vansant David Ward Sr. George Warden Jay Weisenbach Betty Weist Wayne White Bruce Wild Greg Winfree Thomas Witt Thomas Wojcik Robert Young Ron Zaleski John Zambrano

Van Kampen Advertising, Ltd. Koonz Sprinkler Supply Delaware Township Rutgers University Leons Sod Farm, Inc. Hammonton Board of Ed. Lucks Sales Associates Piscataway Twp Board of Ed Pritchard Industries, Inc. Applied Landscape Technologies Dean's Lawn & Landscape Delbarton School Old Bridge Township Brick Township Middlesex County Parks Spring Irrigation Co. Inc. College of Staten Island Scotch Plains Township Cliffside Park Recreation Mendham Borough Environmental Resolutions, Inc. Scotch Plains Township Springfield Public School Cranford Board of Education Allendale Borough Croton, Village Of Rahway Public Works Field Pro Enterprises, LLC Ocean Cty.Utilities Authority Rutgers Univ., Plant/Pathology Lacey Township Pavelec Bros.Golf Course Cnst. Geo.Schofield Co. Inc. Winslow Twp. Board of Ed. Scotch Plains Township Seton Hall University Irrigation Association of NJ Hillsborough Township **DVH** Athletic Turf Moyer & Son, Inc. Millville, City of Cliffside Park Recreation Glen Rock Dept. Public Works Better Materials Corp. Pine Hill Public School Medway, Town of Mahwah Township Mansfield Township Cranford Board of Education Delaware Valley Reg.High Sc Cinnaminson Board of Ed. Glen Rock Public Works County College of Morris Montville Township Springfield Public School STMA Ocean City Bow School District Mahwah Board of Education National Seed Roxbury Township Middlesex County, Dept.Parks Glen Ridge Borough NJ Landscape Contractors As. Holmdel Township Board of Ed. Storr Tractor Co. **DVH** Athletic Turf Cranbury Township Finch Services Inc. Fair Lawn Borough Glen Ridge Borough Monmouth University

Head Groundskeeper - Newark Bears

Specific responsibilities include but are not limited to, the following:

- Responsible for all field maintenance including, fertilization, pesticides, mowing, turf areas, skin area (mound, Plate and infield)
- Manage pre game set up & post game break down of field operations
- Must have knowledge of sand based fields.
- Working knowledge of irrigation systems, sprayer, spreaders, and tractors and mowing equipment.
- Develop and manage relationships with outside vendors
- Negotiate prices and contracts
- Maintain inventories
- Order supplies
- Schedule all maintenance & repairs necessary for field through outside vendors.
- Hire, train and supervisor full-time hourly employees and game day staff.
- Establish and maintain budgets.
- Be a team player. Actively participate in other operational functions as necessary including but not limited to pulling tarp, attending all scheduled events and meetings.

To apply: Send or e-mail cover letter, resume and list of three professional references to: Dean Rivera, General Manager Newark Bears 450 Broad Street Newark, NJ 07102 drivera@newarkbears.com

You're Always Ahead of the Game with a COVERMASTER® Raincover...

"Great Service..., The Best..."

wrote Chip Baker, Asst. Baseball Coach, Florida State University, Tallahassee, FL

Chip's comments confirm what we hear from the many groundskeepers who use a COVERMASTER[®] raincover to keep their fields dry and ready for play.

Call us and we'll gladly tell you more.

The COVERMASTER® Advantage...

- Superior in strength and UV resistance
- Outstanding heat reflective properties
- Light weight easy to handle
- Widest materials for least number of seams
- Largest choice of weights and colors
 Backed by truly dependence and colors
- Backed by truly dependable warranties

TARP MACHINE VIDEO!

Call, fax or e-mail for a free video, material samples and a brochure.





fields are also readily available.

TARP MACHINE[™] lets you roll the cover on and off in minutes.

TARPMATE™ roller comes in 3 lengths with safety end caps.

тм



CALL TOLL FREE 1-800-387-5808

Covermaster.com E-MAIL: info@covermaster.com



COVERMASTER INC., 100 WESTMORE DR. 11-D, REXDALE, ON, M9V 5C3 TEL 416-745-1811 FAX 416-742-6837

continued from page 1

A common mistake committed by many sports field managers is to add infield mix to compensate for an improperly maintained lip.

Lip build up is caused by many factors and is to a large degree site specific in nature and severity. In this article we will consider some of the causes and controls.

The most obvious cause for lip buildup is the movement of infield mix from the infield into the grass perimeter. Improper grooming technique is the most limiting factor in proper lip management. Both wind and water erosion are also contributing factors. Game play is another obvious factor in particle migration and therefore lip buildup. The buildup caused by game play is most obvious at the turf adjacent to first and third base.

There is clearly no maintenance technique capable of preventing material movement within the confines of the infield skin. However, there are considerations and cultural practices, which can help to minimize this movement and its negative effects on safety and playability. There are in addition, procedures that can manage the lip buildup that does occur.

Most importantly, prior to all other maintenance procedures, the infield turf perimeter should be cleared of any infield mix that has accumulated since the last maintenance. This maintenance is essential prior to rain since rain has the ability to wash the mix down into the thatch layer where removal is much more difficult if not impossible.

Procedures available for this purpose are blowing, power washing and brooming. In more severe circumstances the use of a thatch rake is sometimes effective. In view of the fact that you can't keep the infield mix from moving to the turf perimeter, it is sometimes beneficial to move the turf perimeter away from areas of concentrated disturbance caused by game play. By cutting out existing turf and increasing the distance from first and third base to the outer perimeter of the infield, the amount of mix that is deposited into the turf can be greatly reduced. There are different designs conducive to this concept. The number of options is limited only by your imagination.

Grooming technique is most limiting and therefore first on the list of preventative maintenance considerations. Always rake parallel to the foul lines and turf perimeters. When dragging the infield, always stay 6" from the turf. Vary your dragging pattern. Alternate your starting and stopping point. Never contaminate the turf with infield mix for any reason.

The amount of moisture contained by an infield mix, while being maintained within maximum and minimum limits could be considered the glue that holds an infield together and as such is a factor in lip management. The key is to determine these limits. The limits will vary based on site-specific factors. The most important factor to be considered when addressing moisture management is particle size and distribution of your infield mix. What is the physical analysis of your infield mix? What is the sand, silt and clay particle size analysis? Would your mix be considered a sandy mix or would your mix be considered a clayey mix?

The ASTM Standard Guide for Construction and Maintenance of Skinned Areas on Sports Fields has provided guidelines to help in identifying and classifying your particular mix. In general a mix containing 70%-85% sand



size particles and containing 15%-30% clayey mix is considered an acceptable product. The sandier a mix is, typically the less stable it is given the affects of game play. The higher the percentage of sand a infield mix contains, the more difficult it is to maintain moisture at levels sufficient to promote stability.

The more clayey an infield mix is, the more that mix tends to retain moisture and the more effective moisture becomes as a means of stabilization. Let me repeat; this moisture is only beneficial when maintained between site-specific limits. Most all of us are aware that a clayey mix usually takes more time to condition after a heavy rain than a sandy mix. The benefit derived from the ability of a clayey mix to retain moisture is lost in this circumstance because the level of moisture has exceeded the limits of potential benefit.

It should be understood that a mix on the clayey side of the ASTM standard that is allowed to become very dry is somewhat difficult to rewet. Tilling or some other means of cultivation is sometimes necessary as a part of the wetting procedure.

When a level of maintenance is reached which allows for the "scheduled" periodic application of water, a soil amendment such as calcined clay may be beneficial in extending the duration of time between water applications. It must be understood that these products constitute the potential for a double-edged sword.

As was addressed earlier, a sandier mix or a mix with more sand "sized" particles is less stable and is more likely to migrate given the effects game play. If moisture levels are permitted to vary beyond the limits of potential benefit (either too wet or too dry), these products will display characteristics similar in nature to sand. When allowed to dry out, a mix that has been modified with an amendment such as calcined clay will have characteristics similar to a sandier mix, which does not have an amendment added. If allowed to become saturated, a mix amended with a product such as this will move in much the same way as a sandier mix.

For the purposes of this article the characteristics of a dry infield mix are determined almost solely by particle size and nothing more. When discussing the stability of an infield mix, a dry calcined clay particle will differ little from a dry sand particle given the same particle size. If anything, the clay particle will be more mobile due to being lighter and maintaining a lower bulk density. The benefits of products such as calcined clay are only realized in their ability to absorb moisture and aid in maintaining moisture levels between site-specific maximum and minimum limits.

I have in the past used the following comparison to explain the affects of moisture on a sandy infield mix. When walking along the beech an observation can be made. Up on the beach where the sand is dry the conditions are very unstable. You sink into the sand. As you approach the waterline, the sand has more moisture content and as such gains stability and firmness. As you enter into the water and the sand becomes saturated it again looses stability. The moisture in the sand provides stability only between maximum and minimum saturation levels.

Wind erosion is a subtle culprit that can slowly but surely eat away at the integrity of your infield. It is obviously site specific based on the severity and consistency of the wind. As with any erosion problem (wind or rain) wind erosion impacts on the smaller and or lighter particles. For this reason wind erosion has the potential to erode the silt, clay, fine sand and or added amendments from your infield and deposit this material at the turf perimeter adding to the problem of lip buildup. Along with adding to lip buildup, if allowed to persist, wind or rain erosion will destroy the integrity of a clayey mix and leave you with a sand box.

Controls would include providing a windbreak to minimize wind velocity. This can be incorporated into the permanent perimeter fencing. It can also be provided as snow or silt fence utilized during the off-season. If snow or silt fence is utilized as a windbreak during the off-season, remember to keep it away from the turf on the down wind side of the field. If a windbreak is installed to close to the turf it will cause airborne particles to drop right into the turf. As can be observed by the effective use of snow fence in winter storm management, drifting occurs on the downwind side. Maintaining moisture levels within the mix will increase stability of the mix and also minimize erosion.

When discussing erosion of a specific infield mix caused by water (rain), two major factors contribute to the severity of the problem. These two factors are water volume and velocity.

The more water there is and the faster it travels, the more severely it impacts on the stability of the infield mix.

First, consider water volume. The volume of water is the amount of water you are dealing with. Although you cannot control the amount of rain you receive, there are a number of ways to control the volume of water that travels within the confines of an Infield.

1. Cover the infield when it rains. For most of us this is an impossibility

2. As water travels along a linear path it increases in volume. Limit the distance the water travels before exiting the infield and you limit the accumulated volume. By properly grading the infield, you can direct water the shortest distance to the perimeter thereby limiting the volume of water. An example of one such grading plan would be to maintain the pitchers area as the high point of the infield and slope the infield to the perimeter with all bases being approximately level to one another.

3. Limit the concentration of water in specific areas when exiting the field. An example of the very worst grading design which encompasses the very worst of examples #2, #3 and #4 would be a skinned infield with home plate as the low point of the entire infield. As



Only Rain Bird rotors feature Rain Curtain™ Nozzle Technology that delivers uniform water distribution across the entire radius range for green grass results. Gentle, effective close-in watering around the rotor eliminates dry spots without seed washout, and larger water droplets assure consistent coverage, even in the windiest conditions. Install Confidence. Install Rain Bird.



water is funneled to a central location potential volume is increased and therefore the potential severity of erosion is increased. This problem is also magnified due to the distance the water has traveled in order to exit the infield.

4. Eliminate the potential for water to enter the infield from other areas during episodes of rain. If the outfield or foul territory is higher than the infield, water should be channeled away from the infield by some means.

The second player in this game of erosion is velocity or the speed of the water. Water increases in speed or "velocity" with an increase in slope. By minimizing slope you minimize velocity and therefore minimize erosion. Professional fields I have read about maintain around 1/2 % slope. This equates to approximately 51/2 inches of fall from the area at the base of the pitchers mound to the turf radius assuming a 90' radius. I prefer to maintain a slope of between 3/4 % and $1\frac{1}{2}$ % on the infields I take care of I believe, at less than 3/4 % there is too much potential for ponding and above

1½ % there is too much potential for erosion. These tolerances become more critical as the distance to exit increases. Remember, volume and velocity increase with distance as long as the supply (rain) remains constant

The last factor that contributes to the development of a lip that I stumbled on (literally) by accident is the combined affects of freezing and thawing along with the increased development of a thatch layer at the turf perimeter.

In November of last year I returned to an infield I had recently renovated to admire my work. I had completely resodded the perimeters of the infield and for that reason I was certain there was no lip. To my dismay a defined lip had developed within a period of days. The freezing and heaving of the very edge of the sod caused the lip. I believe this honeycombed soil structure provides an avenue for the inwashing of material from the infield. In addition to increasing the volume of soil within the lip, this modified root zone coupled with increased moisture supply at the perimeter of the infield promotes a localized environment conducive to the

Established 1925 **Fertl Soil** Specializing in Topdressing and Construction Soils For Golf Courses and Athletic Fields 548 Rosedale Road Kennett Square, PA 19348 610-444-0496 Brian I Parks of Moyer & Son, Plant Nutrients Offers professional sport turf care products. M.O. S. T. organic for full parks

M.O.S.T. organic fertilizers Moyer Green Gro granular sport turf fertilizers Fertigation & Turflo®Liquid Fertilizers Exclusive grass seed & erosion control products Soil amendments Clay drying materials

For Further Information Call 888-408-5433

development of a concentrated root system. With this proliferation of root development comes an increase in thatch layer and therefore an increase in elevation contributing to a lip.

Depending upon the severity of the lip, there are a number of ways to deal with it after it has established. The most aggressive procedure would be to use a sod cutter and remove the entire area of turf that rises above the desired elevation. The excess material that has accumulated below the sod is removed and the area is either resodded using the existing sod or new sod is brought in for the procedure.

A less aggressive approach to the problem is to dig a shallow trench adjacent to the turf lip and roll the lip into the trench. This procedure is most effective if the lip is very narrow and defined in relation to the desired elevation.

* A procedure that fills the gap between the least invasive (trench and roll) and most invasive (sod cut) procedures is to aggressively core aerate the area of lip, remove the cores and then roll the lip to the desired elevation. There must be enough volume of material removed through the aeration process to allow for the movement of remaining material without increasing compaction. The aeration procedure must penetrate deep enough to provide compaction relief 2" to 3" below the desired finish grade. There must be sufficient soil moisture available so as to allow for movement in the soil but not so much moisture so as to allow for smearing of the soil, which is in fact damaging to the soil structure.

*This idea was contributed by Brian Meola of Washington Township Parks & Recreation (Morris County).

Did You Know?

Phosphorous is the least soluble of the major turf nutrients and as such moves very slowly through most native soil root zones. For this reason much of the phosphorous applied, is not available to the turf roots in the year of application. In new construction, if soil test results report phosphorous as "low" availability, it is advisable to incorporate half of the recommended phosphorous into the root zone prior to seeding and topdress the balance. •

Rutgers Corner — Crabgrass control strategies for sports fields

By Brad Park, Rutgers University — park@aesop.rutgers.edu

"Crabgrass can grow on bowling balls in airless rooms, and there is no known way to kill it that does not involve nuclear weapons" – Dave Barry, Miami Herald

With spring soon to arrive, it is an important time to begin thinking about options for controlling crabgrass. If a significant soil seed bank exists and there are voids in the turfgrass stand which minimize competitive benefits of the turf, as a summer annual, crabgrass will germinate profusely in the spring, mature throughout the summer months, and die in early fall at the first killing frost leaving dead "skeletons' throughout the landscape. Crabgrass seed will typically begin germinating after April 10 in South Jersey and by April 20 in Central and North Jersey. Crabgrass will continue to germinate though mid-July.

Integrated Pest Management (IPM)

Recall that IPM attempts to reduce the risk that pest control strategies may have on the environment and people by incorporating all suitable techniques to maintain pests within acceptable limits. Although it is a common misconception, IPM *does not* entail the elimination of pesticide use.

Simply mowing at a cutting height suitable for the specific turfgrass species or mowing at a frequency such that scalping is avoided can constitute IPM. Improper mowing techniques leading to scalped turf will thin-out turfgrass areas, lead to voids in the stand, and subsequently provide opportunities for crabgrass to encroach. IPM also entails proper fertilization. Under-fertilizing turfgrass will often result in a weak stand, poor turf density, and an environment in which crabgrass can readily invade. Yearly nitrogen requirements per 1000 ft² for cool season turfgrasses used on New Jersey sports fields are: Kentucky bluegrass, 2-5 lbs; perennial ryegrass, 3-5 lbs; tall fescue, 2-4 lbs. High-use sports fields often necessitate the high-end of these nitrogen fertilization guidelines in order to encourage turfgrass recovery from traffic.

Preemergence herbicides: Are they an option?

For sports field managers whose cultural program includes spring overseeding of his or her fields, applying most preemergence herbicide products at the time of seeding will not

only deter crabgrass emergence, it will also inhibit establishment of cool season turf. Products such as p e n d i m e t h a l i n (Pendulum or Pre-M), benefin + trifluralin (Team), prodiamine (Barricade), oxadiazon (Ronstar), and dithiopyr (Dimension) are not viable

options for preemergence crabgrass control if overseeding is a part of the manager's spring program. Depending on the product and the application rate, the residual of these products is such that the seeding of desired cool season turfgrasses may not begin for 2 to 6 months following the application of the herbicide. Additionally, these products many not be used in

newly seeded turf as young turfgrass seedlings are highly susceptible to the phytotoxic effects of these herbicides.

Siduron

Siduron (Tupersan) is a herbicide that is labeled for preemergence crabgrass control in newly seeded Kentucky bluegrass, tall fescue, and perennial ryegrass. Tupersan is

formulated as a wettable powder and should be applied in the spring to coincide with maximum crabgrass germination. The label calls for either a

NATIONAL SEED PROFESSIONAL TURF PRODUCTS

Specializing in Quality Grass Seed To Meet All Your Turf Performance Standards

Call For a Catalog 800-828-5856

Carrying a full line of quality mixtures especially formulated for:

SPORTS & ATHLETIC FIELDS

LOW MAINTENANCE AREAS GENERAL GROUNDS GOLF, LAWN, RECLAMATION

Technical Agronomic Support and Custom Blending Available single application of product at 4.0-12.0 lbs/Acre or sequential applications at 6.0-12.0 lbs/Acre followed by a 4.0-6.0 lbs/Acre application 4 weeks later.

Postemergence herbicides

In order to use the chemical tools available to selectively treat crabgrass postemergence, the sports field manager must be able to accurately identify crabgrass at various seedling stages. Large crabgrass seedlings are characterized by upright growth and leaves that are rolled in the bud, lack auricles, and have a jagged membranous ligule. Large crabgrass leaf blades and sheaths are covered with stiff hairs. Smooth crabgrass is similar to large crabgrass, however it has fewer hairs on its leaf blades and sheaths.

Quinclorac and fenoxaprop

Quinclorac (Drive) and fenoxaprop (Acclaim Extra) are labeled for the selective postemergence control of crabgrass in perennial ryegrass, Kentucky bluegrass, and tall fescue. Quinclorac is effective in controlling young, un-tillered crabgrass seedlings and may be applied up to 0.75 lbs/Acre (1.0 lb Drive/Acre). To increase the efficacy of weed control, the label recommends applying quinclorac with an oil-based adjuvant such crop oil concentrate or methylated seed oil.

Quinclorac may be applied up to 7 days prior to the seeding of tall fescue, Kentucky bluegrass, and perennial ryegrass, at the time of seeding for perennial ryegrass and tall fescue, 7 and 14 days after the emergence of tall fescue, and 1 month after the emergence of Kentucky bluegrass, perennial ryegrass and tall fescue. The label notes that adjuvants should not be added to quinclorac applications to newly seeded turf prior to 28 days after seedling emergence.

Fenoxaprop may be applied at rates ranging from 0.016-0.17 lbs/A (3.5-39.0 fl. oz Acclaim/A) depending on the stage of crabgrass growth and established turfgrass species. For example, 4-5 tiller crabgrass may be treated with fenoxaprop at 0.17 lbs/A (39.0 fl oz Acclaim Extra/Acre) in perennial ryegrass and tall fescue whereas no more than 0.12 lbs of fenoxaprop (28.0 fl oz Acclaim Extra/ Acre) may be applied to 3-4 tiller crabgrass in Kentucky bluegrass turf.

Following applications of fenoxaprop, tall fescue and perennial ryegrass may be seeded immediately.

Following germination of tall fescue and perennial ryegrass, fenoxaprop should not be applied until seedlings have matured for 1 month. Of the cool season turfgrasses used on sports fields in New Jersey, Kentucky bluegrass is the most susceptible to phytotoxic effects associated with fenoxaprop. For example, when utilizing fenoxaprop rates greater than 0.04 lbs/A (9.0 fl oz Acclaim Extra/A), Kentucky bluegrass seedlings must be at least 3 growing months old before fenoxaprop can be applied. Additionally, 21 waiting days should be allowed following the application of fenoxaprop prior to seeding Kentucky bluegrass

Due to the complexity of Drive and Acclaim Extra labeling with respect to crabgrass growth stage susceptibility, individual turfgrass species herbicide tolerances, and turfgrass seeding timings, pesticide labels *must* be thoroughly read and understood prior to the application of these materials.

Literature Cited

Hart, S. 2000. Crabgrass and goosegrass control in cool season turfgrass. Rutgers Coop. Ext., NJ Ag. Exp. Stn., E233. ◆





Puzzle Contest: Win one free admission to Spring Field Day on April 20th. Fill in the puzzle, be the first to fax the correct answers to 908-730-7770. Attached your name, address and phone.

Across

- 1. A measure of engine strength
- Resistance of a fluid to sheer force, ie oil weight
 A form of engine aspiration
- A form of engine aspiration
 A structure which protects an equipment
- operator8. Unit of electrical current used to quantify
- the capacity of an alternator9. Measure of hydraulic pump capacity
- The organization which develops standards
- 12. Measurement of pressure
- 13. Measurement of a battery's starting power

Down

- A transmission with infinite speed control
 Method of pest control that attempts to minimize yet not eliminates the use of pesticides.
- 4. Abbreviation for the program responsible for evaluating and reporting new turf seed varieties.
- 5. The strength of an engine measured in foot pounds
- State department concerned with pesticide regulations.
 Measure of speed travel

ENVIRONMENTAL



• ENGINEERS • SCIENTISTS



Our firm is seeking a Landscape Architect to enhance the firms, planning, and design and construction management capabilities. We are a progressive group of professionals dedicated to client service. Our company is growing and we have recently expanded into a new Burlington County office in Mount Laurel, New Jersey.

We are interested in an individual with a degree in Landscape Architecture and a CAD/design program background. This person will aid in the development of park and recreation projects from concept through ribbon cutting. In addition, a primary function will be the review of development plans for municipal clients.

Our firm is seeking a motivated individual interested in emerging into a new area of the business and enabling this section to develop.

Send resume to Environmental Resolutions, Inc., 525 Fellowship Road, Suite 300, Mount Laurel, New Jersey 08054-3415, attn. Wanda Ford, EOE.

Movement of Nitrogen: Fertilizer in a Turfgrass System

From the January 1999 issue of Sportsturf - *by University of California-Riverside Research Team

Nitrogen (N) aids many plant processes and components. It's necessary for growth and development, appearance, and recuperative ability of all turfgrasses. However, its mobility makes N a potential environmental hazard. In nitrate form, N won't bind with soil or organic colloids. It can move from the application site to ground/surface water or the atmosphere by leaching and runoff, or by volatilization. Our study monitored N movement below the root system of cool-season turfgrasses. We looked at situations where N was applied at high rates and frequent intervals.

Methods

Turfgrass Research Project at the Agricultural Experiment Station of the University of California (U.C.) - Riverside provided study plots of mixed Kentucky bluegrass and perennial ryegrass.

We applied N at 2.5 lbs. per 1000 sq.ft. to Hanford fine sandy loam soil, and reapplied every eight weeks. We

sampled the experimental plots through two consecutive application periods, and performed nitrate analyses with a Technicon Autoanalyzer II.

We used a randomized, complete block of 4-ft. by 6-ft. plots, and performed three replications. Weekly mowing maintained a 2-in. height of cut, and clippings were collected to limit thatch. Sprinkler irrigation replaced soil moisture according to estimates of natural evapotranspiration.

Our nitrogen sources included granular urea (46-0-0), sulfur-coated urea (SCU: 37-0-0), and blood meal (13-0-0). These sources are classified as soluble, slow-release, and natural organic, respectively. They represent a range of nitrate-leaching potential. An untreated control balanced the study.

We collected two samples from each plot every week using Irrometer lysimeters. Samples of tap water from the irrigation source and deionized water



accompanied each batch of leachate samples.

Results

Granular urea provided the highest concentration of nitrate sampled. The concentration peaked 10 to 14 days after application. At no time did nitrate leachate exceed federal safety limits.

Sulfur-coated urea treatments demonstrated significantly less leaching of nitrate than urea during peak leaching times. SCU regularly showed more evidence of leaching than blood meal and the untreated control, but there was no significant difference among the three treatments at any rating date during the study.

Even at very high N fertilization rates, there was little probability of significant nitrate leaching from any of the tested sources. Only urea gave levels that were above tap water content, but these readings still fell below federal guidelines.

Discussion

Other studies found similarly low levels of N leaching. A Michigan State University researcher recovered less than 0.2% of applied N below the turfgrass root system. The N he detected was well below the drinking water standard.

A Nevada study reported a total leachate loss of 1.0% or less for tall fescue and bermudagrass turf, and another study at Cornell University found minimal N leaching.

In contrast, a Washington State University study found that nitrates could leach from newly constructed sand putting greens in golf course applications. In this creeping bentgrass study, leaching was strongly tied to N application rate, and was strongly modified by rooting medium and application frequency. N leached more from pure sand than from a sand-peat medium.

Leaching was much greater in the first year of the study than in the second, possibly due to more extensive rooting in the second season. Modified-sand rooting medium, moderate levels of total annual N, and frequent applications produced the lowest leaching loss (3-5% annually).

Studies show further that gaseous loss of N can be minimized by applying water immediately after application. This ionizes ammonia that can be produced by rapid mineralization, and prevents it from escaping into the air.

Gaseous N loss can also result when microorganisms chemically reduce nitrate. This produces elemental nitrogen and nitrous oxide gasses. Further research is necessary to explore this phenomenon.

Fertilizer nitrogen applied to a dense, mature, well-maintained turf is normally used rapidly by the turfgrass plant and soil microorganisms. There appears to be little chance of downward movement of nitrogen other than on pure sand with immature turf present. The following cultural practices help minimize potential leaching: • Water-in fertilizer immediately following application.

· Do not over-apply N.

• Use low application rates or slow-release sources on sands.

• Avoid over-irrigation directly after application.

*University of California Researchers Victor Gibeault, Marylynn Yates, Jewell Meyer, and Mathew Leonard contributed to the study. Their complete report is published by the University's Cooperative Extension in California Turfgrass Culture Vol. 48, Nos. 1 and 2. •





YOUR "ONE-STOP SOURCE"

March/April 2004

Ask the CSFM

Jim Hermann, CSFM is a Certified Sports Field Manager. Ask Jim questions concerning your baseball/softball fields. E-mail him at hq@sfmanj.org

Question: We just acquired a deep tine aerator last fall. We purchased both ³/₄" hollow and ¹/₂" solid tines. The problem is that the field is so hard and compacted we were not able to aerate below about 3". What should I do?

Answer: In the short term I would suggest deep solid tine aerating as early in the spring and as deeply as possible. In doing so you will be taking advantage of the natural compaction relief provided by the heaving action of the deep frost we have maintained this winter. This will allow for the most effective deep tine aeration possible given your situation. You may have to acquire larger diameter solid tines to achieve maximum depth without bending. Be sure to wait until proper soil conditions persist before attempting your aeration program. Use your soil test probe to pull core samples to confirm proper soil conditions. The soil should be moist but not so moist so as to ribbon when rolled between the thumb and forefinger. It should have the ability to crumble or separate into individual aggregates when pressure is applied.

If timing and usage permit, I would recommend follow-up core aeration in late May. By then, game play will have recompacted much of the playing area. Depending on the depth of your topsoil, I wouldn't recommend core aeration much below 3" or 4". Pulling cores from below this depth will typically bring inferior soil to the surface. Compaction from foot traffic normally does not impact on soil below this depth. In the long term, I would begin to develop a proactive aeration strategy, which anticipates the affects of traffic and seasonal weather patterns. •

Field Tip

Have any necessary infield mix delivered and placed on the infield while the ground is frozen. This will minimize unnecessary handling of material and also minimize the potential for unnecessary damage to surrounding turf areas caused by heavy trucks.

2004 Proud Sponsor Directory

TOTAL CONTROL, INC.

Athletic Field Management & Consulting Jim Hermann, **CSFM** P.O. Box 422, Lebanon, NJ 08833 Ph/Fax: 908-236-9118 jimtc@att.net

US ATHLETIC FIELDS, INC.

Sports Field Maintenance, Renovation & Construction John McKnight and Jim Gilligan P.O Box 38 – Skillman, NJ 08558 609-466-2846 Fax: 609-466-1808 jim@usathleticfields.com

GSI CONSULTANTS - TURFCON DIV.

Dr. Henry Indyk, Sports Field Consultant 732-247-8026

STORR TRACTOR COMPANY

Turf, Irrigation and Ballfield Equipment 3191 Highwqy 22 Somerville, NJ 08876 908-722-9830 Fax: 908-722-9847

WILFRED MAC DONALD, INC

Turf Equipment Specialists Bernie White – Sales Representative 19 Central Blvd., S. Hackensack, NJ 07606 888-831-0891 ex 114 Fax: 201-931-1730 sales@wilfredmacdonald.com

SPORTS TURF SYSTEMS BY SAUL BROS.

Drill & Fill, Deep Tine Aeration Jerry Saul & Danny Saul P.O Box 299 Livingston, NJ 07039 973-983-1141 Fax: 973-983-8845

MENDHAM GARDEN CENTER

Turf Products Mendham – 908-543-4178 Chester – 908-879-5020 Annandale – 908-730-9008

PUT YOUR AD HERE: To become a Proud Sponsor Call 908-730-7770 \$150 FOR ONE YEAR



Storr Tractor Company

Distributors of Commercial Turf Equipment & Irrigation New Jersey State Contract Vendor



The Groundsmaster 4100-D is the most powerful mower in its class with high quality cutting performance.





3191 Route 22 Somerville, NJ 08876 908-722-9830

> Serving the Industry Since 1945



P.O. Box 370 Annandale, NJ 08801

STANDARD MAIL U.S. POSTAGE PAID LEBANON, NJ 08833 PERMIT No. 13

Or Current Occupant

Bradley Park Rutgers University Dept. of Plant **Bio/Pathology** 59 Dudley Road New Brunswick, NJ 08901



Experts on the Field, Partners in the Game.



Wilfred MacDonald is your team when it comes to athletic field equipment. We offer a wide variety of equipment from striping reel and rotary mowers to athletic field conditioners, line stripers, groomers, top dressers, aerifiers and more! Our comprehensive line of Jacobsen, Smithco, Turfco, National and Vertidrain gives you the largest variety of turf equipment to choose from! Contact your sales representative today for a demonstration!

















Wilfred MacDonald, Inc 19 Central Boulevard South Hackensack, NJ 07606 888-831-0891 www.wilfredmacdonald.com

Bernie White Mike Clifford **Tim Kerwin Mike Pelrine**

