testing

Oregon certification standards require 98% pure seed and 85% germination with minimum allowance for weed and crop seed. Penncross growers, through their association, have tightened these requirements even more. Putting green quality Penncross must meet the above standards plus be free of all objectionable weeds and crop seeds including Poa annua. This extra effort by the growers makes Penncross one of the purest turfgrass seeds available for putting greens.

Bentgrass seed is extremely small, with approximately 9-million seeds per pound. Since each seed produces a plant, the seeding rate for even new greens is light compared to larger turfgrass seeds sown on other areas of the golf course.

The general recommendation for seeding new greens is one to two pounds of seed per 1,000 square feet. For overseeding existing greens, one to one and a half pounds per thousand square feet is recommended. Rapid establishment of new greens is proven by letters from enthusiastic golf course superintendents who report newly seeded greens being in play from 60 to 90 days after seeding.

This bentgrass variety does receive scattered complaints from course superintendents who run into maintenance problems, such as the requirement for a tight mowing schedule. The hybrid qualities of Penncross require greater attention: brushing and verticutting are necessary to keep greens in top condition. When properly maintained, it will give that consistent texture required for true putting. The rapid growth also repairs the green from heavy traffic associated with today's courses.

The hybrid characteristics of Penncross testify to the ability of Penncross to crowd out Poa annua from greens. Where Poa is a problem, overseeding with this bentgrass on a regular program will indeed help. In fact, according to a letter from one golf course superintendent, it is the only cure he knows for ridding greens of Poa annua.

Penncross is deep rooted, a quality which must be considered in a golf green grass. This factor makes it both cold and heat tolerant. From the rigors of the Canadian and Alaska winters word comes to the Penncross Association that it came through in great shape. Conversely, golf course superintendent in а California reports an irrigation system breakdown for one week in over 100 degree weather and still the Penncross emerged with no visible harm.

The consistent demand for Penncross seed is evidence of its acceptance throughout the golf world and speaks well for this very special polycross developed by Dr. Musser and Dr. Duich. The Penncross Association has adopted the slogan, "the grass designed for golfers" and the reputation speaks well for that slogan []

Fertile Tiller Increase Noted By Open Field Burning

Open field burning after harvest has helped control disease and weeds and produce better yields in grass seed crops in Oregon's Willamette Valley since the practice began in the mid-1940s

How does the heat improve yield? Oregon State University crop scientists, working with red fescue, have found part of the answer. The answer is important because Willamette Valley growers, who produce 70 percent of the U.S. supply of red fescue and a large part of the world's supply of other grass seed, face a January 1975 deadline. That is when open field burning is scheduled to be banned in Oregon because of air pollution problems.

Dr. David O. Chilcote, OSU crop physiologist, recently reported on the red fescue results at the annual meeting of the American Society of Agronomy.

"The primary response to postharvest residue burning was an increase in the number of fertile tillers (panicles)," he said.

"Not surprising is that competition for sunlight in the fall is one of the keys to plant growth and yield. With field burning, the plant does not elongate, seeking the sunlight. The shorter plant has more tillers in the fall and the tillers produce more seed heads in the spring."

In tests conducted by graduate student Phillip C. Stanwood, plants in a burned area produced 16 tillers per 10 primary shoots in the fall. The rate for the unburned area was one tiller per 10 primary shoots.

"Contrary to the belief that burning causes a shock in plants which stimulates growth, we found that burning, in addition to opening up the stand of plants, also exposes the soil more, so the plant roots and shoots experience a wider range of temperature each day." Chilcote said. This may favor floral induction.

Red fescue was picked for the plot experiments because it is particularly sensitive to burning in terms of seed yield. Similar results were produced in more limited tests with tall fescue, perennial ryegrass, bentgrass and bluegrass.

The importance of sunlight was established by shading burned and unburned plots through the winter. The shade, acting much like stubble left in an unburned field, stopped seed production.

"Our evidence points to light competition as being problably the most important single factor related to crop response and post harvest burning," said Chilcote.



For More Details Circle (128) on Reply Card