The Turfgrass Seed Market

IMPROVED TURFGRASSES

All grasses are Monocotyledons belonging to the family Gramineae. They are constructed of narrow and parallel-veined leaves which grow from a hollow stem, the culm.

The grass family is one of the most numerous and most important plant species to man, with more than 600 genera and 10,000 species.

All cool and warm-season grasses considered turf type today originated outside of the United States. Most cool-season grasses were brought to North America from Europe during colonization, including bentgrass, all fescues, ryegrasses, and bluegrasses. Even crabgrass came from Europe. If you are wondering what grasses are indigenous to North America, they include buffalograss, reed canarygrass, and blue gama.

Warm-season grasses had the following origins; Bermudagrass -Africa, Centipedegrass - S.E. Asia, Zoysia - Asia, St. Augustine - Africa, Bahiagrass - Brazil, and Kikuyugrass - Africa.

Turfseed selection and development is a building process. First collections of turf and forage type grasses are made. Selections from these collections are the first level of improved turfgrasses. These selections are crossed to get first generation hybrids. The most recent group of improved turfgrasses are crosses of first generation hybrids. These are considered second generation hybrids.

The turf seed market has developed into an attractive market with adequate protection for proprietary turfgrasses. The additional push on development of new turfgrass varieties by private enterprise is causing an evaluation of all types of grasses for turfgrass use. The reclamation market and low maintenance turfgrass market are considering grasses which were



Glenn W. Burton Breeder of the "Tif" series of bermudagrasses and originator of radioactive mutation techniques for turfgrasses.

previously excluded from breeding and selection work. Now, all genera and species are being considered for a role in the turf market.

An outline of each genus applicable to turf are needed to grasp the overall scope of turfgrasses today.

COOL SEASON GRASSES

Bentgrasses

Bentgrasses, because of their importance to golf, received much of the initial attention of turf researchers. Bentgrasses are naturally low-growing and tolerate low mowing. Redtop and German creeping bentgrasses were common on golf courses and other turf areas prior to the 40s. Scotts used to sell German bentgrass stolons to golf courses in the 20's.

Much of the early work of the USDA and USGA Green Section involved bentgrasses. Pennsylvania State, the University of Rhode Island, Oregon State Agricultural Experiment Station, and Rutgers all contributed to selection and development of bentgrasses.

There are three primary types of bentgrasses used for turf today; creeping bentgrass, colonial bentgrass and velvet bentgrass.

Creeping bentgrass, Agrostis palustris, is very aggressive when fertilized and irrigated. It is propagated either by seed or stolons. Seeded varieties at one time exhibited a certain degree of segregation which would cause a green to look patchy. The problem has been resolved for the most part with the newer varieties. Improvements in disease suceptibility have also been made with newer varieties. The latest release is Penneagle (1979) developed by Joe Duich at Pennsylvania State University. Penneagle was evaluated for more than 20 years before its release. Penncross was released in 1954 by Penn State and suffered from lack of protection until Tee-2-Green Corp. was established to represent growers and to control purity in the mid-70's. Tee-2-Green also markets Penneagle.

International Seeds, Inc., distributes a Swedish creeping bentgrass called Emerald. It is known as Smaragd in Europe and is owned by W. Weibull in Sweden. Emerald was developed in Europe from progeny of Congressional, a vegetative variety developed in the U.S.

The oldest seed type creeping bentgrass marketed today is Seaside, selected by Oregonian Lyman Carrier and released in 1928.

Colonial bentgrass, Agrostis tenuis, is aggressive but has a lower tendency to creep. The last cultivar to be released in the U.S. was Exeter by Rhode Island in 1963. It too suffered from lack of protection and marketing effort, but will soon be repromoted by Pickseed.



1. Lime is applied to the fields after they have been turned under. Then the fields are plowed again and fine graded, very similar to soil preparation for lawn seeding.



2. Charcoal banding is a process where the seed and liquid fertilizer are layed in rows and sprayed with a band of charcoal covering the seed. The charcoal protects the seed from Karmex or Atrazine applied to kill weeds.



3. Grassland drills are an economical way of reestablishment of the seed crop. This can only be done if the fields are burned to get rid of the straw. Drilling is a one-step. process putting down both seed and fertilizer.

Northrup King distributes Holfior, a colonial bent developed in 1940 by D.J. van der Have of the Netherlands. Highland colonial bentgrass is a public variety released by the Oregon Agricultural Experiment Station in 1934. Production of this seed is carefully watched by an association of growers for quality and supply reasons.

Astoria was also released by Oregon in 1936. It does not have the winter hardiness of Exeter.

Velvet bentgrass, Agrostis canina L., has an extremely fine leaf which gives it a velvet appearance. It is less aggressive than creeping bentgrass but more aggressive than colonial. Velvet bentgrass is more tolerant of acidic soils than the other bentgrasses but prefers well drained and well aerated soil. New England is a prime area for use of velvet bentgrass in the U.S.

Rhode Island's Richard Skogley released Kingstown in 1963 as a public variety. It was the first velvet bentgrass released since the depression. Pickseed intends to market and promote Kingstown soon.

Redtop bentgrass, Agrostis alba L., is a coarse, stemmy bentgrass well adapted for use on poorly drained, infertile soils such as roadsides. It is occasionally used in mixtures for low maintenance areas.

Bluegrasses

Kentucky bluegrass is the most popular turfgrass in North America and as such has more cultivars available than any other turfgrass. The prime factor in bluegrass improvement was the discovery of apomixis, a characteristic which limits cross-pollination of some Kentucky bluegrass varieties. Kentucky bluegrasses which exhibit this characteristic were considered asexual and thus received patent protection prior to the 1970 Plant Variety Protection Act which provided protection to sexually propagated plants.

Dr. C. Reed Funk at Rutgers developed the technique to create Kentucky bluegrass hybrids that were apomictic. His work is part of nearly every Kentucky bluegrass developed since the mid-60's. Most other improved varieties are based upon collection and selection work.

Recently, International Seed, Inc., released a variety of rough bluegrass, *Poa trivialis*, called Sabre. Sabre was developed by Reed Funk and William Dickson at Rutgers for shady, moist areas where a lowgrowing bluegrass is desired and for southern overseeding.

Kentucky bluegrass, Poa pratensis L., is the backbone of turf in the Northern U.S. Releases of improved Kentucky bluegrasses began in 1947 with the official release of Merion. The emphasis behind improvement programs for Kentucky bluegrass was disease resistance, color, low growth habit, texture, and sod strength. Later winter hardiness, shade tolerance, traffic tolerance, establishment rate, and spring and fall color were added to the list.

Merion had been found as a naturally superior Kentucky bluegrass by Superintendent Joe Valentine at Merion Golf Club near Philadelphia in 1936. It has proven itself under low mowing and golfer abuse. Nature did the work, but it took Valentine to find it and the Green Section's Fred Grau and Pennsylvania State's Musser and Duich to refine and evaluate it for the market.

Other Kentucky bluegrasses collected and selected include the European varieties Fylking, Baron, Aquila, Parade, Sydsport, Cheri, Birka, Enmundi, and Rugby. U.S. varieties collected and selected include Warren's A-20 and A-34; Glade, a shade tolerant bluegrass found by Reed Funk in an Albany, NY lawn ; Ram I, which Green Section director Al Radko and Superintendent Ernest Brown found on his golf course in Maine; Scenic, picked out of a field of Merion by seed grower Otto Bohnert; and Newport, found by extension specialists in Oregon.

The selection process continues as an important contributor to new Kentucky bluegrasses. As the germ plasm base is expanded, the number of possible hybrids increases. From selections processed through Rutgers, Funk's team was able to develop the hybrids Adelphi, Bonnieblue, Majestic, America, and Bristol. Rutgers has been involved with the selections Glade, Touchdown, Columbia, Plush, and Brunswick.

O.M. Scott & Sons has developed the hybrid Merit and the selections Vantage, Victa and Windsor.

Dick Bailey, formerly a partner in Turf Seed and now working for Jacklin, found the selection Shasta out of a field of Pennstar. Shasta is considered a bluegrass specifically for the Northwest. Pennstar was a selection by Pennsylvania State. Sodco is a selection developed at Purdue.

SEED PRODUCTION



6. Spot spraying in the spring. The spray is a combination of Roundup and a red dye so one can tell what has been sprayed.



5. Fields are sprayed with selective herbicide in the fall to control broad-leaf weeds. Each swath of the sprayer is 40 feet wide.



4. Fields in early spring. Taller plants are volunteer ryegrass and will be spot sprayed in the spring

Fescues

There is currently a great deal of effort on fescue improvement and development. A number of companies are working to serve the transition zone with an improved tall fescue. Hard fescues are also being studied and two have been released.

There are five types of fescue used for turf; creeping red, chewings, tall, hard and sheep.

Creeping red fescue, Festuca rubra, is a fine leaved fescue which is often used in mixtures with Kentucky bluegrass and perennial ryegrass. It germinates more rapidly than bluegrass, but not as rapidly as perennial ryegrass. Fescues tolerate drought and infertile soil better than both Kentucky bluegrass and perennial ryegrass. Under dry periods on clay soils, the fescue may dominate the stand.

Much of the improvement in creeping red fescues has taken place in Europe. D.J. van der Have of the Netherlands developed Ruby and the Sports Turf Research Institute in Bingley, England developed Dawson. Both of these are marketed in the U.S. by Northrup King. International Seeds Inc. markets Ensylva. In 1954, Pennsylvania State released Pennlawn. Prior to that, Oregon State released Illahee. Pickseed markets Agram from the Netherlands.

A chewings fescue is one that doesn't creep. Chewings fescue, Festuca rubra var. commutata, originated in Europe, but much of the original production took place in New Zealand. In the 30's, much of the chewings fescue on the market came from New Zealand and suffered from poor germination. Rhode Island, Michigan, and New Jersey (Rutgers) experiment stations contributed to the improvement of chewings fescues. Rutgers developed Banner which is marketed by Burlingham. Rhode Island developed Jamestown from material found by Richard Skogley on an abandoned green in Jamaica. NY. Jamestown is marketed by Lofts. Wintergreen was developed at Michigan State from material from the Netherlands. Northrup King markets Wintergreen, and the European variety Atlanta. Turf Seed released Shadow chewings fescue in 1980. International Seeds markets Highlight, developed by Van Engelen of the Netherlands as a chewings type red fescue. Other wellknown chewings fescues are Golfrood from the Netherlands and Cascade developed at Oregon State University in 1964.

Tall fescue, Festuca arundinacea, is a coarse fescue with a bunch type growth habit. However, it has a deep root system and survives on infertile, salty, low maintenance areas such as roadsides and parks.

The primary varieties are Kentucky 31 developed by E.N. Fergus at the University of Kentucky in the late 40's, and Alta, developed in 1947 by Harry Schoth, an agronomist with USDA in Corvallis, OR. Fred Grau assisted in getting Alta planted on a number of highways to prove its advantages. Alta is not a preferred grazing grass for cattle.



C. Reed Funk

The first full-time turfgrass breeder in the U.S. serving the New Jersey Agricultural Experiment Station at Rutgers. Funk was the first to develop hybridization techniques for Kentucky bluegrasses.

Kentucky 31 is often used for turf in the transition zone due to its ability to withstand hot humid summers and acid soil. To provide turf managers in the transition zone with finer-bladed tall fescue, Loft has released Rebel tall fescue and Burlingham has released Falcon, developed in cooperation with Bill Meyer at Turf Seed.

For similar reasons, hard fescue, Festuca ovina var. duriuscula L. Koch, has received attention. In addition to good drought tolerance, it exhibits good shade tolerance. Northrup King markets Scaldis and Scotts markets Biljart developed in the Netherlands. Pickseed markets Tournament from the Netherlands.

Sheep fescue, Festuca ovina L., has fine leaf texture and exists on acid, coarse soils. It has good shade and drought tolerance but appears bunchy.

SEED PRODUCTION



7. Fields are cut and windrowed while seed is still green and less prone to shatter.



8. Whirlwinds can pick up the windrowed grass and carry it up to a mile contaminating nearby fields of other seed types. Whirlwinds and rain can seriously damage the crop in its last days of production.



9. Combines mechanically separate the seed from the straw by a series of rub bars incorporated in the thrashing machine.

Perennial Ryegrass

Next to Kentucky bluegrass, perennial ryegrass, *Lolium perenne*, has received the most attention from breeders and developers. This attention, however, wasn't attracted until the late 60's when new material from Rutgers and Penn State reached the market.

In the mid-60's, production of perennial ryegrasses rarely exceeded 150,000 pounds. Manhattan, Pennfine and a whole new series of perennial ryegrasses made turfgrass seed buyers take note, and in 1980 growers expect a crop of nearly 25 million pounds.

Bob Russell of Adikes is credited for the acceptance of perennial ryegrasses by northeastern golf course superintendents, sod producers and landscape contractors with his NK100 mixtures. The southern overseeding market was first broken by Northrup King and the other members of the Seed Production and Introduction Corp. (SPIC). Lofts, Turf Seed, Pickseed West, and International Seeds have southern overseeding mixtures available also.

Perennial ryegrasses are sexually propagated by crossing and polycrossing. This is similar to bentgrasses in that various parents are grown together in the seed field and crosspollinate to produce the variety of seed.

In the mid-60's, the New Jersey Agricultural Experiment Station (Rutgers) turfgrass breeding program was led by C. Reed Funk. Jerry Pepin was Funk's student at the time. These two men at Rutgers and Joe Duich at Penn State started a revolution with improved ryegrasses.

Pennfine was released by Duich in 1968, after Manhattan had been released from Rutgers. It was a three-clone variety, with two parents originally from Pennsylvania golf courses and one from a grass tennis court. Duich made the decision to hold off marketing Pennfine until the expected Plant Variety Protection Act was passed (1970). Today, Pennfine is succesfully marketed by SPIC.

Manhattan, however, was released prior to the Plant Variety Protection Act and did experience problems early in its marketing. It was first marketed in 1968 by Bill Rose of Turf Seed who had taken six pounds of breeder seed and gotten production started. Today, Manhattan is marketed by Turf Seed and Whitney Dickinson as agents to the Manhattan Ryegrass Growers Association. Rutgers has taken part in some way with an amazing number of perennial ryegrasses. They include: Blazer, Dasher, and Fiesta from Pickseed West; Belle from Burlingham; Derby from International Seeds; Goalie, Delray and NK-100 from Northrup King; Omega from Turf Seed; Pennant from Agricultural Services; Diplomat, Yorktown and Yorktown II from Lofts; and Regal from North American Plant Breeders.

Turf Seed developed Birdie and Citation. Northrup King has developed Eton, Epic, and NK-200. International Seeds has developed Clipper and Scotts has developed Caravelle.

European material includes Loretta from Scotts and Hunter and Elka from International Seeds.

One use of perennial ryegrasses that is receiving a great deal of attention is as a transition grass for the south. It is overseeded in large quantities to keep greens colorful and soft in the winter. Turf Seed has developed a annual/perennial ryegrass for overseeding, called Oragreen.



Jerry Pepin

A student of Reed Funk's in the 60's, Pepin has carried perennial ryegrass improvement from Rutgers to Rudy Patrick and now to International Seeds Inc. of Halsey, Oregon. He is the breeder of Derby, Regal and a number of other improved turfgrasses.

Crownvetch

Crownvetch, Coronilla varia L., is not a grass or monocotyledon. It is a perennial, dicotyledon herbaceous plant with pinkish blossoms that serves to cover and stabilize roadsides and slopes due to its spreading ability and deep root system. Stanford and Turf Seed market Penngift Crownvetch, which was discovered, produced, and promoted by Fred Grau of Penn State. Grasslyn Farms, managed by Fred Grau Jr., produces much of the seed for Stanford to market.

An odd situation with Penngift was when Grasslyn was the only producer of the seed, the Highway Department would not buy from it because it was a monopoly. This led to the development of Chemung and Emerald Crownvetch by the Soil Conservation Service in the early 60's.

Grau discovered the legume on a Pennsylvania farm in 1935. In 1947, he had produced seed on his farm and gave demonstrations of the cover across the state. Burt Musser suggested Grau scarify the seed to improve germination in 1951. Due to the problem with the Highway Department, Grau had to assist in setting up his competition.

WARM SEASON GRASSES

Bahiagrass

Bahiagrass, Paspalum notatum Flugge, was brought to the U.S. from Brazil as a low maintenance turf for semitropical areas. Argentine and Pensacola are varieties developed by the Florida Agricultural Experiment Station in the late 40's.

Hugh Whiting, a private turf breeder in California has developed Adalayd, *Paspalum vaginatum*, to improve the species.

Bermudagrass

Bermudagrass Cynodon dactylon, is the most important warm-season turfgrass in the U.S. It is propagated mainly vegetatively.

Many states have been involved in improving bermudagrass, including Florida, Kansas, Texas, California, South Carolina, Oklahoma, Arkansas, Alabama, Georgia and Maryland. However, their work is overshadowed by the developments of Glenn Burton with the USDA in Tifton, Georgia.

Burton began his work on the "Tif" series in 1946 after being encouraged by Fred Grau and Olaf Aamodt from USDA Beltsville. He collected dwarf pasture bermudagrasses and



Director of Pure Seed Testing in Hubbard, Oregon, and leading turfgrass specialist in the area of stem rust. Meyer is finishing development work on an improved Merion and an improved Manhattan.

crossed them with selections from golf greens. Golfers were complaining that bermuda greens were too coarse. From these hybrids, Burton selected one released as Tiflawn in 1952. But Tiflawn was still too coarse for greens. A finer turf was required.


Burton got hold of a bermudagrass from Africa, Cynodon transvaalensis, a softer, finer variety. He bred the African bermuda with a dense selection of Cynodon dactylon. The result was a sterile, but improved variety which he called Tiffine. It was released in 1953. But the bermuda was sterile and improvement stopped at that point for that turfgrass.

Burton went back to his collection for another *C. dactylon* to breed with *C. transvaalensis.* He selected a bermuda from Charlott Country Club in North Carolina. The cross produced another sterile bermuda which he called Tifgreen. It was released in 1956 and made a much improved bermuda for greens.

Looking for better frost tolerance, Burton made a third cross with *C*. *transvaalensis*. He got what he wanted but it was stiffer than Tifgreen. This bermuda was released in 1960 as Tifway.

Fortunately, Tifgreen produced a vegetative mutant with finer stems, smaller and darker leaves. Burton

Underneath it all... Weather-matic; the hidden secret of beautiful landscaping.



Quiet, Cost-effective operation for medium range with our G Series sprinklers

- Rotary pop-up with sealed gear train.
 Minimal pressure and flow required to
- pop-up and seal.
- Self flushing action removes particles.
 Full and part circle sprinklers can be
- valved together.
- · Easy to service above ground.
- Arc size field-adjustable from 70° to 340°.
- Arc location adjustable during
- sprinkler operation.
- Housing, gears and components of highest quality, space-age plastics.
- Optional features: rubber covers, vandal resistant screws, range adjustment on G40F and G40PC.

For additional information about our G Series rotary sprinklers, send the coupon below.

	MAIL TO: Weather-matic Division TELSCO Industries P.O. Box 18205 Dallas, Texas 75218 Send information on:
Distributor D	G Series Sprinklers Catalog
	G Series Sprinklers Catalo Directory TURF IRRIGATION Manua
NAME	G Series Sprinklers Catalog Directory TURF IRRIGATION Manual
NAME	G Series Sprinklers Catalo Directory TURF IRRIGATION Manua
NAME COMPANY ADDRESS	G Series Sprinklers Catalo Directory TURF IRRIGATION Manua

ODUCTION




10. Field burning to stimulate the plants for next year's production, remove straw residue from the fields, and disease and insect control.



11. Experimental field burning machines were tried to control the volume of burning but they were dropped as too expensive and unreliable.

Photos courtesy of Harry Stalford, International Seeds Inc.



Book Department 9800 Detroit Ave. Cleveland, Ohio 44102.

released Tifdwarf in 1965.

Since 1965, Burton has experimented with ways to cause mutations of the sterile bermudas he has produced. In 1970, he first used radiation to produce mutants and was successful.

This space-age technology has produced more than 150 mutants for evaluation. Burton says he expects to release one or more of the mutants within the next year.

In more conventional methods of selection, good varieties of C. dactylon have been developed. U-3 was one of the first selected from material supplied to USDA in Arlington and later Beltsville by Lester Hall. Hall found the bermuda on his golf course in Savannah, Georgia. It was released as U-3 in 1947. Ormond from Florida, Royal Cape from California, Texturf from Texas, and Tufcote from Maryland are selections of C. dactylon.

Other crosses of C. dactylon and C. transvaalensis have been accomplished by Florida (Everglades), Kansas (Midway), and South Carolina (Pee Dee).

Uganda is a natural selection of C. Transvaalensis.

Centipedegrass

Centipedegrass Eremochloa

ophiuroides, originated from China. It exhibits poor wear tolerance, but provides an adequate turf in warm regions without great care. It exhibits extremely tough resistance to insects and disease which may cause a closer evaluation in the future. It may serve for lower traffic areas on fairways and roughs.

Kikuyugrass

Kikuyugrass, Pennisetum clandestinum is another turfgrass brought from Africa for use in the U.S. It is a tough turfgrass which tolerates high temperatures, low cutting, wear, and some shade. Extended cold weather will damage it however.

St. Augustine

St. Augustine Stenotaphrum secundatum, is second to bermudagrass for warm season turfgrass use. It is an aggressive, lowgrowing, heat tolerant, blue-green turfgrass. Like Kikuvugrass, it will not tolerate extended cold temperature. It forms a good sod and can tolerate some shade. Overfertilization can create severe thatch buildup. Floratine is a variety developed by Florida Agricultural Experiment Station specialists.

