# Landscape Managers' Guide to: Tall Fescues

by William A. Meyer, Turf Seed, Inc., Hubbard, OR

Tall fescue (Festuca arundinacea Schreb.) is becoming increasingly popular as a turfgrass in many areas of the United States. This species has been a very important grass for forage, roadside stabilization and erosion control for many years.

The increased demand for tall fescue for turf has resulted from escalating energy costs to produce fertilizers and pesticides, and water shortages in many areas, especially in the southwestern U.S. Compared to other coolseason grass species, tall fescue is unique due to its deep root system for drought tolerance or avoidance, the ability to persist at reduced soil fertility, and reasonably good tolerance to many pests.

The release of new lower-growing tall fescues starting in the late 1970's has increased the interest in tall fescues for turf.

### Origin and history

Tall fescue is a native of Europe and is found throughout Europe, in Asia, eastern and southern Africa, Madagascar, southern Australia, New Zealand, and in large areas throughout the United States. It tolerates a variety of geographic and climatic conditions, which explains its wide range of adaptation.

This species, now found in most of the 50 states, was introduced in the early 1800's. Because of its morphological resemblance to meadow fescue, it was not classified as a distinct species until 1950, even though it was already being widely used for forage, conservation and roadsides.

The utilization of tall fescue increased a great deal with the development of the first improved cultivars from naturalized American tall fescues. Harry Schoth of Oregon State University developed and released Alta in 1941. Kentucky 31 (KY-31) was developed and released by the University of Kentucky in 1943. Both KY-31 and Alta were developed primarily as forage grasses, but have been used as turfgrasses.

By the 1960's, tall fescue was being recommended for turf by the state of

Tennessee because of its wide adaptation to variable soil pH, rainfall and sunlight.

Until recently, KY-31 has been the most widely used and recommended cultivar for turf, especially in the transition zone of the United States, where other cool-season grasses are not well-adapted and the warm-season grasses are sometimes killed by the cold. This transition zone includes an area south of central Illinois, east of Topeka, KS, north of Macon, GA, and west of the Atlantic coastal plain of the Carolinas. Both Alta and K-31 have been used in the southwestern U.S., especially in California and at higher elevations in Arizona and Nevada.

KY-31 has produced a better turf quality than Alta in most turf trials. In the transition zone the common practice for turf managers is to reseed KY-31 lawns each year to maintain denser stands.

### Growth characteristics

The tall fescues live up to their name when they are allowed to produce seedheads, growing as tall as 4-5 feet.

Tall fescue is basically a bunchgrass, but limited rhizomes can be seen on certain plants, especially when they are grown in light, sandy soils. Breeding work is in progress to develop varieties with increased rhizome production. Thick stands of the present varieties can produce a tough sod if sufficient time is given for a mature turf.

In many northern lawns of Kentucky bluegrass, tall fescue can be a severe weed problem when single, coarse clumps are scattered in a turf area. The leaf blades of KY-31 and other common-type cultivars are generally a light-green color and coarse, with a harsh feel to the hand. Individual plants can be found in old turf stands that are finer-bladed with a softer texture and a reduced vertical growth rate when compared to KY-31.

Most breeding programs to develop improved turf-type varieties have been based on selections from old turf areas that have the ability to develop a denser, finer-bladed turf with improved persistence. The tall fescues usually slow up their growth and go dormant during cold periods earlier than the improved turf-type perennial ryegrasses and later than most Kentucky bluegrasses. Their spring green-up rate is also intermediate to these other two species.

### Breeding turf-types

The main disadvantage of KY-31 and other older tall fescue varieties is they produce a coarse open turf with a light green color. Dr. C. Reed Funk initiated a breeding project at the New Jersey Agricultural Experiment Station to develop tall fescue cultivars for turf purposes. Other breeding programs were started in the 1970's with some programs as cooperative breeding projects with Funk.

Rebel was the first turf-type released from the above programs, followed by Falcon and Olympic. The varieties have leaves approximately 30-40 percent finer than KY-31 and densities of tillers almost twice KY-31. They all have a darker blue-green color than KY-31 with improved persistence in turf. They also have improved disease resistance compared to KY-31.

In most turf trials at both high and low maintenance these new varieties have shown superior turf performance. Other turf-type varieties that have shown improved turf performance over KY-31 are Bonanza, Adventure, Jaguar, Apache, Arid, Mustang, Houndog and Finelawn I. It is interesting to note that some of the later maturing varieties (in seed production) such as Jaguar, Adventure, Bonanza and Apache have generally given superior turf performance than earlier maturing varieties. The varieties Bonanza and Apache possess the darkest genetic blue-green color of the present varieties, followed by Mustang and Olympic.

### **Disease resistance**

The most common and serious disease of tall fescues are Rhizoctonia brown patch and net blotch (Helminthosporium leaf spot). Other diseases occur, but not as widespread as these two.

Net Blotch Under cool cloudy and wet conditions net blotch caused by H. dictvoides Drechs. can be found on tall fescues whereever they grow. In western Oregon this disease is especially severe on seedlings when mowing and traffic occurs during the establishment period. The symptoms are net like patterns of streaks of dark brown tissue on the leaves. When the disease is severe the leaves turn vellowish-brown and die back from the tips. The varieties Bonanza, Mustang, Jaguar and Olympic have shown the best resistance to this disease to date. Breeding programs in New Jersey and Oregon have shown that field selection in space plants can be used effectively to develop more resistant varieties.

Brown Patch Brown patch caused by Rhizoctonia solani Kuhn is a very serious disease of tall fescue. This disease is most severe under hot and humid conditions when the turf is subjected to long wet periods. High fertility, close mowing and poor air circulation increases the severity of this disease. Breeding programs to develop cultivars with increased brown patch resistance have been promising to date. Among the new turf-type varieties, Adventure, Jaguar and Bonanza have shown the best resistance to date. Many of the cultivars from Europe and the varieties Alta, Goar and Fawn have been damaged more severely than KY-31 or the new improved turf-type cultivars.

Brown patch can also become more severe when drought stress precedes humid conditions favoring disease development. Varieties that have improved resistance to drought stress generally show less damage from this disease.

Some varieties are able to recover from brown patch injury better than others. Most varieties recover very rapidly as soon as cool, favorable growing conditions occur. Breeding programs are in progress to develop higher levels of brown patch resistance in new varieties of tall fescue.

Adding five percent by weight of Kentucky bluegrass to a tall fescue turf has shown promise to reduce the development of brown patch. Care must be taken to select bluegrass varieties with only moderate aggressiveness to prevent crowding out of the tall fescue by bluegrass.

It is interesting to note brown patch is not as serious a problem in arid areas of the Southwest as in the transition zone.

Pythium Blight Various species of

pythium can destroy seedlings or mature turf of tall fescue when weather conditions are similar to those favoring brown patch. This disease, called Pythium blight, is usually most severe in poorly drained areas and can rapidly deteriorate turf quality if conditions for disease activity persist for more than a short period.

Koban or Apron fungicides are effective to control Pythium blight as seed treatments. A change in weather conditions to cooler temperatures and reduced humidity will usually halt the development of this disease.

The level of resistance in the present cultivars is questionable considering the number of pythium species involved. Adventure and Falcon had somewhat less disease in a New Jersey trial in 1983.

**Crown Rust** During cool spring and fall periods crown rust caused by Puccinia coronata cda. can occur on tall fescues in many areas of the U.S. The y ellow orange pustules are urediospores of this fungus and in the late fall the crown-like teliospores (hence the name) can occur. Tall fescue is generally more resistant than meadow fescues. Progress has been made in selecting varieties with improved resistance in Western Oregon. Alta and Fawn are quite susceptible while KY-31 and most of the new turftypes are quite resistant.

**Powdery Mildew** Some of the new turf-type varieties which develop very dense leaf canopies in seed production fields have been found to be susceptible to powdery mildew. This disease is usually restricted to the dense center parts of each plant that is severely shaded. It appears that most of the present varieties can be affected when excessive foliage is produced in the springtime. From space plant observations it appears that breeding for improved resistance to this disease is possible. Powdery mildew has not been noted in shade trials to date.

### Heat, drought tolerance

Tall fescue is a very deep rooted species which helps to tap water resources deeper in the soil profile than the other cool-season species. In Nebraska studies by Kopec and others, in two-inch mowed turf they found roots of the new turf-type varieties 48-inches deep and roots of KY-31 54-inches deep. It is encouraging to see that the roots of these denser and fine textured varieties are maintaining rooting depth comparable to the more open and coarser KY-31.

In 1981 studies in western Oregon, the varieties Olympic and Falcon were still maintaining 80 percent or more green color after 70 days of drought compared to 67 percent for hard fescue, 32 percent for chewings fescue, 44 percent for creeping fescue, 40 percent for perennial ryegrass and 16 percent for Kentucky bluegrass.

In water use studies the tall fescues have been found to have a higher water loss rate that most other warm season grasses and some of the cool season species. This high water loss rate has a beneficial cooling effect during heat stress periods as long as the roots can reach available water reserves.

Most of the present varieties of turf-type tall fescues have shown similar response to drought and heat stress to KY-31. Attempts to develop an extremely dwarf and fine variety (similar in texture to turf-type perennial ryegrass or Kentucky bluegrass) have resulted in problems with reduced drought and heat stress and increased disease problems. These very fine and dwarf clones are being crossed with well-adapted varieties with higher levels of summer disease resistance.

### Extreme soil conditions

Tall fescues are able to grow on a wider range of soil conditions than other cool season species. The new turf-type varieties are able to maintain much better density at low fertility than KY-31 at low fertility. The darker colored varieties are more striking at low fertility levels.

Tall fescues have above average tolerance for acidic, alkaline, saline and poorly drained soils. In California trials the varieties Olympic and Jaguar showed improved resistance to iron chlorosis than other varieties.

### Shade tolerance

Many of the new turf-type varieties, especially those with improved net blotch resistance, have shown improved shade tolerance over KY-31. All of the varieties become finer and softer in shade, which makes tall fescue more compatible in mixtures with other species.

In shade situations, the tall fescues can be found very far south because of the cooling effects from the shade.

### Cold tolerance

In trials on the high desert of eastern Oregon the tall fescues have shown very good cold tolerance in space plantings when compared to all of the perennial ryegrasses over four winters. In New Jersey, Pennsylvania and Rhode Island the well-adapted varieties have suffered winter damage under severe ice sheet conditions. The cold hardiness of tall fescues is reduced when they are heavily fertil-

### **Tall Fescue Varieties**

A brief description of the tall fescues presently available or becoming available in the near future.

Adventure tall fescue was developed by Pure-Seed Testing, Inc. of Hubbard, Oregon. It will be distributed by Warren's Turf Nursery. Adventure is an advanced generation synthetic variety derived from selections from old turf areas in the northeastern U.S. It has performed very well in turf trials throughout the U.S. both in full sun and partial shade. Adventure is an attractive, vigorous, leafy, turf-type variety with a medium dark green color, medium-fine leaf texture and good density. It has shown greater vigor and better color at low fertility levels than other tall fescues currently available. Adventure has good resistance to Rhizoctonia brown patch and moderate resistance to net blotch disease. Seed became available in 1983.

Alta tall fescue was developed by the Oregon Agricultural Experiment Station with the cooperation of the Agricultural Research Service of the United States Department of Agriculture. It was released in 1940. Plants were selected out of two introductions from Germany and one commercial seed lot on the basis of improved winter survival, ability to remain green during dry summers in western Oregon, high forage yields, and ability to persist. Alta has heen used extensively for turf in California. It produces a rather coarse, upright turf of moderately low density and rapid leaf elongation. Alta has shown above average susceptibility to net blotch and Rhizoctonia brown patch in turf trials and crown rust in western Oregon. Alta is a commercially available variety.

Apache tall fescue is a new, moderately low growing turf-type variety with a dark blue-green color when compared to most other varieties. This variety was developed by Pure-Seed Testing, Inc. and will be marketed by Turf-Seed, Inc. It is an advanced generation synthetic variety derived from selections from old turf areas in New Jersey. Georgia and Illinois. Apache has shown moderately good resistance to net blotch, brown patch and very good resistance to crown rust. It has shown good density and a medium-fine texture with a reduced rate of leaf elongation. It has shown good traffic, heat and drought tolerance in trials to date and good color retention at low fertility. Commercial seed will become available following the 1984 harvest.

Astro tall fescue is a new, moderately low growing variety developed by Pure-Seed Testing and marketed by Green Seed Co. It is an advanced generation synthetic variety developed from parental clones selected from old turf in North Carolina, Illinois and New Jersey. This variety has a medium dark green color with medium texture and density. Astro has shown moderately good resistance to net blotch, crown rust and brown patch and good heat, cold and drought tolerance and good persistence in turf. Commercial quantities of seed will be available after the 1984 harvest.

Bonanza tall fescue is a new, low growing turf-type developed by Pure-Seed Testing, Inc. and marketed by Cenex, Inc. This variety is a 3-clone synthetic derived from germplasm from New Jersey and Georgia. This variety has a very dark blue-green color when compared to other tall fescues and has a dwarfer, slow leaf elongation rate compared to other varieties. Bonanza has very good resistance to brown patch, net blotch and crown rust. The wear, drought, heat and cold tolerance of this dense variety has been very good in trials to date. It has shown good color retention at low fertility and will be available in very limited quantities after the 1984 harvest

Brookston was developed by North American Plant Breeders and International Seeds, Inc. It is produced and marketed by Stanford Seed of Buffalo, New York. Brookston is a leafy, moderately low growing turf-type tall fescue with a medium dark green color, medium texture and medium density. It is reported to have improved low temperature hardiness. Brookston has very good resistance to net blotch and crown rust. It has shown reduced summer stress tolerance. Limited amounts of seed became available following the 1983 harvest.

Clemfine was developed and released by the South Carolina Agricultural Experiment Station and Loft's Seed, Inc. It originated from germplasm selected from old turfs located in the southeastern United States. Clemfine has a medium green color, coarse texture, and a medium-low density. It has moderatley good resistance to Rhizoctonia brown patch, but only fair resistance to net blotch. Clemfine closely resembles Kentucky 31 tall fescue in many characteristics, but has generally shown improved performance and increased persistence in turf trials. Seed of Clemfine became available after the 1982 harvest.

Falcon tall fescue was developed cooperatively by Pure-Seed Testing, Inc. and E. F. Burlingham & Sons from germplasm obtained from the New Jersey Agricultural Experiment Station. The first certified seed was produced in 1980. Falcon is a leafy, moderately low growing, turf-type variety. It has the ability to produce an attractive, more persistent turf with finer texture and higher density than most of the currently available varieties of tall fescue. It has improved resistance to brown patch, crown rust and net blotch.

Fawn was developed at the Oregon Agricultural Experiment Station and released in 1964. It produces high seed and forage vields in the Willamette Valley where it shows greater height and vigor in early spring and earlier maturity than Alta or Kentucky 31. Fawn produces a coarse, open turf with a rapid rate of leaf elongation. It is susceptible to net blotch and Rhizoctonia brown patch diseases in New Jersey turf trials and to crown rust in western Oregon. Fawn also appears to be less winter hardy and less tolerant of adverse summer conditions than Kentucky 31.

Finelawn I is a new moderately low growing turf-type variety developed by Pure-Seed Testing, Inc. and marketed by Finelawn Lawn Research, Inc. of Madison, Georgia. This is an advanced generation synthetic variety developed from clones derived from old turf areas in Alabama, Georgia, Pennsylvania and New Jersey. This variety has shown moderately good resistance to net blotch, brown patch and crown rust. It has shown good drought, heat and cold tolerance. It became commercially available following the 1983 harvest.

Galway was developed by Northrup-King. It is a medium dark green turftype variety with medium density and a medium coarse texture. The variety has good heat and drought tolerance and improved cold hardiness. It does well in light to moderate shade as well as in full sun. Galway has shown moderate resistance to Rhizoctonia brown patch and net blotch. Galway is a moderately low growing variety intermediate in growth habit between the turftype varieties and Kentucky-31. Commercial seed production has been initiated.

Goar tall fescue was selected at the Imperial Valley Field Station at El Centro, California and developed in cooperation with the Plant Materials Center, Soil Conservation Service, Pleasanton, California and the California Agriculture Experiment Station. The parental germplasm was obtained from Hungary. It was released in 1946. Goar *continued on page 54*  is an early maturing, vigorous, rather coarse bunchgrass adapted to heavy textured alkaline soils. It is reported to show improved growth during high summer temperatures in California. Goar has performed poorly in turf trials with high susceptibility to Helminthosporium blight and Rhizoctonia brown patch. Goar is reported to be susceptible to crown rust.

Houndog was developed and released by International Seeds, Inc. of Halsey, Oregon. Its parental germplasm originated from plants selected from old turfs in Kentucky and Tennessee plus plants selected from Rutgers and Missouri. Houndog is a leafy, persistent, moderately low growing, turf-type variety with a medium dark green color, medium texture, medium density, and a semi-prostrate growth habit. It has good heat and drought tolerance, performs well in shade and shows good color retention in late fall. Houndog has moderate resistance to Rhizoctonia brown patch and net blotch and crown rust. Seed of Houndog became available following the 1983 harvest.

Jaguar was developed by Pure-Seed Testing, Inc. of Hubbard, Oregon. It will be distributed by Garfield Williamson, Inc. of Jersey City, New Jersey. Jaguar is an attractive, leafy turf-type tall fescue with medium density and texture, a medium dark green color and a moderately low growth habit. It has good heat and drought tolerance, good shade adaptation, and very good color retention with low temperatures in late fall. Jaguar has very good resistance to crown rust, Rhizoctonia brown patch and net blotch. It has done very well in trials Seed became available commercially following the 1983 harvest.

Kenhy tall fescue was developed cooperatively by the Kentucky Agricultural Experiment Station and the Agricultural Research Service of the United States Department of Agriculture. It was released in 1976. Kenhy is a synthetic of eleven 42 chromosome derivatives of annual ryegrass X tall fescue hybrids which were selected for vigor, soft lax leaves, and forage with high moisture content during drought stress. Kenhy has a lower stomatal frequency than Kentucky-31, which may contribute to its lower percentage of rolled leaves, greater palatibility, and greener foliage during summer drought stress. Kenhy produces a rather coarse, moderately open turf with a rapid rate of leaf elongation. It has shown improved

ized and mowed closely.

### **Traffic tolerance**

In our wear trials in western Oregon and the Sports Turf Research Institute in Bingley, England, the new improved varieties such as Falcon, Adventure, Mustang, Olympic and Rebel have been found to have superior traffic tolerance than KY-31 and other common type varieties. The crowns of tall fescue are slightly deeper in the soil profile than perennial ryegrasses. This may be the reason why these grasses have shown improved traffic tolerance.

### Insect resistance

Tall fescues generally have better resistance or tolerance to insects than the other cool-season turf species.

In tests in western Oregon billbugs severely damaged many of the Kentucky bluegrasses, perennial ryegrasses and fine fescues. The adjacent plots of new and old varieties of tall fescue showed no damage.

There is a possibility that the endophyte, commonly referred to as *Epichloe typhina*, which has been found to convey insect resistance when it is present in perennial ryegrasses, is also involved in conveying insect re-



tolerance of heat and drought and improved overall turf performance in Kentucky trials compared to Kentucky-31.

Kentucky-31 tall fescue originated as an ecotype selection discovered growing on the William Suiter farm in Menifee County, Kentucky, where it had apparently been growing since 1887 or earlier. The variety was collected, tested and developed by the Kentucky Agricultural Experiment Station and released in 1943. Kentucky-31 is very widely used for both turf and forage purposes in the lower midwest and upper south. It is adapted to a wide range of soil types and shows a better tolerance of climatic extremes than most other tall fescue varieties. Kentucky-31 produces rather coarse, moderately open turf with a rapid rate of leaf elongation. It has moderate tolerance of net blotch, Rhizoctonia brown patch and crown rust and it shows less damage from these disease than Alta, Fawn or Goar.

Marathon is a new, moderately low growing turf-type tall fescue with medium texture, density and a dark green color. This variety was developed by International Seeds, Inc. It was derived from elite parental clones selected in the transition zone of the U.S. Marathon has shown moderately good resistance to Rhizoctonia brown patch and net blotch and good resistance to crown rust. It has exhibited good heat and drought tolerance and is being marketed exclusively by Hubbard Seed Supply, Hubbard, Oregon.

Maverick was developed by Pickseed West Inc. of Tangent, OR, using germplasm from the USDA, Beltsville, MD. It is a moderately low growing, medium-fine textured variety with a pleasing medium dark green color. Maverick has better heat tolerance than most turf tall fescues and has very good shade tolerance. It is adapted to the upper south or transition zone and to low to medium turf management.

Mustang was developed by Pickseed West, Inc. of Tangent, Oregon, using germplasm obtained from the New Jersey Agricultural Experiment Station. It is an attractive, leafy, moderately low growing, turf-type variety with medium texture and density. The variety has very good shade adaptation and good fall low temperature color retention and spring greenup. It has shown very good resistance to net sistance to certain tall fescue varieties. This fungus normally produces no external symptoms and lives only within the tissues of plants by growing between the cells. It is transmitted from one generation to the next by seed. After seed has been stored for two years or more usually the percentage of seeds with viable endophyte goes down to very low levels.

Many of the new turf-type varieties of tall fescue have been found to have low percentages of infection in their seed lots, while KY-31 has been found to have levels over 90 percent.

### Establishment

It is very important that quality seed be used to produce quality turf. Seed that has been certified with a seal and tag on the bag should be used when ever possible. Usually certified seed costs only 5-10 percent more than uncertified, and the consumer can be confident he is getting seed of a variety that the breeder has described. This will help to avoid varietal substitution, poor germination and low purities.

It is also important that there are no objectionable crop (such as orchardgrass, bentgrass, poa trivialis and ryegrass) or weed (annual bluecontinued on page 58





## **TINY TERRA 140**

### The small aerator built to handle the big jobs.

140 tines in 22" of aerating width overall width of 35" to get in the tight places.

This Tiny Terra produces more holes than most aerators do with a 6' width. It can be pulled easily by a 10 horse garden tractor.

> **TERRACARE** *PRODUCTS CO., INC.* P.O. Box 506 • Pardeeville, WI 53954 Phone—608-429-3402 Circle No. 134 on Reader Inquiry Card

Remember the name because you're going to hear a lot about it. Fylking Kentucky bluegrass lawn is something special. It forms a dense, thick turf, more disease and drought resistant. Greens up earlier, stays greener longer. Fylking can be cut as low as ¾ inch, even ½ inch for home putting greens. Proven over many years of international certified testing. Ask your seed distributor for Fylking.

Iking - the World's Fair grass.



Another fine, quality-controlled roduct of Jacklin Seed Company.

### TALL FESCUE from page 52

blotch, and moderately good resistance to Rhizoctonia brown patch. Mustang has good tolerance to close mowing with minimal thatching. Seed is commercially available.

Olympic was developed by Pure-Seed Testing, Inc. of Hubbard, Oregon using germplasm developed at the New Jersey Agricultural Experiment Station. It was released by Turf-Seed. Inc. Most of its parental germplasm was collected from old turfs in Alabama, North Carolina and New Jersey. The first certified seed was produced in 1981. Olympic is an advanced generation synthetic variety derived from the progenies of eight clones. These parental clones were selected from spaced-plant nurseries because of attractive appearance. freedom from disease, softness of leaves, ability to resist leaf roll during hot dry weather, dark green color, and promising seed yield potential. Single plant progenies of selected clones were evaulated in closely mowed turf trials in New Jersey and Oregon. Seedlings from clones showing the best progeny performance were screened for resistance to crown rust, uniform maturity, attractive appearance, and acceptable seed yield. Breeder seed was harvested from the remaining 1004 plants. Olympic is an attractive, leafy, persistent turf-type variety with a dark green color, medium texture and density, and a moderately low growth profile. It has good heat tolerance, performs well in moderate shade and has very good color rentention in late fall. It has good resistance to net blotch and crown rust and moderately good resistance to Rhizoctonia brown patch. It retains an acceptable green color at low nitrogen fertility levels. Olympic has shown less iron chlorosis than other tall fescues on alkaline soils in California. Commercial seed is available.

Rebel tall fescue was developed by Lofts Seed, Inc. from germplasm obtain from the New Jersey Agricultural Experiment Station. Plants collected from old turfs of New Jersey and surrounding states contributed most of the germplasm of Rebel. Additionally, part of the parental germplasm was obtained from a number of accessions received from the United States Plant Introduction program and from trispecies hybrids of tall fescue, meadow fescue, and perennial ryegrass obtained from the United States Regional Pasture Research Laboratory; University Park, Pennsylvania. Clones of the original accessions were initially evaluated in nurseries subjected to frequent close mowing. Single plant progenies of the most promising selections were subsequently subjected to three cycles of phenotypic recurrent selection for persistence, attractiveness, disease resistance, and performance in turf trials maintained at three-quarter-inch cutting height. Rebel is a turf-type variety



**ELEMENT IN ANY COMPLETE** TURF CARE PROGRAM. Let's face it - low price is no substitute for poor product. Challenged by tough turf problems together with rising costs, you just can't afford to play games with products that don't perform consistently

18-5-8 FERTILIZER OR CHEMICAL SUMPL

and economically. You need quality turf products you can depend on.

You need COUNTRY CLUB and GREENSKEEPER PROFESSIONAL TURE PRODUCTS from LEBANON

COUNTRY CLUB AND GREENSKEEPER FERTILIZERS ARE OF THE HIGHEST QUALITY HOMOGENOUS GRADES MADE TODAY - GIVING YOU MORE FEEDING POWER FOR YOUR DOLLARS. What makes homogenous fertilizers better? Each pellet contains a chemicallybalanced combination of N-P-K, regardless of the material's sizing and coverage. The plant root needs only to come in contact with one pellet to receive a balanced diet. Consequently, you get even element distribution.

WHATEVER YOUR TURF NEEDS, THERE'S A COUNTRY CLUB/ **GREENSKEEPER PRODUCT SPECIALLY FORMULATED TO MEET** THEM - ALL WITHIN YOUR BUDGET! Backed with over 35 years of extensive university and end-use testing, we offer you the most complete line of fertilizers and control products available on the market as well as a full range of S.C.U. blends and chemicals.

LEBANON CHEMICAL has the flexibility to formulate exactly the analysis your turf tests indicate to be most beneficial. We are fully equipped to manufacture your private label requirements when needed.

EVERY COUNTRY CLUB AND GREENSKEEPER PRODUCT IS BACKED WITH A TOTAL COMMITMENT TO SERVE YOU BETTER. Our technically experienced staff is eager to assist you in solving turfgrass problems and will design a complete turf care program for you! Plus you're supported by LEBANON'S solid network of distributors throughout the country that ensure you of product availability and prompt service.



that has an attractive, leafy, persistent turf of greater density, finer texture, and a slower rate of vertical growth than most of the commercially available varieties of tall fescue. Rebel has improved resistance to brown patch and moderate resistance to net blotch. Rebel became commercially available following the 1979 harvest.

Tempo is a new moderately growing turf-type tall fescue developed by Agriculture Services Corp. from selections from old turf areas in and around Ocean Shores, Maryland, It has a medium green color with medium texture and density. Tempo has moderate resistance to net blotch with improved drought tolerance. Selection was made for plants with narrower leaves and a slower rate of vertical leaf growth. Commercial seed will be produced and marketed by Normarc, Inc., Tangent, Oregon and will be available Fall of 1984.

Willamette is a new moderately low growing turf-type tall fescue with an attractive moderately dark green color. It has a medium texture and density and was developed by International Seeds. Inc. Willamette was derived from elite clones selected from old turf areas in the midwest and mid-southern U.S. It has shown improved resistance to brown patch and crown rust and moderate resistance to net blotch. This variety has also shown improved heat, cold and drought tolerance and will be marketed by Willamette Seed and Grain Co., Albany, Oregon. 🗆

grass or velvetgrass) seeds in a lot.

Most of the KY-31 is grown as a biproduct of pastures in Missouri. Much of the seed from this production contains orchardgrass or other serious contaminants of tall fescue turf.

Much of the seed produced in western Oregon is produced by professional seed growers and the quality is generally better than the southern production. The two contaminants that can occur in Oregon production are annual and perennial ryegrass. These ryegrasses can cause severe competition during establishment and reduce turf quality. Some of the perennial ryegrasses may persist and cause a clumpy appearance.

Ground preparation, soil fertility, lime, seed placement and irrigation requirements at planting time are similar to other cool-season turf species. The seeds should be placed at a depth of 1/8-1/4 inch. Rolling the ground after seeding aids in making good seed contact and makes the surface smoother. The surface must be kept moist until germination occurs.

When optimum conditions exist,



When Japan's largest fertilizer supplier, Zen-Noh, needed phosphate rock to make fertilizer, they came to ESTECH. Through our joint venture at the Watson Mine in Ft. Meade, Florida, we helped Zen-Noh import over 600,000 tons of phosphate rock into Japan last year.

Which in turn grew a lot of grain. Which in turn fed a lot of people.

You'd think we would have shared this claim to fame before now. But that's the secret of our success...

> At ESTECH, we don't talk. We just do.



We'll share some equally revealing secrets about Professional & Commercial Fertilizers. Call Irv Stacy, National Sales Manager, Par Ex 813/533-1181 sprouts of tall fescue can be observed in 7 days, with turf requiring the first mowing at 2-3 weeks. Seed rates of KY-31 are usually recommended at 8-10 lbs. per 1000 sq. ft. The turf-type tall fescues have increased tillering ability compared to KY-31 and seeding rates of 4-8 lbs. per 1000 sq. ft. can produce a very good turf. Higher seeding rates of tall fescue can be used to give a more rapid turf cover to compete with weedy species. High seeding rates can result in additional irrigation requirements to maintain good turf during periods of drought and heat stress.

### **Cultural practices**

Tall fescues require less nitrogen fertilizer than the improved Kentucky bluegrasses and perennial ryegrasses. Sod growers in southern California have found that sod fields of tall fescue are managed best by cutting the fertilizer rates used for Kentucky bluegrass sod by one half. An annual fertility program of 2-4 lbs of actual nitrogen per 1000 sq. ft. is suggested. The optimum cutting height for tall fescues varies from one-and-one half to three inches depending on the amount of heat stress a turf will face. Cutting heights of one inch or lower can result in severe competition from annual bluegrass, especially under high irrigation and fertility conditions.

Tall fescues have shown good tolerance of the commonly used selective broadleaf and pre-emergence crabgrass herbicides.

The tall fescues are not thatch formers like the bluegrasses and fine fescues and usually don't require dethatching.

Under western Oregon summer drought conditions the tall fescues will stay green and grow 3-4 weeks longer than Kentucky bluegrass and 2 weeks longer than perennial ryegrass.

Tall fescue varieties should be used in blends with other varieties of tall fescue or in mixtures with five percent Kentucky bluegrass. The variety of bluegrass used should have only a moderate level of aggressiveness to prevent crowding out of the tall fescue. This is especially important when the turf is mowed at one-and-one-half inches or loss. Mixtures of tall fescue with perennial ryegrasses and fine fescues have resulted in crowding out of alot of the tall fescue by these species in western Oregon turf trials.

There are many breeding programs in progress at this time to make further improvements in the levels of disease resistance and turf quality of tall fescue. **WT&T**