

WEEDS TREES & TURF

The Magazine of Landscape and Golf Course Management Since 1962

Soil Amendments Are No Help to Tree Survival

Building Longer Life into Turf Equipment

A Professional's Guide to Kentucky Bluegrasses



**TURF DISEASE
CONTROL GUIDE**

Seed Series Part One

KENTUCKY BLUEGRASS

Hearty, aggressive Kentucky Blue is becoming more versatile than ever before.



Look beyond spring. Use DURSBAN to stop fall bugs, too!

Nothing tops DURSBAN* Insecticide for stopping spring-emerging insects in turf. But unless you spray it again in late summer, you could be heading for a fall, come autumn. Because even the long-lasting residual action of DURSBAN won't last quite that long. So you could miss sod webworms and other tardy types.

To protect your customers' lawns from spring through fall, apply DURSBAN Insecticide twice a year—on your first round to control the early risers, and again later in summer to stop the late arrivals.

The cost is reasonable: as little as 46¢ per 1000 sq. ft. application. That makes a double treatment a sound investment for your customers. They get year-long protection against insects. And you avoid the

awesome costs of call-backs and weakened customer relations.

DURSBAN will get just about every bug you want it to...chinch bugs, sod webworms, bill bugs, turfgrass weevils, armyworms, cutworms, ants and more. So look beyond spring. Apply DURSBAN Insecticide twice this year and get the fall bugs, too.

DURSBAN insecticide. In familiar 2E and concentrated 4E liquid formulations, and now 50W wettable powder. See your Dow distributor. And ask about our new "DURSBAN Delivers The Goods" incentive program. Be sure to read and follow all label directions and precautions. Agricultural Products Department, Midland, Michigan 48640.

DURSBAN

Keeps you looking good all year.



Dow Chemical U.S.A.

*Trademark of The Dow Chemical Company.

Circle No. 109 on Reader Inquiry Card

4901

WEEDS TREES & TURF

The Magazine of Landscape and Golf Course Management Since 1962



22 Disease Control Guide: Cool-Season Strategy

Dr. Joe Vargas, turf pathologist, Michigan State University, offers the latest tips on controlling major cool-season diseases. Chemical and cultural control are included in this reference you'll want to keep.

DEPARTMENTS

- 6** News/Trends
- 8** Green Industry News
- 12** Golf Update
- 14** Landscape Update
- 16** Government Update
- 20** Landscape Log
- 72** Problem Solvers
- 75** Events
- 76** Letters
- 82** Classifieds
- 84** Outlook

38 Disease Control Guide: Southern Turf Strategy

Dr. Don Blasingame, extension plant pathologist, Mississippi State University, zeroes in on the major diseases of warm-season turfgrasses and overseeded cool-season grasses.



44 Turf Fungicide Directory With Free Inquiry Card

Compile a reference file of labels and background on turf fungicides by circling the numbers of products listed in this directory on the free information card in the magazine.



46 Professional's Guide to Kentucky Bluegrasses

The difference between varieties of Kentucky bluegrass is significant to professional turf managers. Drs. Reed Funk and Ralph Engel of Rutgers describe strengths and weaknesses of common and proprietary Kentucky bluegrasses.

64 Engineers Build Self-preservation Into Turf Equipment

Manufacturers of professional turf equipment step beyond the basic engine:wheels:blade to provide longer life and greater reliability. New cleaning and cooling devices take equipment beyond the homeowner grade.

Robert L. Edgell, Chairman; Richard Moeller, President; Lars Fladmark, Executive Vice President; Arland Hirman, Treasurer; Thomas Greney, Senior Vice President; Ezra Pincus, Senior Vice President; Pat O'Rourke, Group Vice President; Joe Bilderbach, Vice President; James Gherna, Vice President; George Glenn, Vice President; Harry Ramaley, Vice President.

WEEDS TREES & TURF (ISSN 0043-1753) is published monthly by Harcourt Brace Jovanovich Publications. Corporate and Editorial offices: 7500 Old Oak Boulevard, Cleveland, Ohio 44130. Advertising Offices: 7500 Old Oak Boulevard, Cleveland, Ohio 44130, 111 East Wacker Drive, Chicago, Illinois 60601 and 3091 Maple Drive, Atlanta, Georgia 30305. Accounting, Advertising Production and Circulation offices: 1 East First Street, Duluth, Minnesota 55802. Subscription rates: \$20 per year in the United States; \$25 per year in Canada. All other countries: \$50 per year. Single copies (pre-paid only): \$2 in the U.S.; elsewhere \$4.50; add \$3.00 for shipping and handling per order. Second class postage paid at Duluth, Minnesota 55806 and additional mailing offices. Copyright © 1984 by Harcourt Brace Jovanovich, Inc. All rights reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical including photocopy, recording, or any information storage and retrieval system, without permission in writing from the publisher. Microfilm copies of articles are available through University Microfilm, International, 300 N. Zeeb Road, Ann Arbor, Michigan 48106.

POSTMASTER: Send address changes to WEEDS TREES & TURF, P.O. Box 6198, Duluth, Minnesota 55806-9898.

MC MC

MC



CMC3200

MASTER AND SATELLITE SYSTEM

...the natural evolution of the MC concept!

From the same people who developed the "state-of-the-art," computerized MC Series irrigation control systems now comes the ultimate Master and Satellite System, the CMC3200. In addition to unsurpassed ease and versatility of programming and 12, 18, and 24 station

satellites capable of independent operation, you'll find many unique features available for the first time in a competitively-priced product. Prominent among these is the first reliable cassette tape interface and printer option. For additional information, contact Irri-Trol.

The CMC 3200, see it soon.

IRRI-TROL® mfg., inc.

irrigation control systems

27940 Beale Court, Valencia, California 91355 • (805) 257-2333 • TELEX: 662600 IRTROLMFG VALE

CONTROLLERS • VALVES • VALVE ACTUATORS

Circle No. 253 on Reader Inquiry Card

This may be the only by trees, playgrounds and



At John Deere, we think we've found the perfect designers for our 318, 420 and 430 Lawn and Garden Tractors ... trees, playgrounds and hedgerows.

After all, look what they've put into our line.

Tough, economical 2-cylinder gasoline engines in the 318 and 420 (18 and 20 hp respectively). A 3-cylinder water-cooled diesel engine in the 430 (19 hp).

All three tractors have a 26-inch turning radius to get around easily in tight places. And power steering makes turning nearly effortless.

There's hydrostatic drive that lets you change speed and direction without clutching. And a welded steel frame takes humps and bumps in stride.

For better power delivery and greater durability, these John Deere tractors have differential axles with large cut-steel bevel gears. The 420 and 430 even have a 2-speed rear axle and differential lock for better traction in slippery conditions. And to help you finish big jobs without refueling, the 400 Series tractors have a large 6½-gallon capacity fuel tank.

There's even an optional Category "O" 3-point

line of tractors designed hedgerows.



hitch and a 2,000 rpm rear PTO. And a variety of attachments, including rotary tillers, snow blowers or throwers, front blade, thatchers, 3-point hitch mowers, and center-mounted mowers with 46-, 50- or 60-inch cutting width, depending on tractor.

Plus, you can count on John Deere dealer service. It's as reliable as the equipment he sells.

For the name of your nearest dealer, or a free folder on the John Deere lawn and garden tractor line, call 800-447-9126 toll free (1-800-322-6796 in Illinois) or write

John Deere, Dept. 50/50T, Moline, Illinois 61265.

John Deere lawn and garden tractors. Made to handle the work around trees, playgrounds and hedgerows... quickly and efficiently.



Nothing Runs Like a Deere®

Circle No. 108 on Reader Inquiry Card

by Bruce F. Shank, executive editor

Multi-course ownership grows as TPC eyes 20 courses by 1990

The pattern established by Club Corporation of America and American Golf Corporation is being carried a step further by the Tournament Players Club, which has set a goal to own 20 prestige, stadium-style golf courses within ten years.

TPC courses are designed to attract various tournaments and the revenue television and other media shell out. Pete Dye, primary architect for TPC, has a mean pen when he designs the courses.

Word is most of the 20 courses will be new, but a few, such as Edgewood Country Club, are reconstructed to Dye's specifications. See story in Golf Update.

Low dose pesticides may alter application methods in future

New low-dose herbicides and growth regulators, such as Du Pont's Oust and 3M's Embark are testing current application equipment and technique. These products are used in ounces per acre rather than pounds. Slight miscalculations can have harmful results. The value of the products and the sense of low dose pesticides in regard to environmental concerns, will force rethinking of application technology within the next five years. The resources of companies like Du Pont and 3M will see to that.

Housing starts to hit 1.7 million

The National Association of Home Builders are forecasting more than 1.7 million new homes for 1984. The increase may slow in the second half depending upon interest rate increases following the presidential elections. Apartment construction is expected to fall in some areas due to overbuilding.

Ford exec sees landscape boom

Jack Johnson, general sales manager for Ford Tractor North America, expects a complete recovery in construction by year's end. "The key to revival is housing starts," Johnson told HBJ Group Publisher Bob Earley during Ford's product introduction news conference in San Antonio, TX. The company has introduced new tractor/backhoe/loader models specifically for the landscape construction market.



Jack Johnson

STAFF

Executive Editor
Bruce F. Shank, Cleveland
Managing Editor
Maureen Hrehocik, Cleveland
Assistant Editor
Kevin Cooney, Cleveland
Publisher
Dick Gore, Atlanta
Senior Vice President
Tom Greney, Chicago
Group Publisher
Robert Earley, Cleveland
Production Manager
Kathy Judd, Duluth
Production Supervisor
Marilyn MacDonald, Duluth
Graphic Design
Beth Travis-Betts, Duluth
Circulation Manager
Kristine Bussell, Duluth
Directory Coordinator
Sheryl Albertson, Duluth
Reader Service Manager
Gail Kessler, Duluth
Promotion Manager
Linda Winick, Cleveland

OFFICES

ATLANTA
3091 Maple Drive
Maple Center One Building
Atlanta, GA 30305
(404) 233-1817

CLEVELAND
7500 Old Oak Boulevard
Cleveland, OH 44130
Editorial: (216) 243-8100

CHICAGO
111 East Wacker Drive
Chicago, IL 60601
(312) 938-2344

SEATTLE
1333 N.W. Norcross
Seattle, WA 98177
(206) 363-2864

DULUTH
120 West Second Street
Duluth, MN 55802
(218) 727-8511

MARKETING REPRESENTATIVES

Dick Gore
Atlanta (404) 233-1817

Ron Kempner
Atlanta (404) 233-1817

Joe Kosempa
Cleveland (216) 243-8100

Robert Mierow
Seattle: (206) 363-2864

UNDERCOVER AGENT

TURCAM[®] INSECTICIDE

For Use Only by Professional Applicators on
Ornamental Trees, Shrubs and Turf

ACTIVE INGREDIENT:	Percent by Weight
Bendiocarb: (2,2-dimethyl-1,3-benzodioxol-4-ol methylcarbamate*)	76%
INERT INGREDIENTS:	24%
Total	100%

*Protected by U.S. Patent No. 3,736,338

EPA REGISTRATION NO. 45639-59
EPA EST. NO. 45639-MI-1

KEEP OUT OF REACH OF CHILDREN WARNING

HARMFUL OR FATAL IF SWALLOWED STATEMENT OF PRACTICAL TREATMENT

If swallowed — Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to unconscious person. Get medical attention.

If inhaled — Remove patient to uncontaminated area. Keep patient strictly at rest and transfer to hospital immediately to get medical attention.

If on skin — Wash with soap and warm water.

If in eyes — Flush thoroughly with clean water.

NOTICE TO PHYSICIAN — Obtain a cholinesterase inhibitor as a pre-emptive measure to counteract the effects of the insecticide. It is suggested that the patient be given 1 mg/kg of atropine (1 mg/5 ml) of 1% solution intravenously every 30 minutes until the patient is fully atropinized.

An advanced carbamate insecticide that roots out and ruthlessly destroys white grubs, chinch bugs, sod webworms, mole crickets and other lawn and turf "terrorists". This tough operator doesn't get trapped in thatch, thus assuring positive grub control. TURCAM[®] is odorless...works well in spray equipment...won't damage turf or ornamentals. You'll find that TURCAM packs a federally-approved nationwide label, too.

Powerful reasons for assigning TURCAM to your toughest pest control problems.

For more information on TURCAM, including full labeling and recommendations for use, contact your local distributor or write to:

 **NOR-AM**

NOR-AM CHEMICAL COMPANY

3509 Silverside Road, P.O. Box 7495
Wilmington, DE 19803

*Registered trademark of Schering AG West Germany

Circle No. 130 on Reader Inquiry Card



CHEMICALS

Ortho takes aggressive marketing stance

The Ortho Division of Chevron Chemical Company, which is currently negotiating a takeover of Gulf Oil, has reactivated plans for construction of a fertilizer plant in Wyoming and restructured its Ortho marketing, manufacturing and research organizations to move its marketing management closer to its key markets, says Joseph Eckhart, general manager of the Ortho Agricultural Chemicals Division.

Ortho holds the lead position in consumer gardening products and indications are it intends to do the same in agriculture and professional horticulture. Expanded labels for Orthene and the construction of both a new fertilizer plant in Rock Springs, WY, and a research center in Richmond, CA, indicate the company is aware of the maturing nature of agriculture and the opportunity of specialty markets.

"We are focusing our future activities in research and development on innovative, highly-active products for major crop markets; and we are taking steps to significantly increase new product introductions," Eckhart told Weeds Trees & Turf. "Our market strategy is to improve our operating efficiency and response to customer needs."

Other chemical companies are using Ortho for consumer marketing.

EQUIPMENT

Thieves alter target to farm equipment

For more than a decade equipment thieves have favored construction equipment over farm equipment. That is until this year, says W.E. Rutledge, chairman of the heavy equipment theft committee of the International Association of Auto Theft Investigators.

"Because the construction indus-

try has been depressed, nobody's been working and the contractors were not buying heavy equipment," Rutledge explains. The Association's figures for Missouri show 167 pieces of construction equipment were stolen, compared to 451 pieces of farm equipment. The total loss in the state to large equipment theft is estimated at \$19.7 million. Missouri ranks fourth in the nation in large equipment theft.

Law enforcement officials have traced a large portion of the stolen

equipment to ports on the Gulf of Mexico. They complain that the equipment is easy to steal, hard to trace, and easy to sell because it is not registered.

With construction on the upswing, thieves might return to their old favorite, construction equipment. Also, since the average cost of stolen equipment is \$32,000, it makes sense to take extra precautions as business picks up and work schedules get complicated.



Dinah presented golf architect's award

Dinah Shore was presented the Donald Ross Award for her support of Women's golf during the American Society of Golf Course Architects annual meeting in Palm Springs, CA.

MEETINGS

Arborist reservations needed now for 1985

February 1985 seems like a long way off, but if you are an arborist planning to attend the National Arborist Association annual meeting on St. Thomas, Virgin Islands, you need to send your deposit to the hotel before the end of this month.

The meeting will be held at the Frenchman's Reef Beach Resort, Feb. 3-7, 1985. The area features tennis, golf, sailing, scuba diving, and, of course, a beach.


Bob Felix, NAA executive vice president, says early reservations are needed since there is no such thing as walk-ins on an island and the hotel must coordinate services with ships and planes. For information contact NAA at (516) 221-3082.

TREES

Deterioration of forests noted in Eastern U.S.

Air pollution is the chief suspect for
continued on page 12

1984 PGA putts on PENNCROSS



18th green at Shoal Creek, Shoal Creek, Alabama

Shoal Creek, Alabama

At the rugged Shoal Creek course, host of the 1984 PGA Championship August 16-19, 1984, the touring professionals will again be putting on Penncross creeping bentgrass. Penncross and Penneagle, the Penn Pals, are no strangers to the pros. Most of the recent major championships have been played on one or the other. The Western open at Butler was played on Penneagle fairways and greens. The 1983 US Open at Oakmont was played on Penneagle tees, fairways and greens. Maybe you saw the sixty-foot putt on Penneagle.

So, when the Shoal Creek course was designed, it's no wonder Penncross was specified for the putting surface. It was a 'natural' to choose Penncross bentgrass, with its glorious history, for a course with a sparkling future!

The PENN PALS



Penneagle's Oregon certified blue tags qualify for cash awards to qualifying turf organizations for turf research. Call our toll-free number for details. Marketed by **TEE-2-GREEN Corp.**, PO Box 250 Hubbard, OR 97032 1-800-547-0255 TWX 510-590-0957



IT PAYS TO HAND THIS KIND OF WORK TO ROUNDUP®

Hand-weeding areas like this doesn't make sense today. Not when Roundup® herbicide can do the job faster and better and give you more time to handle more customers.

Here's how: one-shot Roundup controls over 100 weeds right down to the roots, including poison ivy, poison oak and kudzu. One gallon of 2% spray solution can treat up to 30 tree rings for about a nickel each, or 1,000 feet of fenceline for less than \$2.00*.

It also pays to use Roundup when you're working around expensive ornamentals—because Roundup won't wash, leach or carryover in the soil. And

Roundup is virtually odorless and environmentally sound.

Use Roundup wherever you've been hand-weeding—around patios, flower beds, curbs, walkways, driveways, mulched areas, fences and steps. The choice is simple. You can spend valuable time hand-weeding. Or you can use Roundup to help you make more money. Now, that's using your head.

*Based on the cost of Roundup only to treat low growing vegetation.

FOR A FREE TRIMMING AND EDGING GUIDE
FOR ROUNDUP CALL TOLL FREE 800-621-5800.
IN ILLINOIS, CALL 800-972-5858.

ALWAYS READ AND FOLLOW LABEL DIRECTIONS FOR ROUNDUP.

Roundup® is a registered trademark of Monsanto Company.
© Monsanto Company 1984 RSP-4-104D



Monsanto

Circle No. 129 on Reader Inquiry Card



UPDATE

Latest NGF data is out

The National Golf Foundation's latest report shows 56 new golf courses were opened last year and 48 courses were expanded, for a total of 12,197 courses in the U.S.

Florida had the most new courses with 12, 11 connected with real estate developments, and a total of 30 new or expanded courses. Texas claimed second with ten new or expanded courses. Minnesota ranked third with eight, and Arizona, California, and Wisconsin each had six openings.

The future looks as bright with construction starts on 100 new or expanding courses last year. Another 102 are in the planning stage. Florida also leads in these categories, but Michigan and Illinois are showing up near the top of the starts and planning lists.

Growth of public courses has greatly outpaced private courses in the last ten years. Since 1973 the number of daily fee courses has increased by roughly 800 and municipal courses by 400. Meanwhile, the number of private course has increased by 90. Today, there are 4,809 private courses, 5,528 daily fee, and 1,860 municipal.

The number of golfers has increased by five percent in the past five years for a total of 17.8 million. For a complete copy of the report contact the National Golf Foundation, 200 Castlewood Drive, North Palm Beach, FL 33408.

GCSAA, NGF plan joint survey

A nationwide study of golf course maintenance activity will be conducted this summer jointly by the Golf Course Superintendents Association of America and the National Golf Foundation. GCSAA says superintendents of every course in the U.S. will be asked to share information on turfgrass maintenance practices, operating and capital budgets, and equipment use and needs. GCSAA has 5,000 members so the balance will have to come from NGF or other lists.

Questionnaires will be mailed this month. The returns will be compiled by the University of Kansas and results will be available by late summer or early fall, according to GCSAA. NGF President David Hueber says the survey is a pilot program in a long-range cooperative arrangement between the two organizations.

TPC family grows to five

The renovation of Edgewood Country Club, Cromwell, CT, under the direction of architect Pete Dye is nearing completion and renaming to TPC of Connecticut, the fifth club to be owned by the Tournament Players Club. The Greater Hartford Open will be played there in July. A sixth TPC course is under consideration in Potomac, MD. Active discussions are also ongoing in six other locations by TPC. The Tournament Players Club is rapidly becoming a major multi-course company and land owner.

deterioration of forests from the Ohio Valley to the East Coast. Studies by the U.S. Forest Service and Butler University in Indianapolis, IN, point to air pollution to reduced growth rates of conifers and poor foliage production of softwoods. Hardwood trees are also showing symptoms as measured by studies of the growth rings of trees.

Scientists liken the problem to a similar event in Germany where 35 percent of the forest declined or died more than 20 years ago. Trees monitored for timber and paper use have shown substantially less growth than expected during the past 15 to 25 years, says Dr. Authur Johnson of the University of Pennsylvania.

More specific blame has been placed in the Ohio Valley where many coal-burning plants are located. Dr. Orie Loucks, director of the Holcombe Research Institute, Butler University, described symptoms as discoloration of foliage on pine, tulip, poplar, sycamore, white ash, hickory, maples and black oaks. Also mentioned are narrower growth rings and a higher degree of mortality.

PEOPLE

Names in the news

Joe Troll made it 25 years of quarter-backing the Massachusetts Turfgrass Conference in March. Joe may be retiring soon and considering this year's conference, he is going out as a winner.

Bob Mullane, president of Alpine Tree Care, Inc., White Plains, NY, is the new president of the National Arborist Association. A University of Massachusetts graduate, Mullane said his goal is to reach out to others in the industry by attending at least ten industry shows. **Neil Engledow**, Mid-Western Tree Experts, Indianapolis, IN, will follow Mullane as president next year.

David K. Scatterday, has joined Martin Associates, Inc., Prairie View, IL, as a project manager. Scatterday is a landscape architecture graduate from the University of Illinois.

Lyle Borg, has been named sales manager for all Vermeer products by president Stan Vermeer. **Kevin Grooms** is the new industrial product sales manager for Vermeer, in charge of both tree and trencher products.

Gordon Mitchell brings 24 years of sales experience to Brouwer Turf Equipment as marketing manager. Gerry Brouwer, president, said Mitchell will be responsible for product introductions, dealer recruitment, advertising and promotion.

**A great ryegrass
is handsome, tough,
mows nicely and
germinates in 5-7 days.**



turf-type perennial ryegrass

is a great ryegrass.

Of course, the beauty, durability and mowing qualities of Derby turf-type perennial ryegrass aren't its only fine qualities.

Over the years, in a variety of locations and under widely varying conditions, Derby has proven itself a very consistent performer – a grass that will flourish without pampering.

Derby produces a dense, persistent turf when cut to $\frac{3}{16}$ inch for specialized uses such as golf greens, at standard cuts of 1 - 1½ inches for home lawns, parks and playgrounds, or $\frac{3}{4}$ inch on golf course fairways or tees.

It also mixes well with bluegrass or fine fescue, tolerates a wide range of soil types from heavy clay to sandy and retains its deep green color when used as a winter overseeding grass for dormant native Southern grasses.

Derby is registered with the Plant Variety Protection Office. PVA No. 7500009.



INTERNATIONAL SEEDS, INC.

P.O. Box 168, Halsey, Oregon 97348 U.S.A.

Telephone (503) 369-2251 TWX 510/590-0765

Circle No. 116 on Reader Inquiry Card

New study reveals local garden shows pay off for contractors

A new study by *Weeds Trees & Turf* reveals participation by landscape contractors in local garden shows adds to credibility and builds a small, but solid base of referrals during the off-season.

Contractors interviewed spend up to 250 hours and \$35,000 to design, set up, and tear down displays. Garden shows lend credibility to newer contractors, show homeowners the potential of landscaping, and help contractors reestablish contact with former customers and suppliers.

"People recognize immediately you are a good contractor," says Russell Jones, president of J. Franklin Styer Nurseries, Concordville, PA. "Any time you can achieve customer contact, it's very critical, especially in the off-season," Bob Kinney, vice president, Wilmore Gardens, Denver, CO, stressed. "One good customer gained from a show can turn around and give you 25 more customers in the next year," according to Lambeth Marshall, president, Associated Landscape, Inc., Hoyt Court, NC.

Undeniably leads from these shows are residential, but corporate executives have been known to make impressions about contractors while attending garden shows in their area. The complete report will be published in an upcoming issue of *Weeds Trees & Turf*.

Poor acclimation suspected for problems with large ficus

Survival problems with large *Ficus* trees transplanted to interior landscapes is now being linked to inadequate acclimation of field-grown trees to containers by the Interior Landscape Division of the Associated Landscape Contractors of America. Originally, the marl soil the trees were grown in was suspected. ALCA reported the problem this past fall and has tried to find a common link among problem trees.

ALCA says a process of root pruning, container size, proper digging and acclimation all effect the tree's ability to survive on location. The best tree is one that has apparently been growing in a container rather than in the field prior to installation.

Contest entry time is here

Summer is the time to enter many landscape award contests; including ALCA's Fifteenth Annual Environmental Improvement Awards, the Florida Nurserymen and Growers Association Fourteenth Annual Landscape Awards, and the California Landscape Contractors Association Trophy Awards.

The keys to winning landscape awards are knowing the job may be award material before it is started, good before and after photography, thorough records of the job as it progresses, testimony from the owner as to the impact of the project, following the contest rules closely, and presenting the entry material in a neat and attractive fashion.

Mike Bush has been appointed manager of Cypress Gardens' Nursery Division. Bush hopes to expand the attraction's existing collection of rare tropical and subtropical specimen plants. "I believe there is a wealth of plant material in places like Brazil, Argentina, and Paraguay that has yet to be exploited," Bush said.

Exmark Mfg. Co., Beatrice, NE, has promoted Dick Tegtmeier to executive vice president. Bob Martin, president, said the promotion was the result of "remarkable" growth for the fairly new mower manufacturer.

LESCO Inc. has named two territory managers. Les Guedel will be manager of East Coast sales and Jim Johnson will manage North Central sales for the company based in Rocky River, OH.

Ray Lewis and his wife returned from the Las Vegas GCSAA Show with some unpredicted baggage, a Dedoes trailer aerator. The Lewis couple won a drawing during the show.

JOB HUNTING

Turf and landscape recruiter opens doors

A firm specializing in recruiting personnel for the turf and landscape industries has been launched in Littleton, CO. S. Ronald Gaston and Associates will handle national fee-paid positions for golf course superintendents, landscape architects, and other professional landscape positions.

Ron Gaston says his firm is the first of its kind. A pamphlet describing the new service is available by writing the company, P.O. Box 2527, Littleton, CO 80161.

TURF SEED

Danish company's U.S. branch renamed

Daehnfeldt, Inc., is the new name of Pacific Seed Production in Albany, OR, after the Danish seed company decided to clear up any confusion about its presence in the U.S.

L. Daehnfeldt of Odense, Denmark, signed a cooperative agreement with Pacific Seed in 1982 to develop a joint Danish/U.S. breeding and production company. Many European seed companies are active in the seed growing regions of the northwestern U.S., often through U.S. companies. Growing conditions and land are more favorable in the U.S. than in Europe.

The company will continue research and development of turf, for-

GET THE FEELING THEY KNOW SOMETHING YOU DON'T?



They do. It's their job to know things first and then pass that information on to you fast. Things like new turf management techniques, effective methods of insect and weed control, what really works... and what won't.

LOOK FAMILIAR?

They should. You've seen them often at turf shows, seminars, conventions, wherever industry news is in the making. They're the editorial and sales management team of **WEEDS TREES & TURF**. Please meet (seated l. to r.) Ron Kempner, Bruce Shank, Dick Gore, and Maureen Hrehocik; (back row l. to r.) Bob Mierow, Kevin Cooney, Joe Kosempa, and Bob Earley.


THERE'S NO GRASS GROWING UNDER THEIR FEET.

They apologize for not always being in when you call, but great stories are found in the field, not in the office. These pros know the only way to really cover the green industry is to get out and be a part of it. From turf test plots in Oregon to the corridors of our Nation's Capital, they're following leads, surveying markets and interviewing the landscape

professionals who make our industry hum. Then, they use their years of experience to pull it all together, analyze it and present it to you in a crisp, easy-to-read and easy-to-use style. That way, you have the information you need to do your job better, faster and more effectively.

The next time you want to know something in the green industry, give them a call. If you can't reach them at the office, don't worry. They'll reach you in the pages of **WEEDS TREES & TURF**.

WEEDS TREES & TURF

 A HARCOURT BRACE JOVANOVIICH PUBLICATION

3091 Maple Drive, Suite 312, Atlanta, GA 30305
(404) 233-1817

— A GREAT MEDIUM TO GROW PROFITS IN. —



New Du Pont Landscape Fabric. It lets water through to give you healthier beds with less work.


New DuPont Landscape Fabric lets water pass through, reduces wash-away of mulch while it impedes weed growth. So you get healthier, more attractive plant beds with less maintenance work and cost.

Everything you apply for bed care gets to plant roots in the amount you want, where you want it. Water, fertilizers, herbicides and pesticides seep down through this chemically inert fabric to nourish and protect every plant in your bed.

DuPont Landscape fabric is easy to put down with scissors or knife. It comes in four roll sizes from 3- to 12-feet wide.

Call 800-441-7517 for the name of the nearest distributor and more information about DuPont Landscape Fabric. Or write DuPont Company, Room G-40955, Wilmington, DE 19898.

Circle No. 150 on Reader Inquiry Card



Circle
the
Reader
Service
numbers
of those
items of
interest
to you.

For fastest response, use the peel-off label from the front cover.

NAME	_____
TITLE	_____
FIRM	PLACE COVER LABEL HERE
ADDRESS	_____
CITY	_____
STATE	ZIP

101	116	131	146	161	176	191	206	221	236	251	266	281
102	117	132	147	162	177	192	207	222	237	252	267	282
103	118	133	148	163	178	193	208	223	238	253	268	283
104	119	134	149	164	179	194	209	224	239	254	269	284
105	120	135	150	165	180	195	210	225	240	255	270	285
106	121	136	151	166	181	196	211	226	241	256	271	286
107	122	137	152	167	182	197	212	227	242	257	272	287
108	123	138	153	168	183	198	213	228	243	258	273	288
109	124	139	154	169	184	199	214	229	244	259	274	289
110	125	140	155	170	185	200	215	230	245	260	275	290
111	126	141	156	171	186	201	216	231	246	261	276	291
112	127	142	157	172	187	202	217	232	247	262	277	292
113	128	143	158	173	188	203	218	233	248	263	278	293
114	129	144	159	174	189	204	219	234	249	264	279	294
115	130	145	160	175	190	205	220	235	250	265	280	295

WEEDS TREES & TURF

JUNE 1984

This card expires August 15, 1984

MY PRIMARY BUSINESS AT THIS LOCATION IS:
(PLEASE CHECK ONE ONLY IN EITHER A, B OR C)
**A. LANDSCAPING/GROUND CARE AT ONE OF THE FOLLOWING
TYPES OF FACILITIES:**

- 0005 Golf courses
 0010 Sport complexes
 0015 Parks
 0020 Rights-of-way maintenance for highways, railroads & utilities
 0025 Schools, colleges & universities
 0030 Industrial & office parks/plants
 0045 Condominiums/apartments/housing developments/hotels/resorts
 0050 Cemeteries/memorial gardens
 0060 Military installations & prisons
 0065 Airports
 0070 Multiple government/municipal facilities
 Other type of facility (please specify) _____

B. CONTRACTORS/SERVICE COMPANIES/CONSULTANTS:

- 0105 Landscape contractors (installation & maintenance)
 0110 Lawn care service companies
 0125 Landscape architects
 0135 Extension agents/consultants for horticulture
 Other contractor or service
 (please specify) _____

C. SUPPLIERS:

- 0205 Sod growers
 0210 Dealers, Distributors
 Other supplier (please specify) _____

Approximately how many acres of vegetation do you maintain or manage? _____

What is your title? (please specify) _____

I would like to receive (continue receiving) WEEDS TREES & TURF
each month: YES NO

Your Signature: _____ Date: _____



BUSINESS REPLY CARD

FIRST CLASS PERMIT NO. 665 DULUTH, MINNESOTA

POSTAGE WILL BE PAID BY ADDRESSEE

READER SERVICE DEPARTMENT

WEEDS TREES & TURF

POST OFFICE BOX 6049
DULUTH, MINNESOTA 55806-9749

NO POSTAGE
NECESSARY
IF MAILED
IN THE
UNITED STATES



**GET
MORE
FACTS**

age, vegetable, and flower seed varieties, both in Denmark and in Oregon. Daehnfeldt plans "substantial growth and expansion" to expedite sale of supplies from its Denmark operation in the U.S.

INSECTICIDES

Mocap receives label for six more turf insects

Rhone Poulenc has received EPA approval to add six turf pests to its label for Mocap 10 percent granular. First registered for use against nematodes and mole crickets, the label now includes sod webworms, chinch bugs, Japanese beetle, black turfgrass ateanius, European chafer, and the bluegrass billbug.

The application rate for the six pests is half that for mole crickets, 1.25 lbs./acre. Only professional turfmen are permitted to apply the product for the new pests.

"The expanded label shows Mocap's ability to control a broad spectrum of insects, including grubs," says Dan Stahl, turf product manager for Rhone Poulenc.

ASSOCIATIONS

Agriculture secretary to give AAN awards

Secretary of Agriculture John R. Block will host a reception for the winners of the American Association of Nurserymen's 1984 Landscape Awards competition after the awards are presented at a White House ceremony, on May 17.

A custom established by Lady Bird Johnson is for the awards to be presented by the President or the First lady in the Rose Garden. Mrs. Reagan presented the awards to the winners in the White House in 1982.

The awards will be presented two months before the 109th Annual AAN Convention in San Antonio, TX, July 14-17. Lady Bird Johnson will address the conference during the meeting. Mrs. Johnson continues to take a leadership role in beautification of parks and roadsides.

SURVEY

Ontario spends \$275 million on turf

A survey, intended partly to justify expansion of turf extension programs
continued on page 80

GOVERNMENT

UPDATE

Correction on Monsanto/EPA suit

The suit reported in the March issue should not be confused with the patent suits Stauffer has filed against Monsanto.

First, there is no suit between Stauffer and Monsanto in the Supreme Court. The Supreme Court suit is between Monsanto and EPA only. Secondly, Stauffer should have been listed as one of those who has filed an Amicus curiae brief in support of Monsanto's position in the Supreme Court case against the EPA. Finally, Stauffer's opposition is that "Me-to" registrations should be nullified.

Johnson says major changes in FIFRA unlikely this year

In an exclusive letter to *Weeds Trees & Turf*, Edwin Johnson, Director, Office of Pesticide Programs, EPA, said, "Substantive FIFRA amendments are unlikely (this year). Rather than pursuing legislative changes this year, the Administrator is seeking administrative changes and remedies to pesticide regulatory problems through the newly formed Pesticide Advisory Committee."

Simazine receives soil restriction

After Dec. 31, 1984, labels on simazine products will bear a restriction against use of the product in areas where groundwater contamination is likely. This generally means it can not be applied to soil in areas designated Sole Source Aquifers by the Soil Conservation Service.

Surflan has experimental turf label

Elanco's Surflan, currently registered for selective pre-emergence control of annual grasses and broadleaf weeds around ornamentals, has received an experimental use permit for weed control in 16 Southern states.

EPA's Moore claims restrictions needed to prevent contamination

John Moore, assistant administrator for Pesticides and Toxic Substances, claims some pesticides will have to be banned or restricted to protect groundwater from contamination. Moore suggested that EPA could point out which pesticides are potential ground water polluters and let the states handle restrictions. "Soil fumigation technology needs to be rethought," said Moore. Reregistration is taking too long to adequately protect groundwater according to Moore.

Get ready for renovation work

Much of July's work is follow-up on spring work. It is when insecticide, herbicide, and fungicide applications are renewed, evergreens are pruned and shaped, and irrigation is closely monitored.

From a planning standpoint, July is when you should nail down fall renovation work and order the necessary supplies. Do not assume everything you need for renovation work will be available. Sell renovation work early in July and notify your suppliers as soon as possible of your needs. Winterkill replacements earlier in the year and reported shortages in some turf seed may limit the renovation work you can do this year.

Labor planning is especially important, considering how valuable renovation work can be while, at the same time, seasonal labor is departing. Quite often renovation work in August and the fall is more important than work during the busy season. By August, seasonal crews are skilled and more efficient than they would be the following spring. Work is therefore more efficient from both labor and plant standpoints.

Chemical renewal

Preemergence herbicides in plant beds and container plantings may need to be reapplied now. Check the label for the length of time the pre-emergence herbicide you use is effective.

Second applications of turf insecticides may also be needed in July to control severe grub, chinchbug, and greenbug infestations.

Many foliage-feeding and sucking insects are at harmful levels in July. Injury during the summer reduces carbohydrate storage by plants leaving plants more vulnerable to winterkill and reducing bud development. Serious foliage feeders include Japanese beetle, gypsy moth, black vine weevil and bag worms. Control is most effective when insects are small. Controls include Sevin, Turcam, Orthene, Diazinon, and methoxychlor. Sucking insects include spider mites, aphids, white flies, and lace bugs. Mite control requires Dicofol, dymet, or Vendex. Malathion, Orthene, Diazinon, dimethoate and Sevin may be used for the other sucking insects. These materials are also effective against the crawler stages of scale insects active at this time.

Hot, humid weather encourages a number of turf diseases, including pythium, anthracnose of *Poa*

annua, *Fusarium* blight, and brown patch. Since poor drainage is a prime cause of these diseases, make a note to improve drainage and air flow of problem areas this fall. Turf fungicides typically have residual periods of less than three weeks. Bayleton, a fungicide with season-long residual must be applied before diseases start, usually sometime in early June. Not all fungicides are effective on all turf diseases. Check with your chemical supplier, the Disease Control Guide in this issue, or local extension agent for specific control information.

Most ornamental disease control is based on control beginning prior to or following bloom. Roses require treatment every two weeks during the summer for black spot and powdery mildew. Preventative spray programs should be established for susceptible plants.

Irrigation

The role of irrigation in landscape management is being closely evaluated today. Water-sensing devices called tensiometers are being added to irrigation systems to reduce water use. Drip irrigation can greatly reduce water use for ornamentals. Wetting agents are gaining acceptance to correct localized dry spots. Plant breeders are selecting plants which require less water. Water use is no longer taken lightly.

As mentioned in a previous issue, Dr. Joe Vargas, plant pathologist from Michigan State University, is recommending light, mid-day irrigation to encourage an active thatch layer. Vargas believe beneficial organisms, which aid in thatch and disease control, are most effective if the thatch layer is not allowed to dry out. Irrigation should be brief, just to moisten the thatch, at mid-day so leaf tissue does not remain damp for long periods.


Dr. Robert Shearman of the University of Nebraska, is recommending irrigation at the first signs of wilt. Then, Shearman recommends a deep soaking to encourage deep rooting and lower disease potential.

Pruning conifers, pines

The best time to prune conifers and pines is July. Shearing half the candle growth results in a denser plant, and if done every year, produces a dwarf plant. This is helpful to keep plants within their intended size and shape in the landscape design.

Juniper, yew, and privet hedges can be shaped at this time. For best foliage appearance, the lower branches should be longer than the top so that all foliage receives adequate sunlight. Otherwise, lower branches will have sparse foliage. **WT&T**

Landscape Log is written based on previous publications by Dow Gardens Horticulturist Douglas Chapman, the Weed Control from March 1983, and the Disease Control Guide in this issue.



HOW TO IMPROVE YOUR TOUCH AROUND THE GREEN.

PRESENTING THE GREENSMASTER 3.


Of all the mowing jobs on a golf course, mowing the green may require the most delicate touch of all. To give each green a uniform cutting height, appearance and playing characteristic while maneuvering in a tightly defined area.

That's why our Greensmaster 3 has a three-wheel design that allows it to turn within its own length. So you can easily cut up to and around the edges of greens, collars and traps.

And to improve your touch around the green, we've added features to the Greensmaster 3 to make it even more maneuverable.

A new single forward/reverse traction pedal has been added to give the operator greater control in maneuvering this tight-turning three-wheeler.

In addition, the operator station features easy-to-reach controls mounted on a steering



arm that adjusts to the operator's height. And the high-backed cushion seat is also adjustable. So the operator enjoys both greater comfort

and greater command of the machine.

Along with superior maneuverability, the Greensmaster 3 gives a consistently smooth, even cut.

It is still the only greensmower designed with its cutting units isolated from both the traction unit and the grass baskets. Which means it will follow the contours of the green for a superb cut without scalping or mismatching. And deliver a uniform height of cut from 3/32" to 11/16".

Finally, the Greensmaster 3 is still built with the rugged Toro durability that keeps you cutting years longer.

So call your Toro distributor about the Greensmaster 3. The agile mower that gives a quality of cut second to none.



TORO

THE PROFESSIONALS THAT KEEP YOU CUTTING.

"Toro" and "Greensmaster 3" are registered trademarks of the Toro Company, 8111 Lyndale Ave. So., Minneapolis, Minnesota 55420.

Circle No. 141 on Reader Inquiry Card

JUNE 1984/WEEDS TREES & TURF 21

TURF DISEASE GUIDE

Two leading plant pathologists, J.M. Vargas Jr. of Michigan State University and Don Blasingame of Mississippi State University, cut through confusing terminology and present the latest turf disease control methods. Diseases of cool-season turfgrasses, warm-season turfgrasses, and overseeded ryegrasses are discussed. This is a section you'll want to save. An added help is the Turf Fungicide Directory on page 44.

Northern Turf Diseases

by J. M. Vargas, Jr., turfgrass pathologist, Michigan State University

Concepts about turfgrass diseases and their management have gone through many changes in the past few years, including the scientific names of the organisms that cause them.

These diseases, the organisms that cause them, and their cultural, biological and chemical management tools are given in Table 1. The following will be a discussion of the latest developments on cool-season turfgrass diseases.

Dollar spot

Dollar spot is primarily a disease of golf course grasses such as creeping bentgrass and annual bluegrass. It is now believed to be caused by two organisms, a *Lanzia* spp. and a *Moellerodiscus* spp.

Now that two fungi have been identified as the cause of dollar spot, it helps explain some of the confusion that has existed about the occurrence of this disease. It has been considered both a cool weather disease and a warm weather disease. It appears that

dollar spot is both, and that there are two different fungi which caused a disease with similar symptoms. This means that you can have dollar spot at any temperature between 60-85° F.

What is needed is an easy, reliable method to distinguish the difference between these two fungi in the field. Fortunately, dollar spot caused by both fungi appears to be reduced by adequate nitrogen levels and adequate soil moisture levels. For the most part, they are also managed by the same fungicides, although one has to wonder if some of the resistance to some fungicides might not be due to the differential sensitivity of the two fungi species that causes this disease.

Brown patch

Brown patch is also primarily a disease of golf courses, although with the new improved perennial ryegrasses being incorporated into home lawn mixtures, it is also becoming a problem on home lawns.



The disease occurs under hot, humid conditions. It can be culturally managed by reducing the amount of nitrogen applied just prior to the advent of warm weather, increasing air circulation by removing trees or shrubs, and/or by pruning them.

Pythium blight

Pythium blight is also a disease of golf courses, and like brown patch, it is becoming more of a home lawn problem with the incorporation of the improved perennial ryegrass.


There still seems to be some controversy over how many species of pythium are involved in this disease, but regardless of how many or how few there are, they do tend to cause rapid loss of turf in hot, humid weather.

Unlike many diseases where only the foliage is damaged and recovery occurs soon after, Pythium blight usually kills the plant. This means recovery in the infected areas will be slow because it will have to come from

Table 1. TURF DISEASES AND CONTROLS



Pythium blight rapidly strikes ryegrasses and other turfs where drainage is poor in hot and humid weather. Recovery is slow since pythium usually kills the entire plant.



rhizomes or stolons filling in from outside the spots or by germination of annual bluegrass or broad-leaf weeds when the cool weather of the fall returns.

Cultural management of Pythium blight consists of reduced nitrogen levels just prior to the advent of warm weather and improving drainage.

Concerning drainage, in marginal areas of the cool-season grass regions, Pythium blight is only a problem in areas of poor soil drainage, where water stands for prolonged periods. In regions where severe Pythium blight damage occurs, it is always most severe in poor drained soil areas. It goes without saying, that good Pythium blight management begins with improving soil drainage.

As far as chemical management is concerned, there are two systemic fungicides to manage Pythium blight, which are metalaxyl (Subdue, Ridomil, Apron) and propamocarb hydrochloride (Banol). These two fungicides will manage the disease for up to three

Dollar spot damages bentgrass, annual and perennial bluegrass, bermudagrass and zoysiagrass. In the South it is common in the spring and fall. In the north it is active anytime temperatures are between 60 and 85°.

weeks. They appear to be slower acting than chloroneb (Teremec SP) or ethazol (Koban, Terrazole). Little spread of the disease occurs after these systemic fungicides are applied, although the mycelium of the fungus may remain evident on the previously infected tissue for a couple of days. No actual resistance to these two fungicides has been reported, but the possibility exists. It would be wise therefore, to follow each systemic fungicide application with contact (chloroneb or eth-

azol) in case resistance does occur following a systemic fungicide application, the contact fungicide will prevent the resistant strain from devastating the turf.

Anthracnose

Anthracnose caused by *Colletotrichum graminicola* is primarily a disease of annual bluegrass, although it will attack the fine leaf fescues, perennial ryegrasses and seaside creeping bentgrass.

Annual bluegrass dies from an-

TURF

Table 1: TURF DISEASES AND CONTROLS*

Disease	Causal Agent	Hosts	Cultural Control	Chemical Control
Anthrachnose	<i>Collectotrichum graminicola</i>	Annual bluegrass Fine-leaf fescue Kentucky bluegrass Perennial ryegrass	Adequate nitrogen. Cool grass by syringing	Maneb plus zinc sulfate, chlorothalonil, benomyl, thiophanate-methyl, thiophanate, thiophanate-methyl + mancozeb, triadimefon
Brown patch	<i>Rhizoctonia solani</i>	All major turfgrass species	Reduce nitrogen. Remove "dew." Increase air movement.	Mancozeb, maneb + zinc sulfate, chlorothalonil, anilazine, cycloheximide + thiram, benomyl, thiophanate-methyl, thiophanate, thiram, thiophanate-methyl + maneb, cadmium compounds, thiophanate + thiram, PCNB, iprodione, vinclozolin
Dollar spot	<i>Lanzia spp.</i> <i>Moellerodiscus spp.</i>	Annual bluegrass Bahagrass Bermudagrass Centipedegrass Colonial bentgrass Creeping bentgrass Fine-leaf fescues Kentucky bluegrass Perennial ryegrass St. Augustinegrass Zoysiagrass	Increase nitrogen. Remove "dew".	Benomyl, thiophanate, thiophanate-methyl, chlorothalonil, anilazine, cycloheximide + PCNB, cadmium compounds, thiophanate + thiram, thiram, thiabendazole, benomyl, iprodione, thiophanate-methyl + maneb, vinclozolin, triadimefon
Fusarium blight syndrome		Kentucky bluegrass Centipedegrass	Light, daily watering during the summer.	thiophanate-methyl, thiophanate, triadimefon
Helminthosporium Diseases Brown blight Leaf blotch Leaf spot	(<i>Dreschlera</i>) <i>D. siccans</i> <i>D. cynodontis</i> <i>D. sorokinianum</i>	Ryegrass Bermudagrass Bentgrass, Fine-leaf fescue, Kentucky bluegrass	Remove clippings. Raise cutting height. Plant resistant cultivars. Moderate spring nitrogen. Daily irrigation	Mancozeb, chlorothalonil, cycloheximide, anilazine, maneb + zinc sulfate, cycloheximide + thiram, cycloheximide + PCNB, iprodione, vinclozolin
Melting-out Net-blotch Red leaf spot Stem and Crown Necrosis Zonate eye spot	<i>D. poae</i> <i>D. dictyoides</i> <i>D. erythrospilum</i> <i>D. spiciferum</i> <i>D. giganteum</i>	Kentucky bluegrass Fescue Creeping bentgrass Bermudagrass Bermudagrass		
Gaeumannomyces patch (Take all patch)	<i>Gaeumannomyces graminis</i>	Annual bluegrass Colonial bentgrass Creeping bentgrass Kentucky bluegrass Tall fescue Velvet bentgrass	Reduce soil pH. Avoid liming. Use acidic fertilizers. Sulfur.	None.
Pythium blight (cottony blight)	<i>Pythium spp.</i>		Improve soil drainage. Increase air circulation.	Chloroneb, ethazole, metalaxyl, propamocarb, hydrochloride
Red thread	<i>Laetisaria fusiformis</i>	Creeping bentgrass Colonial bentgrass Bermudagrass Annual bluegrass Perennial ryegrass Fine leaf fescues	Increase nitrogen	anilazine, iprodione, triadimefon, vinclozolin, chlorothalonil
The Snow Molds <i>Typhula</i> blight <i>Gerlachia</i> patch	<i>Typhula spp.</i> <i>Gerlachia nivalis</i>	Annual bluegrass Colonial bentgrass Creeping bentgrass Fine-leaf fescues Kentucky bluegrass Perennial ryegrass Tall fescue Velvet bentgrass	Avoid early fall nitrogen fertility that leads to lush growth.	Mercury compounds, PCNB products, chlorothalonil, chloroneb. These products may have to be used in combination for effective snow mold management. Benomyl, Iprodione or Mancozeb will control <i>Gerlachia</i> patch where it occurs alone.
Yellow patch	<i>Rhizoctonia cerealis</i>	Kentucky bluegrass Creeping bentgrass	Nitrogen to promote recovery.	Iprodione

*The order in which fungicides are presented does not imply the order of their effectiveness.

thracnose during heat stress periods of the summer. This is not due to heat alone, as was once believed. If proper cultural management is followed, and effective fungicides are used, annual bluegrass will survive the summer heat stress period like any other perennial.

Good cultural practices consist of deep vertical mowing early in the spring, as soon as growth is initiated for the season. This will allow for the production of new juvenile growth which should be more resistant to heat stress. This should be followed by coring a week or two later for good root growth. A second coring should be made following heavy seedhead production in the late spring to provide an optimum medium for maximum root growth (the coring holes) in the few remaining weeks prior to the heat stress period. More than 70% of the annual bluegrass roots disappear during heavy seedhead production. If only one coring a year can be done, this is the time to do it. A third coring should be made when the cool nights return in late summer and early fall.

Light nitrogen application should be made (i.e. 1/2 lbs. actual nitrogen per 1000 sq ft.) in June, July and August. This produces healthier annual bluegrass and reduces the amount of inoculum produced by *C. graminicola* for subsequent infection during the remainder of the anthracnose season. Finally, one of the effective fungicides should be used to insure healthy turf.

A computer model has been developed, which predicts the occurrence of anthracnose based on average daily temperatures and continuous hours of leaf wetness. Fungicides for the management of anthracnose can now be applied when the disease occurs instead of on a calendar basis.

Snow molds

There are two prevalent snow molds in the United States: *Typhula blight* (gray snow mold)

and *Gerlachia patch* (pink snow mold).

Gerlachia patch *Gerlachia patch* is caused by *Gerlachia nivalis*, formerly known as *Fusarium patch* caused by *Fusarium nivale*. Yes, another name change and this time, they not only changed the scientific name of the organism causing the disease, but the common name of the disease as well!

The disease becomes a problem in the fall when the temperature drops into the low 60's and continues through the spring, until daytime temperature climbs back into the seventies. It is usually first noticed in the shaded areas of the green, tees and fairways.

Gerlachia patch does not need snow cover to become active, only the cool wet weather. Annual bluegrass is especially susceptible to *Gerlachia patch*.

In the spring the disease is often misdiagnosed as copper spot, because of the small copper-colored spot that it causes. However, copper spot is a disease that occurs in warmer weather. Keeping nitrogen at low levels during the time when *Gerlachia patch* may be active is important in helping manage the disease

Typhula blight

Typhula blight is caused by two species, *Typhula incarnata* and *T. ishihariensis*. *T. incarnata* is the primary species in eastern, southern, and regions of the midwest and western U.S. *T. ishihariensis* is most prevalent in the more northern snow mold regions, especially where prolonged periods of permanent snow (two or more months) exist in the mid-western and western U.S.

The two *typhula* species are easily distinguished from each other when observed soon after the snow melts. *T. incarnata* produces grayish spots in the turf, with fairly large uncommon brown sclerotia (a mass of filaments) evident. Whereas, *T. ishihariensis* spots have a reddish cast and contain small, dark black

sclerotia. *Typhula blight* only occurs under snow cover. It does not occur in the cool wet weather of fall and spring, except under leaf piles.

Knowing which species you have is important in chemically managing the disease. Many fungicides, including the mercuries, chloroneb (Teremec SP), PCNB (Terraclor), triadimefon (Bayleton), and chlorothalonil (Daconil) will manage *Typhula blight* caused by *T. incarnata*. They do not all manage *Typhula blight* caused by *T. ishihariensis*.

The picture also is more confusing state by state. For example, in Michigan, the mercuries, PCNB, and chlorothalonil will manage both species, but triadimefon and chloroneb will not manage *Typhula blight* caused by *T. ishihariensis*. In northern Wisconsin and Minnesota, combinations of the mercuries and PCNB are required to manage both species. You should check with your local turfgrass experts to find out the fungicides that are effective in your area.

Gaeumannomyces patch

Gaeumannomyces (take all) patch caused by *Gaeumannomyces graminis* var. *avanaea* was formerly known as *Ophiobolus patch* caused by *O. graminis*. This disease was originally thought to be confined to the Pacific northwest. It has now been reported in the coastal areas of New England, New York, and the mid-Atlantic states, primarily on creeping bentgrass turfs. A *Gaeumannomyces* like organism has also been reported on annual bluegrass in the mid-eastern and mid-western U.S.

In 1983, the disease caused widespread destruction on many annual bluegrass fairways in mid-August and early September. So, *Gaeumannomyces patch* or closely related diseases are now occurring through most of the cool-season grass regions of the United States.

TURF

Lowering the pH through the use of sulfur still appears to be the best way to manage this disease. A word of caution is necessary, the granular sulfur products have been observed to cause injury to the turf the season following application. This injury initially resembles dollar spot. The sprayable sulfurs are just as effective and do not have the bad side effects.

Fusarium blight

There are two current schools of thought on the cause of Fusarium blight. The research at Penn State University (Cole) suggests that a basidiomycete (a type of higher fungi) is involved in causing the "frog-eye" symptom associated with Fusarium blight and that, if the Fusarium fungi are involved at all, they are involved as saprophytes colonizing the dead and dying tissue.

The other school of thought, represented by Cornell University (Smiley) suggests the cause of the "frog-eye" is due to two fungi, *Leptosphaeria korrae* and/or *Phialophora graminicola*. It could be we are actually dealing with three different fungi causing three different diseases, all of which have the same symptom. Time will tell which of these schools of thought are correct or if they both are.

Fusarium blight is a warm weather disease that occurs from late June through early September depending on your location. The disease usually occurs after a week or two of dry weather following a heavy rain.


The characteristic initial symptom is wilted turf in the infected spot. This separates it from six other diseases that have similar "frog-eye" symptoms, like brown patch and yellow patch.

Since there may be as many as

three fungi involved in the Fusarium blight syndrome, it is difficult to make specific recommendations to encompass all of them. The following are the best management recommendations available, although slight variations may exist in different areas of the country.

Cultural management Coring should be done to improve root development, reduce thatch, and eliminate layering caused by two different soil types. Homelawn turf is often grown on poor soil. Many times sod is laid on compacted subsoil because the topsoil was removed during construction.

Nutrient and water uptake are active processes which require adequate oxygen. Coring holes provide an excellent area for root growth with good aeration for proper uptake of nutrients and water.



OTTERBINE DAY OR NIGHT...SUMMER OR WINTER



OTTERBINE® Aerators can help you keep unsightly algal growth and objectional odors under control naturally. The fact that OTTERBINES also create beauty—is just one of the many benefits of using our Spray Sculpture™ Floating Fountains.

**WE CREATE BEAUTY
WHILE SOLVING PROBLEMS**

BAREBO, INC. P.O. BOX 217 • EMMAUS, PA 18049
(215) 965-6018 USA



Circle No. 101 on Reader Inquiry Card



Ad
Council

Photo: Peter B. Kaplan

If you still believe in me, save me.

For nearly a hundred years, the Statue of Liberty has been America's most powerful symbol of freedom and hope. Today the corrosive action of almost a century of weather and salt air has eaten away at the iron framework; etched holes in the copper exterior.

On Ellis Island, where the ancestors of nearly half of all Americans first stepped onto American soil, the Immigration Center is now a hollow ruin.

Inspiring plans have been developed to restore the Statue and to create on Ellis Island a permanent museum celebrating the ethnic diversity of this country of immigrants. But unless restoration is begun now, these two landmarks in our nation's heritage could be closed at the very time America is celebrating their hundredth anniversaries. The 230 million dollars needed to carry out the work is needed now.

All of the money must come from private donations; the federal government is not raising the funds. This is consistent with the Statue's origins. The French people paid for its creation themselves. And America's businesses spearheaded the public contributions that were needed for its construction and for the pedestal.

The torch of liberty is everyone's to cherish. Could we hold up our heads as Americans if we allowed the time to come when she can no longer hold up hers?

Opportunities for Your Company.



You are invited to learn more about the advantages of corporate sponsorship during the nationwide promotions surrounding the restoration project. Write on your letterhead to: The Statue of Liberty-Ellis Island Foundation, Inc., 101 Park Ave, N.Y., N.Y. 10178.



Save these monuments. Send your personal tax deductible donation to: P.O. Box 1986, New York, N.Y. 10018. **The Statue of Liberty-Ellis Island Foundation, Inc.**

TURF

Thatch reduction is best accomplished during the coring operation by breaking up the cores with a vertical mower or power rake, and incorporating the soil back into the thatch layer. Power raking does little for thatch reduction. It removes leaf tissue which

is readily broken down but does nothing to remove the rhizomes and roots which are primarily responsible for thatch formation.

Layering results from one soil of a different type being placed on top of the other, as when a muck sod is placed on mineral subsoil.

In the cool weather of spring and fall it may not be a problem, but under stress conditions of the summer it can become a serious problem. The entire turfgrass root system is restricted to the upper layer during the summer heat stress period. This often means the root systems are no more than an inch in depth.

Obviously, drought stress diseases like "Fusarium blight syndrome" are going to be more severe under such conditions.

Integrating two soil layers over a period of years through a coring program should make for a deeper rooted, healthier turf.

Fertility Nitrogen fertility in the summer months of June, July and August, will reduce the severity of the "Fusarium blight syndrome". Approximately 1/2 lb. of actual nitrogen/1,000 sq ft./month should be adequate.

Irrigation Supplemental irrigation can culturally reduce "Fusarium blight syndrome" if applied on a daily basis. If applied at mid-day it will cool the plants, similar to syringing performed on golf courses during the heat stress period. It also provides water for the short and limited root systems of the infected plants.

If the mat or thatch is kept moist, antagonistic microorganisms may develop, which will prevent the pathogenic fungi from attacking the plants. A daily irrigation program during the summer on infected turf may also cause the build-up of antagonistic microorganisms that destroy the "Fusarium blight syndrome" fungi.

Chemical management Thiophanate(Cleary's 3336), thiophanate-methyl(Fungo 50) and benomyl(Tersan 1991), are good fungicides for the management of the "Fusarium blight syndrome". They all have the same basic chemistry.

The turf area to be treated should be irrigated the night before and the fungicides drenched in before they dry on the foliage. They can be used either cura-

Trailer aerator/dump box package by Dedoes: easy, quick and effective.



You can be sure they'll work for you. Can be pulled by most tractors including International, Toro, Ford, Massey Ferguson, Yanmar, Kubota, John Deere, E-Z-Go, and Cushman, because Dedoes trailer aerator and dump box attach to your hitch.



Optional Dump Box

Newly designed remote control hydraulics for easy hookup and stable saddlebag weight boxes for proper penetration, are two hard-working features that make your job easier. Not only is it easy to operate, you'll get years of time-saving aeration with low maintenance.

The new trailer aerator is available in a variety of tine styles for tees, greens, fairways, lawns, right-of-ways, or institutional grounds.

So if you're responsible for strong, healthy, attractive grass and concerned with saving time and money, it's worth your time to investigate Dedoes quality aerators. Call today for a demonstration.

Let us help you select the best Dedoes Aerator for your application. Call us direct at 800-521-7086

DEDOES INDUSTRIES
INCORPORATED

1060 W. West Maple Rd., P.O. Box 575 OUT OF STATE
Walled Lake, MI 48088 313-624-7710 800-521-7086

Circle No. 107 on Reader Inquiry Card

SPREAD IT

OR

SPRAY IT

... but do it right with LESCO



DEFLECTOR



HOPPER COVER



The LESCO Spreader features corrosion-resistant "Delrin" 100 ST gears; sturdy stainless steel axle, impeller shaft and on/off assembly; adjustable metering slide; pneumatic tires and a bumper handle to protect the impeller. Also available in an electric truckster-mount model.

Optional Accessories:

Jet-Action Deflector—allows straight spreading edge along sidewalks, driveways and ornamental plantings.

Hopper Cover—protects product and allows spreader to be transported without being emptied.



The LESCO Sprayer fits a standard or small-size pickup truck or in a van with either a side or rear door. Available in both gasoline and electric models, the sprayer features an in-line strainer to remove debris and prolong pump life; an electric-return hose reel to eliminate hand-cranking and a low-volume spray wand that is ideal for applying broadleaf herbicides. The gasoline model features a five-horsepower, four-cycle engine. The electric model has a 12 VDC motor-driven pump and a rechargeable battery pack. Optional accessories available for the gasoline model—ask your LESCO sales representative for details.

... And remember it's LESCOSAN* liquid or granular for most efficient crabgrass control.

*Lescosan is Betasan, a registered TM of Stauffer Chemical Company.

Put the LESCO Spreader and Sprayer to work for you.

Call Barb to order:

(800) 321-5325

Nationwide


(800) 362-7413

In Ohio

LESCO


Lesco, Inc.
20005 Lake Road
Rocky River, Ohio 44116
(216) 333-9250

Circle No. 120 on Reader Inquiry Card



We sell the patented ChemLawn Gun.





More proof that the best costs less on tees and greens

It's as true this year as it was last.

Check the chart of comparative fungicide costs and you'll see for yourself why Daconil 2787 Flowable Fungicide is more economical to use than the other leading products.

Using a typical tee and green spray schedule, Daconil 2787 gives you greater savings on a per 1000 sq. ft. basis when it comes to delivering superior control of your most serious diseases — dollar spot (including benomyl-resistant dollar spot), Helminthosporium (leafspot and melting-out) and large brown patch.

In fact, Daconil 2787 has a proven record of providing the most effective control of 10 major turf diseases.

And here are more reasons why it pays to use Daconil 2787. There's no need to add a costly spreader/

sticker. Daconil 2787 already has it built in to assure you of full and even coverage for maximum disease protection.


Daconil 2787 resists wash-off. Which means it keeps on working during heavy rains or watering.

What's more, in over 15 years of continuous use, there has never been a documented case of resistance with Daconil 2787. Even on courses where it was applied at weekly intervals over many years.

The facts are there. The proof is plain to see. Daconil 2787 gives you both effective and economical disease control.

So this season, make it Daconil 2787 on your tees and greens. Because the best costs less.

Daconil 2787 from SDS Biotech.



* Costs based on manufacturer suggested retail unit price as of January, 1984.

Daconil 2787® Flowable Fungicide



Always follow label directions carefully when using turf chemicals.



Agricultural Chemicals Business
SDS Biotech Corporation
7528 Auburn Road, P.O. Box 348
Painesville, Ohio 44077

TURF

tively or preventively.

The fourth fungicide, triadimefon (Bayleton), does not have to be drenched in to be effective. However, it does have to be used preventively. This means it has to be applied before the disease becomes active during the current season. This does not mean Bayleton cannot be used on turf areas that had the "Fusarium blight syndrome" the year before, only that it must be used before the disease becomes active this season.

Since more than one fungus appears to be involved in this syndrome, future research may indicate a difference in the effectiveness of these fungicides on the various fungi causing "Fusarium blight syndrome".

Yellow patch

Yellow patch is caused by *Rhizoctonia cereales* and is a newly recognized disease of Kentucky bluegrass. This is a cool weather disease that occurs in September through November, depending on your location.

The initial symptoms are red to purple leaves on the infected plants. The disease is characterized by "frog-eyes" which resemble the "Fusarium blight syndrome". Consequently, the disease is often mis-identified as Fusarium blight. The main differences between the two diseases are "Fusarium blight syndrome" occurs in warm weather and is characterized by wilting turf in the active spots, whereas yellow patch occurs in the cool weather of the fall and is characterized by the red blades on the infected grass plants.

Cultural management Nitrogen fertility during the growing season is necessary for recovery of the older yellow patch "frog-eyes" that were formed in previous seasons. The effect nitrogen has on development of new yellow patch "frog-eyes" has not been determined.

There are some products which claim to change the chemical and

biological activity of soil and thatch to make it a better environment for biological activity of beneficial microorganisms and healthier plant growth. Several products were tested for their management of *Rhizoctonia* yellow patch and some promoted excellent recovery of older yellow patch "frog-eyes" and prevented new ones from forming. They were Lawn Keeper and Green Majic. It is important to point out these are only preliminary findings and further research is needed to check the repeatability of these results and to determine rates, timing and the exact effect the products are having on disease reduction.

Chemical management Preliminary data suggest that iprodione (Chipco 26019) and fenarimol (Rubigan) will manage *Rhizoctonia* yellow patch. Effective timing and minimal rates have to be determined. It appears nitrogen application will have to be made in conjunction with the fungicide. Otherwise, fungicides may prevent the older "frog-eyes" from becoming active again and new ones from forming. Also, the older "frog-eyes" will not fill in and the maximum benefit from the fungicide treatments will not be realized.

Melting-out

This disease is often incorrectly referred to as leaf spot. To be correct, melting-out is caused by *Drechslera poae* (formerly *Helminthosporium vagans*) and is a disease of Kentucky bluegrass that occurs in the cool, wet weather of spring. The disease starts out as spots on the leaf blades and in a 2-3 week period, rapidly moves down the leaf sheath and into the crowns and roots. The entire grass plant is often killed or severely damaged during this period, which is where the term melting-out arises. The entire stand of Kentucky bluegrass seems to melt away.

Leaf spot, on the other hand, is

a warm weather disease of many grass species caused by the fungus *Drechslera sorokinianum* (formerly *Helminthosporium sativum* and sometimes currently referred to as *Bipolaris sorokinianum*). Are you thoroughly confused now? Don't feel bad, you're not alone.

There are many Kentucky bluegrass cultivars that are resistant to melting-out. The first resistant cultivar was 'Merion' which had excellent resistance to melting-out. Some of the newer Kentucky bluegrass cultivars, i.e. Parade, Baron, Cheri, Majestic, etc., have some resistance to melting-out, although it is not the same excellent resistance 'Merion' had. Consequently, stands of some of the newer Kentucky bluegrass cultivars may be thinned by melting-out in the spring, allowing for invasion by crabgrass, quackgrass, tall fescue and/or broadleaf weeds. This means cultural, biological and chemical management practices to reduce the severity of melting-out will have to be incorporated into your turf management programs.

Helminthosporium melting-out is one of the oldest, most written about, turfgrass diseases. Unfortunately, much of what has been written about the disease, is based on "folk law" and not good scientific data.

First, much of the older literature refers to melting-out having a "leaf spot stage" in the spring, during the cool, wet weather and a "melting-out stage" during the hot weather of the summer. Anyone who is familiar with the disease knows all the damage is done during the cool, wet weather of spring. With the arrival of warm weather the turf begins to recover.

Secondly, practically all the literature says to avoid spring nitrogen, because it will increase the severity of *Helminthosporium* melting-out. It appears the research that led to the erroneous conclusion was based on greenhouse data and not field. At Michigan State University, data from

Give your Elm Trees a new lease on life.

The beautiful elm tree, which is the most important landscape tree in a large part of the industrialized world, has been attacked by a wide array of diseases. Of these, Dutch elm disease (Ded) is probably the most commonly known and also the most devastating.

Now a new fungicide, Phyton-27™ is being offered for Dutch elm disease (Ded) prevention and control. Phyton-27 (pronounced "fight on") is an exciting new water solubilized copper-based chemical system that can be safely injected into a tree.

Field results are in!

Phyton-27 is the *only* product label-recommended for use in trees with up to 30% Ded, and in most cases provides two to three years Ded prevention.

Phyton-27 means good business.

This is the kind of product that homeowners, municipalities, and groundsman have been waiting for...an easy to use economical treatment that can be used as a preventative, or, when combined with good pruning practices, as a therapeutic treatment for trees with Ded ratings up to 30%. Phyton-27 can be used effectively from mid-spring when the leaves are fully developed through early fall.

Your customers will want this product.

The reasonable cost of Phyton-27 and the opportunity for you to offer continuing arbor service to your clients can help you build your business.

We want you to try Phyton-27

Phyton-27 is compatible with most tree injection equipment, and may be used with either trunk or root flare injection techniques. Low cost, durable injection equipment and accessories along with disease identification guides, instructional literature and promotional aids to help you use this product are all available from the makers of Phyton-27.

Call or write for complete information, the tree care season is coming. Don't miss it.



- Please send complete information and prices
 Call me at () _____

Name _____

Title _____

Co./Agency _____

Address _____

City _____

State _____

Zip _____

SOURCE TECHNOLOGY BIOLOGICALS, INC.
2850 Metro Drive, Suite 800, Minneapolis, MN 55420
Phone (612) 854-9426 • TOLL-FREE: 1-800-ELM-TREE

PhytonTM
27

Circle No. 136 on Reader Inquiry Card

TURF



ESTECH SECRET NO. 8

“ The hot
Japanese import.

When Japan's largest fertilizer supplier, Zen-Noh, needed phosphate rock to make fertilizer, they came to ESTECH. Through our joint venture at the Watson Mine in Ft. Meade, Florida, we helped Zen-Noh import over 600,000 tons of phosphate rock into Japan last year.

Which in turn grew a lot of grain.

Which in turn fed a lot of people.

You'd think we would have shared this claim to fame before now. But that's the secret of our success...

At ESTECH,
we don't talk.
We just do.”

Estech, Inc.

We'll share some equally revealing secrets
about Professional & Commercial Fertilizers.
Call Irv Stacy, National Sales Manager, Par Ex 813/294-7793

the last four years suggests just the opposite. Nitrogen in the spring actually reduces the severity of *Helminthosporium* melting-out. We recommend two nitrogen applications during the spring period to help manage *Helminthosporium* melting-out. Each application should be between 1/2-1 lb. of actual nitrogen/1000 sq. ft.

The third management practice is biological in nature. It consists of daily irrigation to keep the mat or thatch moist, to encourage the build up antagonistic microorganisms that prevent the fungus *D. poae* from sporulating, or germinating, or infecting. There is also a possibility that these antagonistic microorganisms may even destroy *D. poae*. While the details have not been worked out, the results have shown a dramatic reduction in the amount of *Helminthosporium* melting-out, where light daily irrigation has been applied.

The actual concept may be hard to grasp since the disease occurs under cool, wet weather conditions, but apparently just a few days without rain, allows the top of the thatch to become dry and allows the *D. Poae* fungus to grow and infect these grass plants. You aren't irrigating the turf, you are irrigating the thatch to keep it moist.

Following good cultural and biological practices will help improve the disease management obtained with the fungicides. For the people in the lawn care business, there are now three excellent fungicides which will manage *Helminthosporium* melting-out during the 3-4 weeks it is normally a problem in the spring; iprodione(Chipco 26019), vinclozolin(Vorlan), and chlorothalonil(Daconil). There is a possibility that anilazine(Dyrene) may also manage the disease for the desired period of time, although more research is needed. Remember, applying these fungicides with a little nitrogen will make them more effective.

WT&T

Spread our wings and get the job done faster!

If you're responsible for maintaining a large spread or highway right-of-ways, Woods Batwing® rotary cutters can help you cut your big jobs down to size.

And when Batwing® goes to bat for you, it does more than mow. These heavy-duty cutters will also tackle the toughest weeds, clear light brush and shred stalks as well.

Five Batwing® models are available for cutting from a 10' to 20' swath, and each is engineered from the ground up to take the punishment of continuous commercial mowing.

Find out more about the safe, convenient and dependable design features of Woods Batwing® rotary cutters. Send for your *free* catalog today.



Division of Hesston Corporation

Oregon, Illinois 61061

Woods mowers are backed by 36 years of dependable quality

Circle No. 148 on Reader Inquiry Card




PREVENTIVE TREATMENTS WITH BAYLETON CAN KEEP YOUR

Fungi. Never have so many been fought off for so long by one product. ®BAYLETON Turf fungicide, now in water dispersible granular form for less dust and easier measuring. It works.

Preventive or curative applications of BAYLETON give grass the guts to resist dollar spot, copper spot, powdery mildew, red thread, and rusts. Preventive applications of BAYLETON will effectively control

anthracnose/*Poa annua* decline, stripe smut, Fusarium blight, Fusarium patch, plus pink and gray snow mold. Control lasts up to 30 days or longer, depending on environmental conditions. Use it on greens by

BAYLETON is a Reg. TM of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen.



HEY, WHY DON'T
WE TURN BROWN
LIKE WE DID LAST
SUMMER?

THE
SHADOW
KNOWS.

TURF'S GREEN BEAUTY FROM TURNING INTO BROWN UGLY.

all means—and fairways, too. With its lasting residual action, it provides the most cost effective control for summer diseases.

BAYLETON works so well for so long because it gives you systemic activity through

both foliar and root uptake. Use recommended rates and follow label directions.

BAYLETON Turf fungicide. It can prevent your turf's green beauty from turning into brown ugly. See your turf chemicals distributor.



Mobay Chemical Corporation
Specialty Products Group
Box 4913, Kansas City, MO 64120

TURF DISEASE GUIDE

The southeastern United States is blessed with a wide range of choices when it comes to turfgrass varieties.

The predominant turf species used in this region is bermudagrass; however, there are five other warm season grasses used extensively for turf purposes. These include St. Augustine, zoysiagrass, centipedegrass, carpetgrass and bahiagrass.

During the winter months, when warm season grasses are brown and dormant, various annual and perennial turfgrasses are used as temporary overseeded grasses.

Although most of the serious

diseases of southern turf are caused by fungi, other agents such as bacteria, viruses, and nematodes can cause serious problems to certain grasses.

Turf producers in the south cannot depend solely on fungicides for disease control. Good variety selection, cultural and fertility practices are very important in disease control. No amount of fungicide will compensate for poor fertility and cultural practices.

Brown patch

Brown patch is the most common turf disease occurring in the southeastern United States. Al-

though St. Augustine and zoysiagrass are the most susceptible cultivars, even the more tolerant centipede, bermudagrass and ryegrass are frequently damaged by this fungus.

Brown patch is favored by warm, moist weather when nighttime temperatures are relatively cool. Therefore, in certain areas of the south, brown patch can and does occur any month of the year.

In the upper regions of the south the most favorable conditions for disease development usually occur from late April through mid-October.

Symptoms of brown patch on warm season grasses are some-

Southern Turf Diseases

by Don Blasingame, extension plant pathologist, Mississippi State University



what different from the symptoms that are described for cool season grasses. Even though the grass is usually killed in a circular pattern, many times the characteristic smoke ring is not seen on southern turf. Also, under certain environmental conditions the fungus may cause a gradual thinning of the turf over a rather large area instead of killing in a circular pattern.

There are several factors that tend to make the grass more susceptible to brown patch. One of these is the excessive application of nitrogen fertilizer. This promotes a lush growth of grass that is readily attacked. Another condition that leads to severe disease development is watering late in the afternoon and allowing the grass to remain wet for long periods of time. The excessive accumulation of thatch creates a most favorable environment for development of brown patch and many other diseases that are caused by fungi.

Fungicides are best used on a preventive schedule. Once symptoms develop control can be difficult.

Dollar spot

Dollar spot is a fungus disease common in the southeast on many species of grasses, particularly on bentgrass, bermudagrass, zoysiagrass, and annual and perennial bluegrasses.

Dollar spot is a disease in which symptoms are different on certain warm season grasses than those noted on cool season grasses.

On the finer textured grasses, such as bermudagrass and zoysiagrass, the grasses are killed in small patches two to three inches in diameter. Under severe conditions these patches may coalesce so that the turf has a mottled appearance. Blades of grass at

Spring dead spot attacks

bermudagrass and zoysiagrass while dormant, becoming evident during spring greenup.



Brown patch is the most common turf disease in the Southeast. It affects primarily St. Augustine and zoysiagrass in the South and bentgrass and perennial ryegrasses in the North. It is prevalent during warm, moist days with cool nights.

the outer edges of the infected area develop tan spots with reddish-brown margins.

On the coarser warm-season grasses the turf is killed in larger patches that may range up to a foot in diameter.

Dollar spot is prevalent during periods of mild weather during the spring and fall months. Unlike brown patch, dollar spot is retarded by high levels of nitrogen fertilizer.

Because excess nitrogen tends to favor the development of brown patch and other diseases, discretion must be used in applying nitrogen. Watering should be performed only in the early morning so the foliage can dry quickly. Fungicides can be used to help bring the diseases under control once it gets established.

Leaf spots

There are a number of fungi that cause leaf spots on many of the southern grasses. Regardless of the causal agent, these leaf spots on southern grasses are similar

and so are the control measures.

Helminthosporium leaf spots (Melting Out) Bermudagrass and ryegrass are most severely affected by helminthosporium infections, although the fungus can survive on centipedegrass and St. Augustine.

Infection can occur over a wide range of temperature but usually is more severe when temperatures are 70° to 95°. Milder temperatures in the spring and fall are more favorable for infection.

Helminthosporium causes small, dark colored spots or flecks on the leaves and sheaths. Leaf spots are usually more numerous near the collar of the leaf blades. Severely affected leaves wither and die and the turf frequently becomes brown and thin.

Symptoms on overseeded ryegrass are altogether different. Although leaf spots may occur, this same helminthosporium can cause severe crown rot. This causes a yellowing and discoloration of the grass and a general

TURF

thinning of the turf.

Fertilize with adequate levels of nitrogen and potassium if helminthosporium diseases become a problem. With careful management, apply fungicides recommended for helminthosporium blight control.

Gray leaf spots St. Augustine is the primary host for gray leaf spot. The disease occurs throughout the lower south during warm, humid weather.

Spots on the leaf blades are the most visible symptom but sheath and stem lesions also occur. Leaf spots begin as olive green to brown, water-soaked spots as small as a pin head. These enlarge rapidly and form a circular to elongate lesion that are brown to ash colored with purple margins.

The disease occurs during moderate to warm weather accompanied by high relative hu-

midity. Severity of the disease is enhanced by applications of nitrogen fertilizer and is more of a problem in shaded areas where the grass remains wet from dew.

Treatment with a fungicide may become necessary if the disease outbreak is severe and accompanied by prolonged periods of wet favorable weather.

Rust

Puccinia species infect a number of grasses grown in the south, including ryegrass, zoysiagrass, bluegrass, fescue, bermudagrass and St. Augustine. Zoysiagrass and bluegrass are the most often affected grasses.

Susceptibility varies with the variety. Fungus infection is favored by minimum and maximum temperatures of 50° to 70° F. respectively. For this reason, the disease does not usually cause

severe damage over an extended period. It is likely to be more severe in shaded areas during rainy, humid weather. Affected turf will appear unthrifty and begin to thin.

The disease is characterized by the presence of pustules on the leaf blades. These pustules range from bright orange to cinnamon-brown in color depending upon the species of fungus present.

Certain varieties of ryegrass are extremely susceptible to rust and sometimes severe damage can occur. On warm season grasses, zoysiagrass, especially Meyer and Emerald varieties, are most severely affected by rust.

Fertilize to stimulate grass growth, mow on a four to five day schedule and catch clippings. If necessary, a fungicide may be applied to help reduce the amount of disease present.

SOLO'S JET SET

THE JET SET IS "IN" —
WE WANT YOU IN TOO!


Right now, people all over the country are asking for SOLO quality sprayers, because they are:

- I**Nvaluable for agriculture, horticulture, pest control, sanitary and industrial application, or just around the house and campsite.
- I**Ndifferent to corrosion - made of high-impact plastic with VITON seals.
- I**NDispensable for applying many cleaning solutions, all approved pest control and agri-chemical formulas, including herbicides.
- I**Ncomparable in capabilities and service.

SOLO products, supported by strong marketing programs and co-operative advertising support will boost your sales. Contact us today for detailed information.



HANJET 455




JETPAK 425

SOLO

SOLO INC., 5100 Chestnut Ave., P.O. Box 5030, Newport News, Va. 23605-0030
(804) 245-4228 Telex 823-664

Circle No. 135 on Reader Inquiry Card

HOW TO SELL LAWN DISEASE CONTROL, AND DELIVER IT.



HELMINTHOSPORIUM LEAF SPOT



SCLEROTINIA DOLLAR SPOT



RHIZOCTONIA BROWN PATCH



FUSARIUM PATCH

Turn disease problems into profits with CHIPCO® 26019 fungicide.

Selling disease control is a great way to boost profits and attract new customers.

And CHIPCO® 26019 is the ideal product to help you do both.

The selling proposition: a beautiful, disease-free lawn.

Everything you do for a customer is designed to create a beautiful lawn. Turf diseases look bad...and that can make *you* look bad. A programmed approach includes an effective, long-lasting fungicide like CHIPCO 26019 keeps your customers satisfied.

CHIPCO 26019 is the only fungicide you need.

It can be used safely on all lawn grasses nationwide to control the major turf diseases like leaf spot, dollar spot, fusarium and brown patch. Highly effective and long-lasting, CHIPCO 26019 fungicide fits into your treatment schedule and reduces callbacks.

In short, when you've got CHIPCO 26019 on the shelf, you're ready for *business*.


Send for FREE booklet on how to sell disease control.

We've put together a booklet that details lots of effective techniques and tools for selling disease control to your customers...and delivering it with effective, long-lasting CHIPCO 26019 fungicide.

Turf Fungicide
CHIPCO®
26019

TAKING CARE OF BUSINESS

Send to: Rhône-Poulenc Inc.
CHIPCO Fungicide Lawn Care Center
P.O. Box 125 Black Horse Lane
Monmouth Junction, NJ 08852



Please send FREE *How to Sell Lawn Disease Control* booklet to:

NAME _____

TITLE _____

COMPANY _____

ADDRESS _____

CITY _____ STATE _____ ZIP _____

TURF

Spring dead spot

Spring dead spot is a serious disease of bermudagrass in certain parts of the upper Sunbelt. Generally speaking, it is found on bermudagrass or zoysiagrass under high maintenance.

Damage to the turf apparently occurs during the dormant season, and when greenup occurs in the spring, there are areas a few inches to several feet in diameter where the sod is completely dead.

The causal agent for spring dead spot has not been identified. The only control procedures recommended at the present time are good cultural practices and limiting the use of nitrogen fertilizer especially late in the growing season.

Research has shown that fungicides can limit the damage. However, at the present time only

two fungicides are labeled and these may be limited to use in certain states.

St. Augustine decline

St. Augustine decline (SAD) is caused by a virus. It causes a mosaic-type of chlorosis of the leaf blades that may resemble a nutrient deficiency or mite feeding. Evidently there are several strains of the virus since there is a great range in damage to St. Augustine.

To this point, the disease has only been recorded in Arkansas, Texas, Louisiana and Mississippi. There are no chemicals available for the control of this disease.

There are several varieties of St. Augustine that are resistant to the virus and can be used in areas where the disease is a potential problem. Floratam was the first variety released that has resis-

tance to SAD. It is also resistant to chinch bugs. It has poor cold tolerance and should be used only in the lower south. Seville is resistant to SAD and is more shade tolerant than common St. Augustine. Raleigh is resistant to SAD and has good winter hardiness.

Downy mildew of St. Augustine

Downy mildew of St. Augustine was first described on common St. Augustine in Texas in 1969. Since then the disease has spread and has been identified in Arkansas, Louisiana and Mississippi.

Downy mildew appears as white, raised, linear streaks that develop parallel to the mid-veins of the leaf. Streaks appear in the spring and remain throughout the summer, giving the leaves a yellow appearance with some death toward the tips. Severe dis-



Systemic Seed Treatment Fungicide controls *Pythium* up to 21 days.

Apron® and Subdue® are Reg. T.M.'s of CIBA-GEIGY

Apron®, a systemic seed treatment fungicide from Ciba-Geigy, works on contact to control *Pythium* on the outer surface of turfgrass seed.

As the seed starts to germinate, Apron is systemically translocated throughout the entire growing seedling. This assures you of maximum emergence and seedling establishment for up to 21 days after planting.

Because Apron is a systemic fungicide, you get better and longer *Pythium* control. And surprisingly, it costs no more than the ordinary seed treatment being used today.


After seedlings are established, you should follow up with fungicide treatments of Subdue® to maintain *Pythium* control in turf.

Call your seed distributor and ask for Apron treated turfgrass seed for this year's overseeding.

Gustafson

17400 Dallas North Parkway
Dallas, Texas 75252
(214) 931-8899

Choosing a turfgrass seed




Marvelgreen winter overseeding mixture used at Hilton Head, South Carolina. *(Upper)*

Sodding New England Patriots' practice field with Baron Kentucky bluegrass. *(Lower)*

Home lawn in New Jersey overseeded with Baron and RAM I Kentucky bluegrasses. *(Upper)*

Kansas City Chiefs' practice field, overseeded with Baron Kentucky bluegrass. *(Lower)*

Here's what some



Arnold Palmer views turf from two different angles: as a golfer and as a golf course owner.

As owner of the Latrobe Country Club in Pennsylvania, Arnold Palmer works very closely with the superintendent, his brother, Jerry. "The new Palmer turf-type perennial ryegrass," says Arnold, "has done a good job here. We've used Palmer when reseeding tees, and on heavily-trafficked areas in the fairways and roughs.

"At Florida's Bay Hill Club, Superintendent Jim Ellison and I work closely on our overseeding program. We've been using new turf-type perennial ryegrasses, like Palmer, there too. We've got that course in the best shape ever now. And that's been evident by the comments we get during the Bay Hill Classic.

"When I work with Ed Seay, head architect for my Palmer Course Design Company, we are faced with a whole set of grassing problems for each location. I can't be involved on a day-to-day basis with all my business ventures. But I do know this: some of the new turf varieties released by Lofts have enabled us to improve courses like never before."

There are many decisions to be made when the Palmer Course Design Company redesigns or builds a course. Choosing the best grass for the existing conditions is a major one.



Ed Seay, Golf Course Architect, Palmer Course Design Company, Jacksonville Beach, Florida, describes the grasses he recommends for golf courses around the world.

From Montana to Japan, Ed Seay's design expertise is in demand throughout the world. And the seed is important to him too, as a finishing touch to the quality of each course.

"We've developed some standard mixtures that have worked very well for our


clients in many different areas. For example, where cool-season grasses are well-adapted, and bluegrasses can be used on roughs, we recommend a mixture of 65% Kentucky bluegrass, either Baron, Nassau, RAM I or Georgetown, 10% Jamestown chewings fescue, 10% Reliant hard fescue and 15% Palmer perennial ryegrass. This blend is very adaptable, performing well in sun or shade.

"Where bluegrass is used on fairways and tees, we suggest 85% Kentucky bluegrass plus 15% Palmer perennial ryegrass. On fairways and tees we add Mystic to the bluegrass portions because it's very aggressive. That's good for recovery from divot scars and heavy traffic. And from experience we know Baron, Nassau, RAM I and Georgetown hold up under lots of traffic.

"With all the details of designing a new course, or renovating an existing course, it just makes good sense to follow up with a top-quality seeding program."



of the experts are doing for



Jim Ellison, Superintendent of Bay Hill Club, Orlando, Florida, discusses his winter overseeding program for the Bay Hill Classic.

"As host to the world's top golfers, I need turf with excellent playing qualities. And because we're on nationwide TV, my course has to look great too.

"Although the tournament doesn't occur until early March, we start getting ready in the fall with our winter overseeding program. We overseed the bermudagrass on the greens, tees, tee and green banks, and all the roughs. This adds up to about 100 acres.

"The first week of November we clip the bermudagrass real close. In the roughs we brush it with a street-sweeper

to make it stand straight up. "Next we overseed. For the roughs, tee banks and green banks we overseed with Palmer perennial ryegrass at the rate of 200 lbs./acre.

"On putting greens we use a mixture of 60% Palmer turf-type perennial ryegrass, 25% Jamestown chewings fescue, and 15% Sabre *Poa trivialis*. Our seeding rate is 35 lbs./1,000 sq. ft.

"We decided not to overseed the fairways because during winter the cooler weather slows the bermudagrass growth rate, making an excellent playing surface. The light green bermudagrass, contrasted against the dark

green ryegrass roughs, gives the course a nice definition.


"The tricky part of all this preparation is that we have to do it at the height of the tourist season. While we're working, we've got golfers playing more than 300 rounds a day. But, with all that going on, we still get great results! Our turf ends up with the qualities the pros look for — like fast, smooth putting greens and rich color.

"Arnold Palmer, who's owner of this course, is very pleased with our program. He likes the way the course looks; and he is especially pleased with the putting quality of the greens."



At Bay Hill, the overseeding program plays a significant part in the way the course looks and how it plays. Jim Ellison, Superintendent, feels they're equally important, so he chooses his grass seed carefully.

... golf courses ... home lawns ... sod production




Mike McGehee, Ironwood Country Club, Palm Springs, CA, tells how he ended up with a great-looking course despite poor growing conditions.

Superintendent Mike McGehee was faced with terrible soil conditions for growing grass. "The course is built on an alluvial fan. Surrounded by mountains, weather alternates between 120° and 30°. The soil is extremely rocky and alkaline, and it's very hard to grow any kind of grass. The grass tends to be chlorotic so we have to constantly supply it with fertilizer and iron to keep it green. We can't even prepare a seedbed properly because the ground is too rocky to renovate.

"To provide a playing surface for the winter tourist season, we have to overseed bermuda-

grass with ryegrass. We used annual ryegrass for years. But, the playing conditions were not up to the heavy demands of our busy season. So we tried Palmer perennial ryegrass, and it proved to be the best overseeding program we had ever had. For the first time we had beautiful green grass. Our members were raving over the improved conditions.

"I can't imagine any course having worse soil conditions than Ironwood. I really believe that if Palmer perennial ryegrass can look this good here, it ought to do even better in other areas."



At Ironwood Country Club in Palm Springs, California, it's no easy task to keep the course in a favorable condition. Palmer perennial ryegrass has made the job easier.



Joe DeSantis, Royal Lawns, Pinebrook, N.J., talks about programs his lawn service company has found successful.

"My customers are very impatient. They want a beautiful lawn, and they want it fast! Here in the Northeast that gives me only a limited time to turn a lawn around. And often lawns are so poor they have to be completely reseeded. We've used almost one million pounds of seed in the 12 years we've been in business.

"The most critical time for lawn care is in the fall. And it's the best time to seed. In our fall seeding program, we use a blend of 70% Kentucky bluegrass, 20% Palmer perennial ryegrass, and 10% Jamestown chewings fescue. For the bluegrass, we use

Baron or RAM I, depending on conditions. Where there's shade, we use RAM I. It resists powdery mildew, which is a common problem in shady areas. In areas that are not primarily shaded, we use Baron; it looks good in all kinds of conditions.

"For fast germination, we have especially good results with Lofts Triplex Ryegrass Mixture. It's a blend of Prelude, Palmer and Yorktown II turf-type perennial ryegrasses.

"Many people only consider ryegrass a filler; but that's a misconception. We've found

these new, turf-type perennial ryegrasses to be as attractive and persistent as bluegrass; in some cases they perform even better. They blend well with bluegrasses. They can tolerate the incredible abuse a lawn can get from a homeowner's family and pets.

"We have a carefully planned, professional lawn maintenance program that takes many factors into consideration. We feel that if we start with top-quality seed, educate our customers on watering and mowing, and then keep our chemical programs going, we're almost guaranteed success."



The lawn care business can be tough. The proper use of chemicals is, of course, very important. But reseeding is very often required and the choice of seed makes all the difference in the world. Here again, the newer perennial turf-type ryegrasses are proving very successful.



Jack Kidwell, Richmond, Virginia, relates how the new turf-type tall fescues have helped his sod business.

With over 1,000 acres, Jack Kidwell is Virginia's largest sod producer. And success doesn't come easily in the Washington area and southern Virginia, where the worst turf problems are created by the long, hot summers and *Fusarium roseum*, a disease that's tough on bluegrass.

"For many years, we relied primarily on Kentucky bluegrasses, KY-31 tall fescue and bermudagrass. But the new turf-type tall fescues have really changed things for us. So much so, that now tall fescue represents more than half our annual production of cool-season grasses.

"One particular turf-type tall fescue that's worked very well for us is Rebel. We've found it far superior to KY-31, with respect to texture, persistence and all-around performance. And it forms such a

strong sod that we don't have to use netting other than for early harvest conditions.

"For us, Rebel has pleased everyone right down the line. I like the way it grows and harvests, my customers like its improved performance and their customers get the benefits of good looks, dense growth and easy maintenance.

"As far as Kentucky bluegrass, Virginia Polytechnic Institute and the University of Maryland have endorsed this new Georgetown. I've been using it and it's doing very well in our hot, humid climate. In fact, they've been testing it specifically for hot weather performance and it's doing exceptionally well."



The introduction of the new turf-type tall fescues, like Rebel, has opened new markets for the sod business. The fine, dense growth and all-around performance keep customers and growers happy.

Lofts Proprietary Turfgrass Varieties:

Kentucky Bluegrasses	Perennial Ryegrasses	Fine Fescues	Meadow Fescue
Baron	Palmer	Jamestown Chewings	Beaumont
Nassau	Prelude	Reliant Hard	
RAM I	Yorktown II	Intermediate Ryegrass	Tall Fescues
Georgetown	Cowboy	Agree	Rebel
Mystic	Repell		Clemfline
	Diplomat		

Lofts proprietary varieties are used in their Marvelgreen winter overseeding mixtures.



Lofts Inc.

Bound Brook, New Jersey 08805
(201) 356-8700 • (800) 526-3890

Lofts/Maryland
Beltsville, MD 20705
(301) 937-9292

Lofts/New York
Cambridge, NY 12816
(518) 677-8808

Lofts/New England
Arlington, MA 02174
(617) 648-7550

Great Western Seed Co., Inc.
Albany, OR 97321
(503) 926-5892

Sunbelt Seeds Inc.
Norcross, GA 30094
(404) 448-9932

ease occurs in grass that is grown in flood plains or poorly drained areas.

The white-streak symptom in early stages is easily confused with the virus disease, St. Augustine decline. However, the virus symptoms are more yellow in color and more mottled than striped.

Downy mildew has been difficult to control with most common turf fungicides. The cultural practices recommended for control are to maintain good drainage so that no free water stands on areas where St. Augustine is grown.

Fairy rings

Fairy rings generally appear in lawns and other turf areas as circles or arcs of dark-green, fast-growing grass during the spring and early summer. A ring of thin dead grass may develop on one or

both sides of this circle.

The disease is caused by one of several soil-inhabiting fungi that commonly produce mushrooms. Mushrooms that sometimes appear in the ring are the fruiting bodies of these fungi. Stimulation of the grass is due to the release of nutrients from the organic breakdown of the thatch by the growing fungus.

It is difficult to control fairy ring. Two general approaches may be considered. The first is removal. Remove infected grass and soil to a depth of 12 inches or more in a band several feet on each side of the affected area and replace with clean soil.

Another approach is to suppress the disease. For low maintenance grass areas, increase the water and fertilization program to stimulate the declining grass inside the ring.

Symptoms of the ring can be masked by pumping large quantities of water into this area. There are no chemicals labeled for the control of this disease.

Slime molds

Slime molds are a group of organisms which create considerable concern among gardeners and those interested in maintaining a good quality turf. These molds cover above-ground plant parts with a dusty gray-black or dirty yellow mass.

When you look closely at this growth you see small round balls scattered over the plant. If you rub these between your fingers a sooty-like powder emerges. This sooty-like powder is the spores of the fungus.

Slime molds normally live in the soil where they feed on decaying organic matter. When the



Versatile. All-hydraulic. Air cooled (gas or LP) or electric.

Economical Sweeping

The Tennant 240 is a popular, high speed, easy-to-maintain sweeper. All-hydraulic; no belts, chains or transmission. Sweeps a 53-in. path at speeds to 10 mph. Picks up fine sand and bulky debris in 1 pass. Vacuum/filter controls dust.

High-Low dump option. Hopper empties into 5-ft container or into floor-level compactor.

Converts to scrubber in 15 min. using optional attachment. Lease and time sales plans available. For free brochure, write: Tennant Co., 701 No. Lilac Dr., Minneapolis, Minnesota 55422.



TENNANT MAINTENANCE SYSTEMS
sweepers • scrubbers • scarifiers • floor coatings

Circle No. 138 on Reader Inquiry Card

BEAUTY LAWN ZOYSIA SOD

(Meyer Z-52) for your
FAIRWAYS, ATHLETIC FIELDS, PARKS

- Zoysia saves big on water, fertilizer, insecticides, fungicides, mowing.
- Zoysia provides an outstanding playing surface for fairways.
- Stands up to hard usage.
- Truck-trailer shipping throughout U.S.

ROW PLANTING SERVICE

100 Mile Radius of Louisville, Ky.



Jay and John Frick
The Zoysia Men



The Solution to Old Problems

BEAUTY LAWN ZOYSIA

603 Union Road • Lebanon, Ohio 45036
Near Cincinnati, Ohio (513) 424-2052

Our 35th Year

Circle No. 102 on Reader Inquiry Card

JUNE 1984/WEEDS TREES & TURF 43

TURF

slime mold is ready to reproduce, it grows up on to the grass blades so that the spores may be spread great distances. Its only purpose of selecting plant parts above the soil line is to distribute the spores over a further distance than it would be able to from the soil surface.

Slime molds do not feed on living plants. They only use them for support during reproduction.

Slime molds occur during wet weather throughout the spring, summer and fall. They disappear rapidly as soon as it becomes dry and chemical control is usually not necessary.

Pythium blight

Pythium blight can be a devastating disease on overseeded ryegrasses; however, bermudagrass and the other warm-season grasses can be affected to a lesser degree.

An abundance of moisture is required for pythium blight development. In addition, the disease is favored by warm temperatures.

Affected grass is killed rapidly in spots two to four inches in diameter. These spots may develop into streaks so that large areas of turf are damaged.

During early stages of development the affected grass appears wilted and greasy. At times the affected turf spots may have a cottony appearance due to the abundant growth of the fungus. For this reason the disease is frequently referred to as cottony blight.

Certain species of pythium can also cause root rot on turfgrasses. Due to the restricted root function the plants become chlorotic and the turf begins to thin.

On overseeded grasses the disease can be limited by using treated seed and delaying the overseeding until as late as possible during the fall. Water as little as possible during periods of favorable disease activity. The perennial ryegrasses are not as susceptible as annual. Under severe disease pressure chemical control may be required.

WT&T

DIRECTORY

Turf Fungicide Directory

Common Name	Brand Name	Company	Circle No.
anilazine	Dymec 50	Gordon	201
	Dyrene	Mobay	202
	Ortho Dyrene	Ortho/Chevron	203
	Proturf Fung. III	Scotts	204
benomyl	Proturf Fung. DSB	Scotts	205
	Tersan 1991	Du Pont	206
cadmium	Caddy	Cleary	207
	Cadtrete	Cleary	208
	Cadminate	Mallinckrodt	209
	Kromad	Mallinckrodt	210
chloroneb	Proturf Fung. II	Scotts	211
	Teremec SP	Gordons	212
	Terreneb SP	Kincaid	241
chlorothalonil	Daconil 2787	SDS Biotech	213
	Proturf 10IV	Scotts	214
cycloheximide	Acti-dione TGF	Tuco/Upjohn	215
ethazol	Koban	Mallinckrodt	216
	Terrazole	Olin	217
fenarimol	Rubigan	Elanco	218
iprodione	Chipco 26019	Rhone Poulenc	219
	Proturf Fung. 6	Scotts	220
mancozeb	Fore	Rohm and Haas	221
	Formec 80	Gordons	222
maneb	Tersan LSR	Du Pont	223
mercuries	Calo-Clor	Mallinckrodt	224
	Calo-Gran	Mallinckrodt	225
metalaxyl	Subdue	Ciba Geigy	226
PCNB	Terraclor 75	Olin	227
PMA(PMAS)	PMA, PMAS	Cleary	228
PMA plus Thiram	Proturf Broad Spectrum Fung.	Scotts	229
propamocarb	Banol	Tuco/Upjohn	230
thiophanate	Cleary's 3336	Cleary	231
thiophanate methy	Fungo 50	Mallinckrodt	232
	Proturf Systemic	Scotts	233
thiram	Chipco Thiram 75	Rhone Poulenc	234
	Spotrete	Cleary	235
thiophanate plus thiram	Bromosan	Cleary	236
thiophanate-methyl plus maneb	Duosan	Mallinckrodt	237
triadimefon	Bayleton	Mobay	238
	Proturf Fung. 7	Scotts	239
vinclozolin	Vorlan	Mallinckrodt	240



When your job depends on stopping gypsy moths and other pests, depend on ORTHENE.

ORTHENE® Tree & Ornamental Spray (common name: **acephate**) is the professional way to kill gypsy moth larvae... plus a wide range of other insects. It kills on contact, then keeps killing with residual action.

What other insects? ORTHENE also works on aphids, bagworms, birch leaf-miner, tent caterpillar, Douglas fir tussock moth larvae, webworms, scales, California oakworm, spring and fall cankerworms, Nantucket pine tip moth larvae and adult root weevil.

See your ORTHO Distributor or send in the coupon for more information — make your job easier, with ORTHENE.

AVOID ACCIDENTS: For safety, read the entire label including precautionary statements. Use all chemicals only as directed. Copyright © 1984 Chevron Chemical Company. All rights reserved.



Mail to:
R. G. Gras,
Chevron Chemical
Company,
575 Market Street,
San Francisco,
CA 94105

Please send me more information on ORTHENE.

Name _____

Company _____

Street Address _____

City/State/Zip _____

ORTHENE®

Circle No. 104 on Reader Inquiry Card

Kentucky Bluegrasses and Their Culture

by C. Reed Funk, professor of turfgrass breeding, and R. E. Engel, professor of turfgrass management, Cook College, Rutgers University, New Brunswick, NJ

Selection of turf seed can make a huge difference in the appearance and maintenance requirement of a turf area. Differences between varieties are significant to the professional. Keep this article and the next two parts of the seed series, as a reference for future seed purchases.



Kentucky bluegrass, *Poa pratensis* L., is the most important lawn grass in the northern half of the United States. It is hardy, aggressive, persistent, attractive and widely adapted.

New lower-growing Kentucky bluegrass varieties have been developed which produce a more attractive, durable, persistent turf under a wide range of environmental conditions. These are making this species more useful to the turfgrass industry.

Origin and adaptation

Kentucky bluegrass is native to the Old World and occurs naturally throughout the temperate regions of Europe and Asia. Early colonists brought the grass to North America in seed mixtures, hay and bedding. It was disseminated rapidly by men, birds and other animals.

Much of the bluegrass found on millions of acres of fertile pastures, roadsides, and other open areas developed without seeding by man. Its ability to colonize is one reason for its widespread occurrence.

Like other cool-season grasses, Kentucky bluegrass grows best during the cool months of spring and fall. Studies at Beltsville, MD, and Kingston, RI, have shown some root growth will occur throughout much of the winter in

unfrozen soil if fertility and soil pH are adequate.

With prolonged summer drought, Kentucky bluegrass may become dormant and turn brown. However, it usually recovers and resumes growth quickly with the return of cooler temperatures and favorable soil moisture. Instances of poor recovery from summer drought are usually associated with insects, thatch, excessive density, insufficient rhizomes, disease, and management practices which include excessive applications of nitrogen fertilizer and/or close mowing.

The development of Kentucky bluegrass varieties with greater tolerance to the long hot summers of the transition zone from Southern New Jersey, Washington, D. C., Cincinnati, Louisville, St. Louis and westward is a real challenge to the turfgrass breeder.

Kentucky bluegrass is best adapted to well-limed, fertile loam soils and cool exposures. In humid regions the soil pH should be corrected to 6.0 to 6.8 for optimum performance although some more acid tolerant cultivars, such as Fylking and Victa, may persist on soils with a pH slightly below 5.0. Under arid soil conditions in dry land areas Kentucky bluegrass thrives on soils having a pH as high as 8.0 if irrigation is provided.

Growth of Kentucky bluegrass is best on well-drained soils. However, it is considerably more tolerant of poor soil drainage than the fine fescues. Helminthosporium leaf spot and crown rot can be especially damaging to susceptible varieties on poorly drained soils.

Kentucky bluegrass is not as well adapted to the extremely sandy coastal plain soils as the fescues, or zoysiagrass unless such soils are properly modified with appropriate additions of organic matter, lime, fertilizer and use of some water. It is also moderately intolerant of excess salt accumulations.

A well-limed, vigorous Kentucky bluegrass sod is noted for increasing organic matter content and improving physical structure of soil. Nevertheless, excessive traffic and poor management may weaken the turf and favor the invasion of species more tolerant of compacted soil conditions such as *Poa annua* and knotweed. Friable soils of good physical condition enhance the ease of establishment of Kentucky bluegrass.

Fertility response

Kentucky bluegrass responds well to generous fall fertilization. Minimal spring and summer fertilization is usually best when



Off-type grass plants in the field are killed by roguing crews (left), so the seed crop purity is maintained.



Seedheads emerge on Kentucky bluegrass (left), and later just prior to cutting and swathing in the field.

summer stress is severe.

Turf should be fertilized primarily to improve color and density or to heal injury. This can be done most effectively from September through late fall. Short days and cool fall temperatures stimulate tiller production and root growth. They also reduce the rate of leaf blade elongation and cause the plant to grow in a more decumbent (spreading) manner.

In contrast, during the long days in May and June rapid leaf elongation of Kentucky bluegrass occurs and plants are upright.

Fertilizer applications in the fall do not increase mowing as much as the same fertilizer rates applied during the spring growing season. Also, sparse turfs typically increase their density more

following fall fertilization. Late fall fertilization of Kentucky bluegrass promotes better winter color and also assures early spring green-up.

Excessive nitrogen, that stimulates Kentucky bluegrass in late spring and summer, prevents it from developing the physiologically-hardened condition that helps it survive heat and drought stress. Lush spring growth from high fertility also intensifies damage from the *Helminthosporium* leaf spot and crown rot disease.

In short, fall fertilization of turf causes less turf injury during stress, requires less mowing, gives adequate color and gives better turf density than spring fertilization.

Adaptation to shade

Kentucky bluegrass normally performs best in full sun and light shade. In warmer areas it may tolerate afternoon shading with good air movement. In fact, the cooling effect of light shade may reduce injury from chinch bugs, *Fusarium* blight, heat and drought. In warmer regions, Kentucky bluegrass normally does not occur in full sun.

Shaded areas with restricted air movement in wet climates result in slow drying of the turf and a hot humid microenvironment which weakens the grass and provides conditions favorable for disease development.

Moderate to heavy shading of Kentucky bluegrass reduces carbohydrate food reserves, restricts

Kentucky Bluegrass Varieties

Adelphi (Adikes, Jacklin) is a moderately low-growing, turf-type bluegrass with a very attractive, dark green color which is maintained throughout the entire growing season. It has shown good resistance to leaf spot, Fusarium blight, most races of stripe smut and rust and has moderate resistance to dollar spot. *Adelphi* is a hybrid between a fairway selection from the Bellevue Country Club and *Belturf*.

America (Pickseed) originated as a single, highly apomictic plant. It was selected from the open pollinated progeny of a highly sexual hybrid. This hybrid was obtained from a progeny of the cross 'Bellevue' x 'Belturf'. *America* is a leafy, low-growing, turf-type bluegrass capable of producing an attractive, compact, fine-textured turf of high density and dark color. *America* has shown good resistance to leaf spot and leaf rust. It has shown less damage from stripe rust than many Kentucky bluegrass varieties.

Arboretum (Mangelsdorf) was selected at the Missouri Botanical Garden from old pastures and lawns in Missouri and neighboring states. It is an erect-growing variety highly susceptible to the Helminthosporium leaf spot and crown rot disease. It is useful for low maintenance turf.

Banff (Pickseed) was selected from a closely-mowed turf in Canada. This moderately low-growing, turf-type variety has medium texture and a bright, medium dark green color. *Banff* has excellent early spring color. It has good resistance to leaf spot and most rusts and above average resistance to dollar spot and stripe smut.

Baron (Lofts) was developed in Holland. It has rather broad leaves, a moderately low-growing, turf-type growth habit and a medium dark green color. *Baron* has shown moderately good resistance to leaf spot and has been widely accepted as a good bluegrass variety in many areas throughout the world. The variety has shown only moderate resistance to leaf rust, stem rust, dollar spot and powdery mildew under New Jersey conditions. *Baron* is moderately slow to become green in the spring. It has a large seed and rather good seedling vigor. *Baron* produces high seed yields.

Birka (Burlingham) was developed in Sweden. This variety has a medium fine texture, a moderately low turf-type growth habit and a moderately dark green color. *Birka* has shown good resistance to leaf spot, stripe smut and powdery mildew in New Jersey tests. It is moderately slow to green-up in the spring. The variety is susceptible to stem rust.

Bonnieblue (Burlingham) is a hybrid between the selection from the Bellevue Country Club and *Penstar*. This moderately low-growing, turf-type variety has good resistance to leaf spot, stripe smut and leaf and rusts. It has a bright, rather dark green color and becomes green early in the spring.

Bristol is a hybrid between a fairway selection from the Bellevue Country Club near Syracuse, NY, and *Anheuser Dwarf*. This variety has a rich, dark green color, wide leaves and rather decumbent growth habit with a moderately slow rate of vertical growth. *Bristol* has good resistance to leaf spot and red thread, and moderately good resistance to stripe smut, dollar spot and most races of powdery mildew.

Challenger is a moderately low-growing, leafy, turf-type variety with medium-fine leaves, medium high density, and a very attractive, bright, dark green color. It has excellent early spring color and the ability to stay green into late fall. *Challenger* has shown good resistance to Helminthosporium leaf spot and melting-out, leaf rust, stem rust, stripe smut, and dollar spot. *Challenger* is a hybrid between NJE P-123, a selection from Lafayette Park in Washington, D. C., and PSU K106, a selection found in northern Kentucky.

Cheri (Jacklin) was developed in Sweden. This variety has medium-broad leaves, a moderately low, turf-type growth habit and a medium dark green color. *Cheri* has shown moderately good resistance to Helminthosporium leaf spot and crown rot disease. It has shown only moderate resistance to leaf rust, stem rust, dollar spot and powdery mildew under New Jersey conditions. *Cheri* is moderately slow to greenup in the spring. *Cheri* has large seed and rather good seedling vigor.

Columbia (Turf-Seed) was selected from an old, non-irrigated, moderately low-maintenance turf near Frederick, MD. This moderately low-growing, turf-type variety has medium texture, good density, and a bright, medium dark green color. *Columbia* has an exceptionally attractive early spring color, the ability to stay green into late fall, and the capability of maintaining good winter color in protected locations. *Columbia* has shown good or moderately good resistance to leaf spot, leaf rust, stem rust, dollar spot, stripe smut and Fusarium blight. Turf produced may have a high proportion of stemmy reproductive tillers in late spring and early summer.

Common Kentucky bluegrass, *South Dakota Certified*, is a source of Kentucky bluegrass harvested from nature
continued on page 50

development of roots, rhizomes and tillers, and causes long thin succulent leaves. Such turf is predisposed to diseases, intolerant of wear and less able to recover from injury. Kentucky bluegrass sod laid in intense shade roots slowly and usually fails in 1 to 3 years with the intense shade and wetness of climates such as New Jersey (or the Northeast).

Where shade occurs, Kentucky bluegrass seed content of a mixture should be reduced, but not omitted. This will give a blending of Kentucky bluegrass between the sun and light shade areas.

There is some variation in the shade tolerance of Kentucky bluegrass varieties. *Merion*, which has been one of the better varieties for general use, is highly susceptible to powdery mildew, a disease which is very damaging to susceptible varieties growing in shade, but of little consequence in full sunlight.

Selection for mildew resistance has been of primary importance in breeding shade tolerant bluegrass varieties. *Warren's A-34*, *Eclipse*, *Bristol*, *Benverde*, *Touchdown*, *Nugget*, *Ram 1*, *Birka* and *Glade* are bluegrass varieties with moderate-to-good mildew resistance. A number of promising experiments also appear to have excellent mildew resistance. However, it must be pointed out that different pathogenic races of powdery mildew develop naturally which would cause some of these selections to become infected.

Shade tolerant Kentucky bluegrasses should also have good resistance to leaf spot and other diseases. Also, they must have the ability to photosynthesize enough food to give tillering, generous rooting, rhizome development and carbohydrate storage.

The ability of some of the fine fescues to tolerate tree root competition and the acid infertile soil conditions frequently associated with shaded locations contributes to their success as a shade tolerant component of a turfgrass mixture.

Disease resistance

The present varieties of Kentucky bluegrass show substantial differences in resistance to common turf diseases. Use of disease resis-



Winner by two: Sun and Shade.

Don't handicap yourself with a sun only grass seed. Put Glade, the sun and shade champion in your mix. Glade's already proven. That takes the risk out of who you bet on.

Glade — from your local wholesale distributor.

Glade
Grass

Kentucky bluegrass

U.S. Plant Patent 3151
License in Canada No. 2133

*Another fine, quality-controlled
product of Jacklin Seed Company.*

ralized or native stands. However, some studies indicated natural stands of bluegrass in South Dakota do not contain as much genetic diversity as found in the famous bluegrass region of Kentucky. Kenblue and Park have visually outperformed South Dakota Certified Kentucky bluegrass in New Jersey tests. Studies by Dr. Glen Wood in Vermont showed that bluegrass from the Kentucky areas produced turf more resistant to weed invasion than bluegrass obtained from South Dakota.

Delta was selected in Canada. It is similar in growth habit and appearance to common Kentucky bluegrass and is also highly susceptible to the Helminthosporium leaf spot and crown rot disease. In earlier years *Delta* generally performed as well as common Kentucky bluegrass in turf tests at Rutgers. However, during the past few seasons the performance of *Delta* has been poor.

Eclipse (Jacklin) is a highly apomictic hybrid selected from the progeny of the cross 64-765-4 x Anhesuser Dwarf. The female parent, 64-765-4, was selected from the progeny of the cross SP-1 x Belturf. *Eclipse* is a low-growing, leafy, turf-type variety capable of producing an attractive, dark green turf of good density, good vigor and medium texture. *Eclipse* has demonstrated good or

moderately good resistance to leaf spot, leaf rust, stem rust, powdery mildew, stripe smut, red thread, and dollar spot. It has performed well in shade trials.

Enmundi (International Seeds) is a leafy, attractive, moderately low growing variety developed in Holland. The variety has shown good resistance to leaf spot, stripe smut and Fusarium blight in New Jersey tests. Low seed yields are limiting the use of *Enmundi*.

Fylking (Jacklin) was developed in Sweden. This turf-type variety has good resistance to the Helminthosporium leaf spot and melting out disease. *Fylking* is more resistant to stripe smut, stem rust, leaf rust and powdery mildew than *Merion*. It is occasionally damaged by dollar spot and Fusarium blight. *Fylking* produces an attractive, dense, moderately low-growing turf of a rather fine texture. It maintains this leafy appearance during seed head setting time in May and June when many other bluegrasses become quite stemmy. An attractive, rich dark, green color is developed in early spring which is maintained into late fall and under moderately adverse. *Fylking* is moderately tolerant of close mowing. However, cutting the grass at a height of 1-1/4 inches will favor vigorous growth and help prevent weed invasion. The

variety has rather fine leaves which tend to lean at higher cutting heights, thus a neater appearance is attained with moderately close mowing.

Geronimo (Jacklin) was developed by Mommersteeg International of Vlijmen, Holland. It is a moderately dark green, turf-type variety, with medium wide leaves, and medium density. It has moderately good disease resistance to the Helminthosporium leaf spot and crown rot disease.

Glade (Jacklin) is a moderately fine-textured, dark green selection obtained from an old lawn in Albany, New York. It has shown excellent resistance to stripe smut, many races of powdery mildew, and leaf and stem rust. It has moderate resistance to leaf spot. *Glade* is an aggressive, turf-type bluegrass with a relatively slow rate of vertical growth. This variety has shown good seedling vigor. It has performed well in blends and mixtures with other Kentucky bluegrass, ryegrass and fine fescue varieties. It has shown some tolerance of moderate shade. *Glade* is moderately slow in spring green up.

Georgetown (Lofts) was selected from an old turf in western Oregon. It is a moderately low-growing, turf-type variety with medium texture and a

continued on page 54

NEW ... HARLEY

Double Roller Landscape POWER RAKE




PULVERIZES, LEVELS & RAKES ROCK, roots and other trash. Double Roller gives deeper penetration and allows moving of several windrows into one for faster pickup. **New scarifier.** Also ask us about Harley Rock Pickers.



(701) 252-9300

Toll Free 1-800-437-9779



TERRA 320

320 tines in a 5 Ft. width

Built for the wide open spaces.

You can put in a lot of holes in a short time—Over 14,000 holes per minute—Approx. 3 acres per hr.

TERRACARE PRODUCTS CO., INC.

P.O. Box 506

Pardeeville, WI 53954

Phone - 608-429-3402

tant varieties along with good management practices is the safest and most economical means of controlling many important turfgrass disease problems.

Resistance to disease is a prime consideration in selecting any bluegrass variety.

Melting out disease The most damaging disease of Kentucky bluegrass in New Jersey is a leaf spot and crown rot caused by the fungus, *Helminthosporium vagans*, commonly called "melting out". This disease appears on leaf blades and sheaths as circular to elongate, purplish to brown spots with straw-colored centers. Some lesions extend the entire width of the leaf, especially on the finer-bladed varieties, causing the portion the leaf blade above the affected area to wither and die.

During severe attacks, especially in late spring and early summer, the fungus causes severe leaf die-back and extensive crown rotting which leads to a melting-out condition. The weak plants and thin turf cover allow invasion by

crabgrass and other unwanted weeds before the bluegrass recovers.

Under New Jersey conditions, *Helminthosporium vagans* produces abundant spores during the cool, cloudy, wet season from October through April. Moderate disease buildup may occur in the fall, persist through the winter, and subsequently intensify into severe damage in May and early June. Disease activity and spore production decrease from May through September. If the turf has not been damaged too severely it will recover significantly at this time, provided growing conditions are favorable.

The severity of the *Helminthosporium* leaf spot and crown rot is greatly influenced by various management practices. Disease injury is more severe with close mowing than high mowing. The shorter cut removes more photosynthetic surface thus limiting carbohydrate development. This weakens the turf, making it more susceptible to damage and

less capable of recovery.

Plants receiving low to moderate levels of nitrogen fertilizer often show greater numbers of leaf spot lesions when examined in March or April. However, during late May and early June when the melting-out or crown rot phase of the disease is most severe, turf receiving high rates of nitrogen fertilizer suffers the greatest damage.

Low light intensity due to either shade or cloudy weather also lowers carbohydrate reserves and increases disease damage.

The best methods of controlling this disease are: the use of resistant varieties such as Merion, Eclipse, Bristol and Touchdown, higher cutting heights, and avoiding excessive nitrogen fertilization during the spring season. Varieties such as Park, Delta, Arboretum, S-21 and Kenblue perform little, if any, better than Common Kentucky bluegrass under New Jersey conditions due to their high susceptibility to *Helminthosporium vagans*.

We Made It Simple!

- **QUALITY** — Built with Swiss clocklike precision.
- **LIGHTWEIGHT** — With the comfort of the user in mind.
- **ACCESSIBLE** — A durable self-lubricating and external pump assembly.
- **SIMPLICITY** — Insures cleanliness and longer wear. No tools necessary to disassemble and repair.
- **PRICING** — Here's one instance where you pay less for the best.



BIRCHMEIER

has the
Back Pack
Sprayer
for you —

AVAILABLE IN
5 GAL. (20 K)
3½ GAL. (15 K)
2½ GAL. (10 K)


Technical data	FLOX 2½ gal.	IRIS 3½ gal.	SENIOR 5 gal.
Weight #	11	11.4	12.1
Tank Capacity	2.6 gals.	3.9 gals.	5.2 gals.
Max. Working Pressure	84 PSI	84 PSI	84 PSI

FOR MORE INFORMATION


TREBOR CORPORATION

4047-A JONESBORO ROAD • FOREST PARK, GA. 30050 • 404-366-0957

The fast, efficient, low cost way to turfgrass aeration



NOW* - Topdress and/or overseed simultaneously with our seed-box attachment.
- "Special" no-tear bent grass rollers.



Care and maintenance with the "Aer-Way" will enhance the appearance of your turfgrass, and keep the traffic and playing areas of your parks, lawns and sportsfields greener.

The "Aer-Way" accomplishes this by cutting through thatch and hard pan allowing air, water and fertilizers to get into the ground where it does the most good, stimulating root growth and producing thick green turf. The tufts of grass produced by aeration will disappear with the first cutting.

Contact your nearest Holland office to arrange for a demonstration.

Holland Equipment Limited
P.O. Box 339
20 Phoebe Street
Norwich, Ontario N0J 1P0
Telephone: (519) 863-3414

Holland Hitch Western, Ltd.
Cloverdale, B.C.
V3S 5K1
Telephone: (604) 574-7491

Circle No. 144 on Reader Inquiry Card

Circle No. 115 on Reader Inquiry Card

JUNE 1984/WEEDS TREES & TURF 51

Stripe smut Stripe smut caused by *Ustilago striiformis* is a widespread disease of Kentucky bluegrass that has caused serious damage to many turf areas. It appears to be most serious in the Middle Atlantic states but can cause damage throughout the United States. Apparently, it is not a serious problem in Europe. Stripe smut has been observed for many years and can be found in many mature bluegrass stands.

The increased seriousness of stripe smut has probably resulted from the use of susceptible varieties such as Merion, Newport, and Windsor.

Spores present in the soil or carried on the seed germinate and systemically infect tillers and young seedlings. Long, narrow, gray to black stripes develop on the leaves. The gray stripes are unruptured sori. The black streaks result when the smut sori rupture and liberate mature spores. Following rupture of the sori, infected leaves curl from the tip downward and become shredded. Such tillers then die and disappear during periods of winter and summer stress. The result is a progressive weakening and deterioration of the turf in highly susceptible varieties.

Temporary periods of partial

recovery may occur. Most new stands of the susceptible varieties, if infected, do not deteriorate seriously until they are three to four years old which suggests the disease organism infects the plants slowly. Some turfs escape damage for longer periods.

Turf infected with stripe smut becomes much more susceptible to leaf spot and other diseases. It is also less likely to recover after periods of environmental stress. Resistant varieties offer the most practical means of control.

Some of the systemic fungicides are useful for control of stripe smut in turf.

Fusarium blight The *Fusarium* blight disease is causing serious damage to the more lush bluegrass turf in warmer regions. *Fusarium* blight is generally most severe during periods of high atmospheric humidity with daytime air temperatures of 80° to 95° F and night temperatures above 70° F. Temporary drought stress, high nitrogen levels and heavy thatch appear to contribute to the severity of this disease.

Control with fungicides such as benomyl requires proper timing and is expensive. More information is needed on varietal resistance to this disease(s) and on the stability of such resistance.

Observations at New Brunswick during the summers of 1972 through 1976 showed that varieties differed substantially in susceptibility to *Fusarium* blight. Under conditions of high nitrogen fertility and 3/4-inch mowing, the common types of Kentucky bluegrass including Park, Kenblue, Arboretum and South Dakota Certified showed extensive damage from *Fusarium* blight disease. Merion showed moderate damage. Fylking, Nugget, Geronimo, Delft, Modena and Enita had more damage than Merion. Windsor, Columbia, Enmundi, Parade, Adelphi, Rugby, Sydsport, Majestic, Vantage and Glade showed fair to good resistance in these tests.

Developing the required *Fusarium* blight resistance in Kentucky bluegrass will require intensive research.

Rusts (*Puccinia* and *Uromyces* spp.) A number of genera, species and races of rust infect Kentucky bluegrass. A variety resistant to one species or race of rust may be highly susceptible to another.

Stem rust (*Puccinia graminis*) causes considerable discoloration of susceptible varieties in many parts of the United States.

The disease is normally serious with poor vegetative growth under conditions such as low fertility and moisture stress. An improvement in growing conditions usually brings effective control as new leaf blades are removed by mowing before infection develops.

Merion and Touchdown are susceptible to stem rust, whereas Plush, Eclipse, Majestic, Adelphi, Bristol and Bonnieblue show moderate resistance to present races.

Leaf rust (*Puccinia poae-nemoralis*) is common on Kentucky bluegrass but is normally of concern on only the most susceptible varieties such as Vantage.

Dollar spot Dollar spot is a fungus disease caused by *Sclerotinia homoeocarpa*. While it is severe on many other turfgrasses, this disease is increasing in importance because of the widespread use of irrigation and susceptible varieties.

On Kentucky bluegrass it forms

Planting a new production field in rows. A band of activated charcoal protects the pure seed from herbicides used to eliminate weeds and old grass plants in the field.



COLUMBIA

CHALLENGER

GALAXY

MIDNIGHT

KENTUCKY BLUEGRASS BLEND

For the finest bluegrass turf on earth . . . reach for the stars



Columbia produces a moderately low growing, medium textured turf with a bright medium dark green color.

Columbia has exceptional Fusarium blight resistance and tolerance to stripe rust, good leaf spot, leaf rust, stem rust, dollar spot and stripe smut resistance. Columbia has good winter color and early spring green-up.



Midnight is the darkest green bluegrass available today. Midnight's low growing, compact dense turf has

good heat and cold tolerance. Midnight offers good resistance to leaf spot, stem rust, stripe smut and dollar spot. Midnight adds deep, dark green color to Galaxy . . . or any turf.



Challenger has a bright dark green color and is low growing with medium fine leaves and medium high

density. Challenger has good tolerance to stripe rust, and resistance to Helminthosporium leaf spot, leaf and stem rust, stripe smut, dollar spot and Fusarium blight. Challenger is fast emerging for quicker establishment.

Bluegrass blend rankings calculated by combining actual variety rankings at thirty-seven locations in the 1980 National Kentucky Bluegrass Trials. 1982 Data. 1 = Best

125	100	75	50	25	0
Galaxy Blend					COLUMBIA (25)
					MIDNIGHT (1)
					CHALLENGER (16)
					CHERI (20)
					ADELPHI (15)
					GLADE (24)
TOUCHDOWN (40)					
PARADE (46)					
AMERICA (11)					
					GLADE (24)
					RAM-1 (22)
					BARON (21)

GALAXY

qualifies for the Turf-Seed Blue Tag redemption program. Call or write for details.



P.O. Box 250, Hubbard, OR 97032
503/981-9574 TWX 510-590-0957

bright, medium dark green color. Georgetown has excellent early spring color. It is similar in appearance and performance to Parade, Haga, Banff, Rugby, Trenton, and Columbia in turf trials at Rutgers. Georgetown has good resistance to leafspot and most rusts and above average resistance to dollar spot and stripe smut.

Haga (Burlingham, Jacklin) was developed at the Weibullsholm Plant Breeding Institute in Landskrona, Sweden. Its experimental designation was WW AG 463. Haga is a moderately dark green, turf-type Kentucky bluegrass with medium high density, medium texture, and moderate aggressiveness. It has excellent cool temperature color retention and an exceptionally attractive early spring color. Late spring and early summer turf quality can be adversely affected by a stemmy appearance resulting from an abundance of reproductive tillers. Haga has good resistance to leaf spot, most leaf and stem rusts, and above average resistance to dollar spot and stripe smut.

Kenblue represents an attempt to reconstitute the type of common Kentucky bluegrass formerly harvested from naturally occurring stands of the famous bluegrass region of Kentucky. *Kenblue* represents a blend of seed harvested from selected seed fields of 8 to 15 years standing, situated on 12 farms located in seven central Kentucky counties. After blending, part of the seed was used to establish a breeders seed block at the University farm in Kentucky. The remainder was distributed to producers of certified seed. The first certified seed was harvested in 1967. In recent years, seed growers in Washington and Oregon have selected varieties with high seed production potential. As a result, some seed currently being sold as common Kentucky bluegrass is actually seed of one or more of these high seed producing varieties. Such seed lacks the genetic diversity of seed harvested from naturalized stands. In some situations, especially in short term tests, these varieties may perform better than common bluegrass. However, knowing the true variety has many advantages. Purchase of seed of Certified *Kenblue* Kentucky bluegrass will assure a consumer that he is getting bluegrass of known origin and wide genetic diversity.

Majestic (Burlingham) is a moderately low-growing, turf-type bluegrass with a rich dark green color and rather prostrate leaf blades. It has shown good resistance to leaf spot and leaf rust and moderate resistance to dollar spot, stem rust, and stripe smut. *Majestic* has excellent color especially during the cool seasons of spring and fall. It greens up early in the spring.

Merion originated from a single plant selection made by Joseph Valentine of

the Merion Golf Club, Ardmore, Pennsylvania in 1936. Until the early 1970's, *Merion* was the only commercially available variety with good resistance to the *Helminthosporium* leaf spot and crown rot disease. This quality made *Merion* the Cadillac of bluegrass varieties for well-maintained turf areas. Under conditions of high fertility and moderately close mowing, common Kentucky bluegrass and other susceptible varieties often show over 80% browning and thinning from melting-out under conditions where *Merion* normally shows little damage.

Compared with common Kentucky bluegrass, *Merion* has wider leaves, lower growth, darker green color when properly fertilized, higher resistance to *Helminthosporium vagans* and greater tolerance for close mowing. When properly managed, *Merion* can produce an attractive, dense, vigorous turf, highly resistant to weed invasion and capable of withstanding moderate wear.

Certain weaknesses of *Merion* are also recognized. Stripe smut, which frequently damages older stands, is the most serious problem. *Merion* is also highly susceptible to powdery mildew and should not be used in heavily shaded areas where it usually fails within several years. Stem rust susceptibility can be a problem, especially under conditions unfavorable to good growth. *Merion* may also be damaged by *Fusarium* blight. The latter disease is associated with turf weakened by high temperatures, excessive nitrogen, thatch accumulation, close mowing and with prolonged drought or improper watering. Many other bluegrass varieties can be damaged extensively under similar conditions. Susceptibility to dollar spot under low fertility conditions is becoming an increasing problem.

Merion is normally a vigorous, aggressive grass that will generally crowd out most other bluegrass varieties mixed with it under conditions of adequate fertility, frequent mowing and vigorous growth. Healthy, well-maintained *Merion* turf is seldom invaded by crabgrass or other weeds, but when weakened by stripe smut it is not aggressive.

Merion frequently requires and normally tolerates more fertilizer than common Kentucky bluegrass. However, excessive fertility is to be avoided, especially in older stands.

Merit (Full Circle) is a moderately low-growing, turf-type variety with a medium coarse texture, and medium density. *Merit* has moderate resistance to the *Helminthosporium* leaf spot and crown rot disease. It has moderate color retention during low temperatures and medium spring green-up. *Merit* has large seed and above average seedling vigor.

Midnight (Turf-Seed) originated as a

continued on page 56

small circular spots (two to five inches in diameter) which may merge to form large, irregular areas. The spots become a light straw color and dead grass occupies the center. Fine, white, cobwebby mold growth may be seen in these spots on mornings when a heavy dew is present and the fungus is active. Lesions on individual leaves appear as bleached areas extending the width of the leaf with constricted margins and a chocolate-brown border.

Moderate temperatures (68-86° F) and thick thatch are favorable for dollar spot buildup. Turf deficient in nitrogen tends to show more damage from dollar spot than turf which is adequately fertilized.

Different races of dollar spot respond differently to fungicide control and may attack varieties in a differential manner.

Insects

White grubs frequently cause extensive and serious damage to Kentucky bluegrass turfs as well as most other turfgrasses. These grubs are the larvae of several species of beetles and chafers including the Japanese beetle, the May and June beetles, the European chafer, and the northern masked chafer. Fully grown larvae are 1 to 2-cm long, white to grayish, with brownish heads and six distinct legs.

Severe infestations feeding on grass roots can cause serious thinning and death of the sod especially under drought stress. Birds, moles, and skunks actively feed on grubs tearing up the sod as they search for them.

Products containing resting spores of the milky disease organism, *Bacillus popilliae*, have been useful in keeping the Japanese beetle under moderate control.

Kentucky bluegrass is often severely damaged or completely killed by larvae of the bluegrass billbug. During May and June, adult billbugs lay eggs in grass stems near the crown. Upon hatching, the larvae are legless, white, fat, up to 1 cm in length and are present during July to mid-August. These larvae feed within the grass stem and later on the

MERIT

If they held
an **Olympics**
for Bluegrass,
Merit would
win the **Gold**

Consistently rated a winner for turf quality, color and sod strength, it has also been cited for its resistance to both leaf and dollar spot.

A variety with outstanding seedling vigor, it has won high ratings in shade trials and has excellent spring color.

The record proves Merit is a performer. It was ranked Number 1 for turf quality in a three-year University of California trial in competition with 43 varieties, topping Adelphi, Baron, Columbia, Cheri, Victa, Rugby, Sydsport, Ram I, and Parade.

It has also outperformed Baron, Nugget, Fylking, Adelphi, Glade, and Bonnieblue in various trials such as 1972 NE-57, in New York, in an eight-year Purdue study, a three-year Ohio trial and in a four-year Missouri test.

Note: Merit is marketed in Canada as "Regent Kentucky Bluegrass."

Merit...it's a Kentucky Bluegrass worth looking into


FULL CIRCLE, INC.

P.O. Box 49 Madras, Oregon 97741 (503) 475-3877



A Subsidiary of Cenex

Circle No. 113 on Reader Inquiry Card



single, highly apomictic aberrant plant selected from the open-pollinated progeny of F64-603, a selection made from an old lawn located near the Natural History Museum in Washington, D. C. Midnight is a persistent, low-growing, turf-type variety with the ability to produce a compact, dense turf with medium fine texture, a slow leaf extension rate, and a very dark green color. It has good heat and cold tolerance, fair shade adaptation, a slow spring greenup rate, and moderate low temperature color retention in late fall. Midnight has shown good resistance to leafspot, stem rust, strip smut, and dollar spot.

Monopoly (Pioneer Hi-Bred) was developed by Mommersteeg International of Vlijmen, Holland. Monopoly is a vigorous turf-type Kentucky bluegrass with medium texture, medium density, and a medium green color. It has shown excellent wear tolerance in a recent medium maintenance turf trial. Monopoly has moderately good resistance to leaf spot and dollar spot, and is promoted for its rapid germination.

Mystic (Lofts) was selected by Ralph E. Engel and Al Caravella from the 8th fairway of the Echo Lake Country Club, Westfield, NJ, where it had survived repeated sodium arsenite treatments used to destroy annual bluegrass in a fairway renovation program. An apparently identical plant was also found growing as a very large, attractive turf on the 14th fairway of the Seawane C.C. of Hewlett Harbor, NY, by Frank Curra and A.R. Mazur. Mystic is a low-growing, fine-leaved, turf-type Kentucky bluegrass. It displays a bright, medium-green color, and can produce a dense, attractive turf that is highly aggressive and which competes well with annual bluegrass. This makes it well suited for use on golf course fairways and tees. It has shown good resistance to powdery mildew and stripe smut, and moderate resistance to the *Helminthosporium* leaf spot and crown rot disease. Mystic is susceptible to dollar spot.

Nassau (Jacklin, Lofts) is a highly apomictic hybrid selected from the progeny of NJE P-59 x Baron. Nassau is a moderately low-growing, turf-type variety with medium wide leaves, and a medium dark green color. It is capable of producing an attractive turf of medium density. It can become very stemmy during late spring. Nassau has shown good resistance to the *Helminthosporium* leaf spot and crown rot disease. It has also shown above average resistance to dollar spot, red thread, pink snow mold, leaf rust, and stem rust. Nassau has attractive early spring color.

Newport originated as a single apomictic plant selected from coastal bluffs near Newport, OR. It is a moderately low-growing variety with rather dark green leaves and fairly good establish-

ment and turf-forming characteristics during the first two or three years. Newport has demonstrated good resistance to most current races of stem rust and powdery mildew, but is susceptible to leaf spot and stripe smut. Turf stands of Newport become very stemmy at seedhead setting time in June.

Older plantings of Newport have been heavily invaded by weeds and have frequently shown poor recovery from drought. Newport is often a short-lived type under New Jersey conditions, performing better than common Kentucky bluegrass during the first two or three years of a test; but it does poorly or dies as it ages. Its best use appears to be in blends with other more persistent bluegrass varieties.

Nugget (Pickseed) is a very distinctive variety. Nugget was found growing in Hope, Alaska. This variety produces a very dense, compact, rather low-growing turf which can be extremely attractive, especially in mid- to late-spring. Nugget has good to excellent resistance to leaf spot, most races of powdery mildew and leaf rust. Unfortunately, it appears to be susceptible to stem rust, dollar spot, *Fusarium* blight and aphids. This variety has good tolerance of close mowing and moderate shade when free of disease and insect damage. Nugget is very slow to start growing in the spring and has very poor early spring color in temperate climates. Nugget has frequently looked promising in preliminary turf trials throughout the northern part of the United States and Canada. Its performance in more southern areas has been very erratic.

Parade was developed in Holland. This moderately low growing, turf-type variety has medium texture and a pleasing moderately dark green color. Parade has good resistance to leaf spot, leaf rust and stem rust. It has above average resistance to dollar spot and stripe smut. Parade has excellent early spring color. Early summer turf quality is often adversely affected by an abundance of reproductive tillers.

Park resulted from an extensive selection and testing program initiated in Minnesota in 1937. A large number of plants were collected from old pastures and wasteland areas and tested for breeding behavior and agronomic performance. The 15 best apomictic strains were blended to produce Park, which was released in 1957. The variety has excellent seedling vigor and has shown moderate resistance to stripe smut, leaf rust and stem rust. Some of the component strains have good resistance to powdery mildew. Park is similar in appearance and growth habit to common Kentucky bluegrass and is susceptible to *Helminthosporium* leaf spot. Under New Jersey conditions, Park has often shown some advantages over common Kentucky bluegrass.

Plush (FFR Cooperative) originated as a single, highly apomictic plant selected from the lawn of Warinaco Park in Union County, NJ. Plush is an aggressive, persistent, moderately low growing, leafy, turf-type Kentucky bluegrass with medium texture, good vigor, medium high density, and a medium green color. It has shown good resistance to stripe smut and dollar spot and moderately good resistance to leaf spot and stem rust. Plush has shown good heat and drought tolerance.

Ram I (Jacklin, Lofts) was discovered growing on a putting green of the Webhannet Golf Club in Kennebunk Beach, Maine. Ram I is a moderately low-growing, leafy, turf-type cultivar with a medium texture and a rich, dark green color. Ram I has shown good tolerance of moderately close mowing and good early spring color. It has moderate resistance to leaf spot and stem rust, good resistance to stripe smut and most races of powdery mildew. It has moderate susceptibility to leaf rust and dollar spot.

Rugby (SPIC) is a moderately low-growing, turf-type variety with medium texture, good density, and a bright, medium dark green color. Rugby has a very attractive early spring color, the ability to stay green into late fall, and can maintain good winter color in protected locations. Rugby has shown good, or moderately good resistance to leaf spot, leaf rust, stem rust, dollar spot, stripe smut and *Fusarium* blight. Turf produced by Rugby is generally very stemmy during its reproductive period in late spring and early summer.

Sydsport (Burlingham) was developed in Sweden where it is reported to have good tolerance of the wear and abuse received on athletic fields. It has medium wide leaves and can produce a rather tight, dense sod of a medium light green color. Sydsport appears to have moderately good resistance to leaf spot and stripe smut but high susceptibility to dollar spot has been observed in some tests.

Touchdown (Pickseed) is a fairway selection from the National Golf Links of America located on Long Island. It has excellent resistance to leaf spot, stripe smut, leaf rust and most races of powdery mildew but is moderately susceptible to stem rust and dollar spot. Touchdown is a very aggressive turf-type variety with medium texture and a moderately dark green color. Like Warren's A-34 and Mystic, Touchdown has an excellent record of being able to compete well against annual bluegrass in closely mowed tests at Rutgers. These very aggressive varieties will normally dominate in blends and usually produce more thatch. Touchdown also shows promise of good performance in moderate shade.

continued on page 62

crown, bases of adjacent stems, and on roots and rhizomes.

Sod damaged by larvae is easily pulled from the soil with stems breaking off at the crown. The presence of fine, white, sawdust-like material at the base of each severed tiller is further evidence of billbug feeding. Billbug damage is most severe if turf is under moderate drought stress.

Studies at Nebraska and New Jersey show significant differences in cultivar resistance. Generally, the narrow-leaved, early maturing common types show less injury. However, some improved turf-type cultivars also show promising resistance.

Sod webworms are frequently destructive in lawns. In the warmer months, adult, grayish-white to beige moths frequently fly over lawns and pastures in late evening and drop eggs into the turf. Larvae live in silk-lined tunnels in the thatch during the day and feed on grass blades during the night.

Cutworms are the larvae of a number of species of night-flying moths. Cutworm larvae feed at night on grass leaves which they chew off close to the soil surface. Birds, particularly starlings, frequently visit turf infested with sod webworm and cutworm larvae and create numerous small circular holes in the sod.

Chinchbugs prefer sunny, hot, dry conditions and often cause serious damage, especially on south and west facing slopes. Plant injury occurs as a result of the insect sucking fluids from the plant. At the same time they inject salivary fluids. This disrupts the water-conducting tissues causing plants to wilt and turn yellow or brown. A fungus, *Beauveria* sp., is normally effective in reducing populations of chinch bugs when moisture conditions are adequate for fungal growth.

The adult frit fly is a very small, shiny, black fly. Larvae (maggots) hatch from eggs layed on leaves and in leaf sheaths. They immediately tunnel into and feed within grass stems. Initially the bluegrass has a yellow appearance as younger leaves emerging from infested tillers are damaged. As feeding progresses

FYLKING
FYLKING
FYLKING
FYLKING
FYLKING
FYLKING
FYLKING
FYLKING

Remember the name because you're going to hear a lot about it. Fylking Kentucky bluegrass lawn is something special. It forms a dense, thick turf, more disease and drought resistant. Greens up earlier, stays greener longer. Fylking can be cut as low as $\frac{3}{4}$ inch, even $\frac{1}{2}$ inch for home putting greens. Proven over many years of international certified testing. Ask your seed distributor for Fylking.


Fylking — the World's Fair grass.

FYLKING
KENTUCKY BLUEGRASS

U.S. Plant Patent 2887

Another fine, quality-controlled
product of Jacklin Seed Company.

BOOKSTORE



625-ADVANCES IN TURFGRASS ENTOMOLOGY edited by H.D. Niemczyk and B.G. Joyner
A complete account of the facts presented at the 1980 Symposium on Turfgrass Insects. 200 photographs, tables and graphs make this volume an indispensable reference for anyone connected with the turfgrass industry or research. **\$24.95**

010-ADVANCES IN TURFGRASS PATHOLOGY by Dr. B.G. Joyner & Dr. P. Larsen
Leading U.S. turf pathologists report on turfgrass diseases, pythium blight, snow molds, fairy rings, leaf spot of Kentucky Bluegrass in Minnesota, initial and field fungicide screening, turfgrass disease resistance, etc. Contains new ideas on how to combat turfgrass problems. **\$27.95**

655-TURFGRASS: SCIENCE AND CULTURE LABORATORY MANUAL by Beard, DiPaola, Johns and Karnok
Class tested for over three years, this manual provides fourteen exercises which can be easily adapted to your particular course structure. Exercises involve students in vegetative and seed identification, equipment and irrigation system selection and operation, problem solving of typical math problems involved in turfgrass operations and the diagnosis of problems with emphasis on weeds, diseases and insects. Encompasses both warm and cool season turfgrass. **\$12.95**

645-MANAGEMENT OF TURFGRASS DISEASES by J.M. Vargas
Identifies turfgrass diseases by description and illustration. Includes a holistic approach to healthy turf and lawns. Presents practical management strategies for golf courses, lawns and athletic fields. 204 pages. Illustrated **\$24.95**

615-TURF MANAGEMENT FOR GOLF COURSES by James B. Beard
Written by an eminent turfgrass researcher, this USGA sponsored text is an ideal reference and "how to" guide. Details all phases of golf course operations including course design and construction, turf management, course administration, irrigation, equipment and disease and pest control. Fully illustrated. **\$45.00**

110, 120-TURF MANAGERS' HANDBOOK by Drs. William Daniel and Ray Freeborg
This specially designed manual by leading turf specialists is a comprehensive, organized approach to turfgrass science and care. An easy, on-the-job reference for planning, purchasing, hiring, construction and plant selection. **\$28.95 hardcover, \$23.95 paperback**



665-ARBORICULTURE: THE CARE OF TREES, SHRUBS AND VINES IN THE LANDSCAPE by Richard W. Harris
Provides comprehensive coverage of complete planting, site analysis, preparation and special planting methods, fully-detailed coverage of fertilization, irrigation and pruning guidelines on preventative maintenance, repair and chemical control, how-tos of diagnosing plant problems, practical data on non-infectious disorders, diseases, insects and related pests and pest management. **\$34.95**



345-COST DATA FOR LANDSCAPE CONSTRUCTION 1984 Kathleen W. Kerr, Editor
An updated unit cost data reference for designers and cost estimators. Developed to fill the tremendous need for detailed landscape construction cost data. Laid out in easy-to-use CSI format. Annual. **\$28.75**

395-LANDSCAPE ARCHITECTURE by John Ormsbee Simonds
A Manual of Site Planning and Design
This long-awaited second edition outlines and analyzes the complete landscape process from site selection to finished project. In simple and clear terms it describes various planning constraints imposed by the forms, forces and features of the natural and built landscape. **\$34.95**

300-LANDSCAPE DESIGN: A PRACTICAL APPROACH by Leroy Hannebaum
Geared for the commercial designer/salesperson, this is a one-stop guide to the landscape design process. Covers the entire highly competitive field including design analysis techniques, pointers on land forms, specialized business landscaping methods, environmental design guidelines, specifications estimations, bids. **\$19.95**

360-RESIDENTIAL LANDSCAPING I by Theodore D. Walker
Provides an in-depth discussion of the planning, design and construction phases of residential landscaping. Illustrated with the work of professional landscape architects. Covers everything from analyzing the site to constructing the landscape. **\$22.50**

405-WOODY ORNAMENTALS by Partyka, Joyner, Rimelspach, Carver
Illustrates plant identification characteristics. Organized in two basic sections: plant identification and plant disorders, this text utilizes 430 color photos, 430 line drawings and 45 black and white photos to simplify identification. Goes into detail on plant identification and description as well as plant problems such as diseases, chemicals, insects, animals and physiological disorders. **\$27.00**




800-THE GOLF COURSE by Geoffrey S. Cornish and Ronald E. Whitten
The first book ever to give the art of golf course design its due, and golf course architects the credit and recognition they deserve. 320 pages and approximately 150 color and black and white photographs. Traces the history and evolution of the golf course, analyzes the great courses, shows how they were designed and constructed. **\$35.00**

675-HILLIER'S MANUAL OF TREES & SHRUBS by H.G. Hillier
This 5th edition is a unique reference source when selecting new shrubs and distinguishing between those already in the garden. Provides descriptions of over 8,000 plants with information on color, size and habit. Sections on nomenclature and classification, planting, pruning and garden planning, plus an illustrated glossary, add to the usefulness of a book that will prove invaluable to both beginners and experienced gardeners. **\$20.50**

635-IRRIGATION PRINCIPLES AND PRACTICES by Hansen, Israelsen and Stringham
A new fourth edition of this highly successful textbook presents essential concepts pertaining to water conveyance, application, storage in the soil and use by the plants. Basic underlying principles that govern irrigation practices are stressed. Generalized concepts are discussed and newly formulated practices are examined. **\$39.95**

BOOKSTORE



575-MODE OF ACTION OF HERBICIDES by Floyd M. Ashton and Alden S. Crafts
Provides worldwide body of information on each class of herbicide. Cross-indexed tables of common and trade names of herbicides are included. New herbicides accepted since 1972 are listed in this revised second edition. Excellent practical reference for specialists in field of weed science. **\$58.95**

790-RECREATION PLANNING AND DESIGN by Seymour M. Gold
A comprehensive look at recreation needs for parks and how they can design the park facility for the community. Book's content can help justify construction and maintenance needs. **\$39.50**

565-WEEDS by Walter C. Muenscher
Second edition. Premier text for identification and basic natural history for weeds found in the continental United States and Canada. Ecological data on weed biology combined with excellent keys and plant descriptions makes this an essential reference book. **\$37.50**

ADDITIONAL TITLES

- 340 - CONSTRUCTION DESIGN FOR LANDSCAPE ARCHITECTS \$39.50
- 410 - DISEASES & PESTS OF ORNAMENTAL PLANTS \$32.50
- 660 - DISEASES OF SHADE TREES \$23.50
- 610 - DISEASES OF TURFGRASSES \$30.00
- 350 - HANDBOOK OF LANDSCAPE ARCHITECTURAL CONSTRUCTION \$48.50
- 510 - HORTUS THIRD \$125.00
- 690 - INSECTS THAT FEED ON TREES & SHRUBS \$47.50
- 370 - LANDSCAPE OPERATIONS: MANAGEMENT, METHODS & MATERIALS \$20.95

- 545 - MODERN WEED CONTROL \$21.50
- 700 - THE PRUNING MANUAL \$15.95
- 720 - SHRUB IDENTIFICATION \$8.00
- 750 - TREE IDENTIFICATION \$9.95
- 760 - TREE MAINTENANCE \$35.00
- 640 - TURF IRRIGATION MANUAL \$22.95
- 620 - TURF MANAGEMENT HANDBOOK \$18.00
- 650 - TURFGRASS MANAGEMENT \$21.95
- 630 - TURFGRASS:SCIENCE & CULTURE \$27.95
- 570 - WESTCOTT'S PLANT DISEASE HANDBOOK \$36.50

CLOSEOUTS

ORDER THESE TITLES AT SPECIAL REDUCED PRICES!

- 455 - THE GRAFTER'S HANDBOOK \$16.95
- 460 - GREENHOUSE ENVIRONMENT \$21.20
- 335 - LANDSCAPE DESIGN THAT SAVES ENERGY \$8.50

Mail this coupon to: Book Sales
Harcourt Brace Jovanovich Publications
One East First Street, Duluth, MN 55802

Name _____
Street Address _____
P.O. Box Number _____
City/State/Zip _____
Signature _____ Date _____
Phone Number _____
Purchase Order Number _____

Please send me the following books. I have enclosed payment* for the total amount.

Please charge to my Visa, MasterCard or American Express (circle one)
Account Number _____ Expiration Date _____

BOOK NUMBER AND TITLE	QUANTITY	PRICE	TOTAL PRICE

*Please add \$3.00 per order plus \$1.00 per additional copy for postage and handling.

Please allow 6-8 weeks for delivery.
Prices subject to change.
Quantity rates available on request.

(postage & handling)

Total Enclosed _____

WTT 64

National bluegrass test shows regional differences

Performance of Kentucky bluegrass varieties varies by location. To point out regional differences, the National Turfgrass Evaluation Program was developed in 1980. Jack Murray, agronomist, USDA Agricultural Research Service, Beltsville, MD, was selected national coordinator.

Thirty-seven test gardens were set up in 22 states and two provinces of Canada for the trial. Data for 1982 and 1983 has been collected.

The tests judge the varieties by location, and by a number of other factors; including disease resistance, seasonal condition, leaf texture, density, drought tolerance, sod strength, and insect resistance.

Murray cautions that the data is just for the second year of a multi-year evaluation and that seed companies presented seed from just one lot for the tests. Variation by year and between seed lots should be considered. Although varieties were listed in order from best to worst, the difference between some may be considered insignificant.

When data from all locations and for all months was combined and averaged, those with the best quality overall were

Eclipse, Midnight, Bristol, Adelphi, America, Banff, Baron, Enmundi, Majestic, Challenger, Ram-1, Sydsport, Bonnieblue, Cheri, Columbia, Glade, Rugby, and Victa. A number of experimental varieties were included among the top national performers.

The leaders in the home state, Kentucky, were America, Eclipse, Banff, A-20, Nassau, Merion, Mystic, and Monopoly.

Many bluegrasses performed well in Delaware. The best were Midnight, Bristol, Enmundi, Mystic, Kimono, Bonnieblue, Challenger, Majestic, and America. Maryland was led by Merit, Victa, Baron, and Kimono. Virginia's four locations gave the edge to Eclipse, America, Wabash, Columbia, Cheri, Enmundi, Adelphi, Bristol and Midnight.

Georgia's best bluegrasses were Columbia, Birka, Enoble, Monopoly, Sydsport, Adelphi, Majestic, Midnight, and Baron. North Carolina leaders were Baron, Enmundi, Cheri, Glade, and Parade.

Iowa's favorites were Midnight, Majestic, Challenger, Ram-1, Glade, Bonnieblue, Victa, Merit, Nassau, Nuggett, and Bristol. Nebraska, with four test areas, favored Enoble,

Majestic, Wabash, and Geronimo.

Eclipse, Adelphi, Sydsport, Bonnieblue, Touchdown, A-20, and Enmundi performed best in Illinois. Baron, Challenger, Glade, Columbia, Touchdown, Adelphi, Enmundi, Ram-1, Cheri, and Victa showed them in Missouri. Midnight took Kansas by storm with closest rivals Bristol, Eclipse, Challenger, Bonnieblue, Victa, America, and Majestic.

New Jersey results were better than New York's. The leaders in New Jersey were Eclipse, Challenger, Midnight, Bonnieblue, Kimono, Glade and Majestic. New York's two locations favored Eclipse, Monopoly, Midnight, Sydsport, Cheri, and Wabash.

In the Northwest, where most of the seed is grown, Oregon winners were Eclipse, Sydsport, Columbia, Midnight, Majestic, Challenger, and A-20. Washington leaders were Ram-1, Rugby, Banff, Shasta, and Fylking.

For the complete report, write USDA Science and Education Administration, Plant Genetics and Germplasm Institute, Beltsville, MD 20705. □
Written by Bruce F. Shank, executive editor

these infected tillers die. Careful examination of the tiller base will reveal one or more active, small white maggots with two characteristic mouth hooks visible with a good hand lens.

The greenbug aphid has recently become a serious pest of Kentucky bluegrass in a number of locations throughout the mid-western United States. The piercing mouthparts of the greenbug are inserted into leaf blades to suck out plant fluids. This probing and the injection of salivary fluids causes leaf tissues to turn yellow to light orange, then even darker orange and finally to brown as the plant may die under very heavy

infestations. Scientists at Beltsville, MD, are developing Kentucky bluegrasses with good resistance to the greenbug aphid.

Growth habit and turf-forming properties

The growth habit of Kentucky bluegrass is influenced greatly by day length, light intensity and temperature.

During short days, Kentucky bluegrass assumes a more decumbent (spreading) growth habit, has a slower rate of leaf elongation, and tillering is more abundant. During long days, growth is more erect and leaf elongation is more rapid. Reproductive development

also occurs during the long days of late spring.

High light intensity increases photosynthesis and promotes the development of thick, sturdy leaves and a deep green color. Low light intensity produces weak, thin, etiolated (pale) plants with a rapid rate of leaf elongation.

Common Kentucky bluegrass and varieties such as Park, Delta and Kenblue have a rather erect growth habit with a rapid rate of leaf elongation. Such varieties do not tolerate high nitrogen fertility and close mowing, especially during the spring and summer seasons.

During the long days of spring and summer these varieties make noticeably taller growth. This results in the removal of a higher percentage of the leaf area and makes maintenance of good turf in late spring and summer difficult. Carbohydrate food reserves are depleted and such varieties become highly susceptible to damage from the Helminthosporium leaf spot and crown rot disease.

Varieties such as Nugget, America, Eclipse and Glade appear to exhibit the short day length response of decumbent growth and slow leaf elongation through much more of the year than the common type bluegrass varieties. Additional research related to differential varietal growth response to day length should be of great value in breeding bluegrasses with better turf-forming properties and reduced mowing requirement.

Tolerance of close mowing

For golf course fairways, the turf should make an attractive, uniform carpet which is dense enough to give a good lie to the ball. It must also be able to heal divots rapidly, tolerate considerable traffic and resist the invasion of annual bluegrass.

Frequent, close mowing, adequate fertility and water are needed to produce the firm, dense turf required to support the ball above the soil surface. A dense turf has a much higher population of tillers per unit area which causes increased competition between tillers. This, plus severe defoliation by low fairway mowing and ample fertilization weakens the grass. It develops a less extensive root system and is more subject to drought damage and disease attack.

Close cutting and frequent watering encourages rooting above the soil surface and thatch buildup. This favors many disease organisms. Also, damage from disease is more apparent on an otherwise attractive, uniform closely cut turf.

Kentucky bluegrasses have some tolerance of the close mowing and other factors associated with the production of the dense,

firm, aggressive turf desired on fairways. They have the best chance of success with high light intensity, cool temperatures and moderate humidity.

In less favorable climates, improved varieties and better management are needed for successful results. Many of the current Kentucky bluegrass varieties including Nugget, Warren's A-20, Touchdown, Bonnieblue, Eclipse, Birka, Fylking, Majestic, Merion, Adelphi, Glade, Sydsport, Victa, Cheri, RAM 1, and Baron have characteristics which make them more suitable for close-cut fairways than Common Kentucky bluegrass and other erect-growing leaf-spot-susceptible varieties. Unfortunately, each of these varieties has some weakness.

Proper blending of seeds of these improved varieties might help but will not solve all the potential problems associated with fairway turf. Those who use the turf-type ryegrasses as a major fairway grass will find maintaining Kentucky bluegrass in the mixture helpful. New Kentucky bluegrass selections collected from close-cut areas and those generated in hybridization programs give promise of additional improvement.

Heat tolerance

Kentucky bluegrasses with greater tolerance of summer heat and drought conditions common to the transition zone would be of great benefit.

Most of our attractive, dense, lower-growing, turf-type varieties were selected in the cool summer climate of Northern Europe and from other breeding and evaluation tests located in cool environments. Many of these varieties are often disappointing in southern trials.

An extensive program to collect and evaluate adapted germplasm from summer stress areas of the Mid-Atlantic areas should provide varieties with improved summer performance and dependability. Under conditions of moderately low nitrogen fertility and high cut, varieties that typify common types, such as Kenblue, have survived well in the transition zone.

PROFIT MACHINES!

Landscaping professionals everywhere are rapidly expanding their business with FINN's T80 HydroSeeder and B50 Mulch Spreader. These sturdy workhorses have greatly increased their day's work production without extra man-power—and that means their profits are rapidly expanding, too!



T80 HydroSeeder

The T80 keeps right on going with its 800-working-gallon capacity. One tank-load can seed, fertilize, and mulch up to 1/4 acre—or up to 3 acres with seed and fertilizer only at distances up to 90 feet. Standard equipment includes a variable speed agitator and centrifugal pump. Result: The T80 will mix, suspend, and spray heavier concentrations of dry solids, powders, liquids, and fiber mulches. Among its options is a high-pressure pump for spraying herbicides or fungicides. Available as skid or trailer mounted.



B50 Mulch Spreader

The B50 Mulch Spreader can separate and spread up to 5 tons of straw or hay mulch per hour at ranges up to 55 feet. Its outstanding ground-coverage uniformity saves 50% material costs over hand-spreading methods. Available as skid or trailer mounted.

These FINN profit machines are designed for low operating costs, dependability, and low maintenance. Call or write for all the facts plus information on our complete product line. Remember—they're backed by a half-century of FINN's dedication to quality and service!

FINN CORPORATION

P.O. BOX 8068 • 2525 DUCK CREEK RD.
CINCINNATI, OHIO 45208 U.S.A.
PLANTS: CINCINNATI, OHIO • KNOXVILLE, TENNESSEE
PHONE (513) 871-2529 TOLL FREE (800) 543-7166

Troy was selected from a seed lot introduced from Turkey. It is a tall, erect-growing grass highly susceptible to leaf spot. Troy has been useful as a pasture grass in Montana region but performs very poorly as a turfgrass.

Vantage (International Seeds) was developed by O.M. Scott and Sons of Marysville, OH. Vantage is a persistent Kentucky bluegrass with deep spreading rhizomes, a medium dark green color and a medium texture. It has very good heat and drought tolerance. Vantage has performed well in higher cut, medium maintenance turfs in the Middle Atlantic region of the U.S. where it has shown good resistance to the Fusarium blight disease. However, this same variety has been seriously damaged by Fusarium blight disease in California trials. Vantage has good resistance to stripe smut and dollar spot and moderate resistance to leaf spot. It is susceptible to leaf and stem rusts.

Victa (Scotts) was developed by O. M. Scotts. It has medium broad leaves, a moderately low-growing, turf-type growth habit and a medium dark green color. The variety has shown moder-

ately good resistance to leaf spot. It has shown moderate resistance to leaf rust, stem rust, dollar spot and powdery mildew in New Jersey tests. Victa is moderately slow in spring green-up. It has large seed and rather good seedling vigor.

Wabash was developed at Purdue University. It is a vigorous variety with exceptionally good rhizome development and ability to recover from stress. It produces a turf of medium density, medium wide leaves, and a bright, medium green color. Wabash often shows substantial damage from the Helminthosporium leaf spot and melting-out disease, especially when mowed closely. However, it recovers well and looks very attractive by fall. It showed the best fall recovery of all bluegrasses studied in a test at North Brunswick which had received severe summer stress.

Warren's A-34 (Warren's) is a vigorous, disease resistant variety with somewhat better shade tolerance than most other Kentucky bluegrass varieties. When maintained at a 2-inch growing height, it will tolerate 65% shading.

A-34 does rather well in full sun, producing a dense, medium green turf with moderately good resistance to stripe smut, powdery mildew and leaf spot. It also performed well in wear tolerance trials in Michigan.

Varietal Blends Admitted weakness of all currently available bluegrass varieties has caused many turf workers to recommend the use of varietal blends for better lawns, fairways and most other types of turf. It is hoped that the weakness of one variety will be covered up by a complementary strength of other variety. This may or may not be true depending upon a number of complex ecological factors. We need much more research data on ecology, long-term performance and regional adaptation of bluegrass blends.

Research at Rutgers strongly suggests that varieties with good resistance to both stripe smut and Helminthosporium leaf spot should be included in all turfgrass blends recommended for use on intensely maintained turf areas. Also, one or more should have high tolerance of close-cut unless the turf will be mowed high. □



Seed samples are tested in a lab after cleaning. Contents are reported on the seed label.

Under conditions of somewhat closer mowing and higher fertility the lower-growing, wider leafed, open types having extensive deep rhizomes, such as Vantage, have performed better. Merion Kentucky bluegrass has shown above average summer performance when managed properly and when disease is not a problem.

Color

Visitors at experimental plantings of Kentucky bluegrass selections and hybrids are impressed by the great diversity of shades of green observed.

Mystic has a very attractive bright light green color. Adelphi and Glade have bright, dark colors. Some selections like Bonnieblue, Parade, Columbia and Majestic retain excellent color into the winter and green-up early in the spring. Others like Midnight and Baron go dormant in late fall and green-up later in the spring. Still others like Nugget green-up very slowly in the spring.

Many types show a pronounced purplish cast in late fall, winter and early spring, whereas some, such as Parade, Columbia, Rugby, and Bonnieblue appear to lack this purplish pigment. **WT&T**

SYDSPORT

Kentucky
Bluegrass



HAGA

Beautiful, low maintenance turfs . . .

SYDSPORT and HAGA; two beautiful Kentucky Bluegrass varieties from the world famous turfgrass firm, E.F. Burlingham & Sons.

Both exhibit good seedling vigor and rapid development, resulting in a turf that greens up early in the spring and retains its bright green color throughout the entire growing season.

If you are concerned with wear tolerance and recovery from damage, seed SYDSPORT or HAGA to protect your high wear turf areas . . . and your reputation.

FOR MORE INFORMATION CONTACT:



E.F. Burlingham & Sons.

P.O. Box 217 • Forest Grove, OR 97116
(503) 357-2141 • Telex: 36-0274
Cable: Burlingham

Building Self-Preservation Into Turf Equipment

by Thomas M. Carter, Vice President, Engineering, Jacobsen Division of Textron Inc.



Overheating can destroy an engine. Simple air intake systems are being improved with devices which prefilter ambient air from a high location and ducts it to the engine. The result is longer engine life.

The evolution in turf equipment hasn't been confined to improving or expanding the tasks these machines perform, despite their growing sophistication.

Along with doing the work expected of them better and in less time, turf and grounds care people now more than ever demand a longer trouble-free life with reduced maintenance as an added bonus.

Public and private mandates haven't changed for attractive and useful parks, golf courses, recreational and other turf areas. Only the budgets have.

The turf equipment engineer's assignment has been very clear: design machines that do more, last longer and are easier to maintain.

While hydraulics and other systems and components that together mow, aerate, sweep or

perform many other chores, have been constantly improved, nothing moves without power.

Evaluate power

In other words, the demands on our engines have never been greater, making it mandatory for a variety of power plants to undergo a process of intense evaluation before one is selected.

Tough duty

In the same way, the turf machine user should evaluate the engine with the same care he applies to assessing overall function and key features like hydraulic versus conventional mowing.

The importance of engine evaluation can hardly be overstated when one considers the tough world of turf care. These machines are run for long periods of time under varying load condi-

tions. Oftentimes, operators have little or no sensitivity to mechanical devices, overloading their machines and routinely subjecting them to other abuses. And, as budgets tighten, regular maintenance sometimes suffers.

On top of all that, turf machines operate in harsh environments of dust, dirt and other particulates that are often made even worse by high ambient temperatures and a surface seemingly designed to test every fastener.

Seasonality plays a role, too, with year-around turf use for some regions and six months of service for others.

There are the so-called "systems machines", such as our Turfcat II line, which with attachments like a dozer blade or snow thrower are used for clearing walks and drives in winter — or sweeping anytime of the year with a rotary broom. These versatile machines and the diverse and frequently harsh conditions under which they operate call for engines with a high degree of self-preservation designed and built into them.

Smooth power

In mid-sized turf equipment perhaps the first item for your evaluation is the number of cylinders. Two-, three- and four-cylinder engines — gas or diesel — will outwear and be smoother than single-cylinder power. Though there are exceptions to the rule, generally multi-cylinder engines are more sophisticated with features that add up to better performance and increased longevity. But, no matter how many cylinders, an engine will depend on the machine around it and system components to be designed in such a way as to help assure a long, productive life.

FUNGISOL[®]



FUNGISOL[®] **The first fungicide developed** **specifically for tree diseases**

FUNGISOL. Just registered for arborists. EPA has just registered FUNGISOL (contains DEBC) for suppression of a broad range of fungus diseases on shade trees and ornamentals.

New, systemic FUNGISOL has been tested for 8 years. It's the only fungicide developed specifically for the Mauguet Tree Injection Process.

FUNGISOL is registered for use against fungi which occur in the complex causing oak decline in such diverse areas as Texas, Florida, Pennsylvania, Illinois, Nebraska and California. Also registered for use against pine pitch canker.

In most instances the rate of recovery is greatly enhanced when STEMIX[®] or Zinc STEMIX is applied with FUNGISOL, usually on the same feeder tube. In some situations, INJECT-A-CIDE[®] has also been applied at the same time to control an insect problem.

A research report which highlights the results of a broad range of field testing is available from your Mauguet distributor or by writing:

J.J. MAUGUET COMPANY

P.O. Box 3422 • Burbank, California 91504 (800) 423-2699

Mauguet Process Patented In U.S. And Abroad


Why a Lawn Genie does more work.

The Lawn Genie Pick-up Mower is available in ten models from 36" to 72" engine or PTO driven, lift or pull type.

Quick flip deflector panel easily changes from pickup to ground discharge.

Reversible cutting knives quickly interchange with thatching blades for real versatility.

Adjustable, full-width gauge roller assures even cutting.



Spring, summer, fall. The Lawn Genie from M-C keeps working.


Quick dump hopper empties from the tractor seat.

Large 8⁵/₈" diameter balanced rotor for smooth, dependable operation.

Ask your local M-C Dealer for a demonstration or call or write M-C for more information.



Mathews Equipment Co.
Crystal Lake, IL 60014
815/459-2210



Circle No. 123 on Reader Inquiry Card

Carburetion

Gas and diesel engines alike require a relatively specific ratio of clean air-to-fuel to function properly, so the air cleaner for each power plant should be sized to filter out damaging dust and dirt of turf care while allowing proper aspiration. One way to decrease performance and reduce engine life is to upset that ratio by not replacing the air filter at proper intervals and letting it become clogged.

To avoid clogging turf equipment engineers have gone a step further from the traditional placement of the air cleaner atop the carburetor on the engine. We take in air through a screened opening just behind the operator's seat and then duct it to the cleaner. The cleaner itself is non-traditional, too, because we use a large, industrial unit.

Cooling

Cooling the engines of turf equipment is far more involved than keeping the temperature in line on cars, trucks or even many agricultural machines. It's not enough

to draw air in the conventional manner for either liquid- or air-cooled power plants. Turf machines require controlled air via special ducting, using pre-cleaners to remove dirt that would eventually build up on radiator cores or cooling fins, thus raising engine temperatures to damaging levels.

In addition to looking for ducting and pre-cleaners, check for the position of the air intake. It should draw from the ambient atmosphere; not from air preheated by the engine.

Lubrication

Lawn and garden tractor engine technology simply can't be used for turf machines. Usually that type of power has splash lubrication. That may be quite satisfactory for several hours of periodic work, but the duty cycle of the turf machine requires full flow lubrication to make certain all moving parts constantly receive a quantity of oil that relates to the work being performed. As power demands and rpm increase, so does the flow of oil to reciprocating

and rotating parts.

But, engine oil can do more than lubricate. It also can cool, so watch for added touches like a larger remote oil filter and possibly an oil cooler. Check for a warning light on the control panel to let you know operating temperature has reached a level that, if sustained, could damage the engine. The latest machines may even have heat sensors that signal the operator.

Inside the engine

You can easily see engine peripherals — cooling, carburetion, exhaust system, and so forth. But what you can't see beyond the new paint of the engine may be even more important to performance, life span, and maintenance.

Take the valves. In addition to admitting the air-fuel mixture and exhausting combustion gases, valves maintain compression, and thus, the power you required when you specified the machine.

To make certain power remains consistent, you'll want to look for a variety of features, such

Circle
the
Reader
Service
numbers
of those
items of
interest
to you.

For fastest response, use the peel-off label from the front cover.

NAME	
TITLE	
FIRM	PLACE COVER LABEL HERE
ADDRESS	
CITY	
STATE	ZIP

101	116	131	146	161	176	191	206	221	236	251	266	281
102	117	132	147	162	177	192	207	222	237	252	267	282
103	118	133	148	163	178	193	208	223	238	253	268	283
104	119	134	149	164	179	194	209	224	239	254	269	284
105	120	135	150	165	180	195	210	225	240	255	270	285
106	121	136	151	166	181	196	211	226	241	256	271	286
107	122	137	152	167	182	197	212	227	242	257	272	287
108	123	138	153	168	183	198	213	228	243	258	273	288
109	124	139	154	169	184	199	214	229	244	259	274	289
110	125	140	155	170	185	200	215	230	245	260	275	290
111	126	141	156	171	186	201	216	231	246	261	276	291
112	127	142	157	172	187	202	217	232	247	262	277	292
113	128	143	158	173	188	203	218	233	248	263	278	293
114	129	144	159	174	189	204	219	234	249	264	279	294
115	130	145	160	175	190	205	220	235	250	265	280	295

WEEDS TREES & TURF

JUNE 1984
This card expires August 15, 19

MY PRIMARY BUSINESS AT THIS LOCATION IS:
(PLEASE CHECK ONE ONLY IN EITHER A, B OR C)
**A. LANDSCAPING/GROUND CARE AT ONE OF THE FOLLOWING
TYPES OF FACILITIES:**

- 0005 Golf courses
0010 Sport complexes
0015 Parks
0020 Rights-of-way maintenance for highways, railroads & utilities
0025 Schools, colleges & universities
0030 Industrial & office parks/plants
0045 Condominiums/apartments/housing developments/hotels/resorts
0050 Cemeteries/memorial gardens
0060 Military installations & prisons
0065 Airports
0070 Multiple government/municipal facilities
 Other type of facility (please specify)

B. CONTRACTORS/SERVICE COMPANIES/CONSULTANTS:

- 0105 Landscape contractors (installation & maintenance)
0110 Lawn care service companies
0125 Landscape architects
0135 Extension agents/consultants for horticulture
 Other contractor or service
(please specify) _____

C. SUPPLIERS:

- 0205 Sod growers
0210 Dealers, Distributors
 Other supplier (please specify) _____

Approximately how many acres of vegetation do you maintain or
manage? _____

What is your title? (please specify) _____

I would like to receive (continue receiving) WEEDS TREES & TURF
each month: YES NO

Your Signature: _____ Date: _____



BUSINESS REPLY CARD

FIRST CLASS PERMIT NO. 665 DULUTH, MINNESOTA

POSTAGE WILL BE PAID BY ADDRESSEE

READER SERVICE DEPARTMENT

WEEDS TREES & TURF

POST OFFICE BOX 6049
DULUTH, MINNESOTA 55806-9749

NO POSTAGE
NECESSARY
IF MAILED
IN THE
UNITED STATES



**GET
MORE
FACTS**

**The Professional Lawn Care
Association of America
1984 Conference & Trade Show
November 12-15, Tampa, Florida**



PLCAA '84 Tampa
NOVEMBER 12-15

MORE IN '84! Mark your calendar. It's not too early to start planning for the lawn care industry's biggest week of the year — the Professional Lawn Care Association of America's 1984 Show and Conference.


And what a week it will be! PLCAA '84 is shaping up to be the biggest, best gathering of powerhouse speakers and exhibitors in our five-year history. The seminar program has been expanded to offer over 16 hours of educational programming hosted by a new lineup of outstanding speakers, problem solvers all.

In addition, last year's popular "Workshop" series will offer even more hours of "how to" advice on business basics like purchasing, business expansion, and promotional techniques. Subjects aimed at boosting the bottom line of your operation.

IT'S SHOWTIME, FOLKS! Virtually every major vendor and supplier to the lawn care industry will be among the over 100 exhibitors at PLCAA '84. Longer show hours and more floor space in the beautiful Curtis Hickson Convention Center will give showgoers an unparalleled opportunity to see, touch, and compare all that's new in lawn care.

BRING THE FAMILY! Take advantage of Tampa, heart of Florida's funland. PLCAA '84 offers features, options, and vacation opportunities designed with fun in mind. Features like a special Spouse Program full of fun and surprises. Options like a week-long post-conference Caribbean cruise on the luxurious New Amsterdam, newest ship in the Holland America Line. Opportunities like a chance to visit those Florida wonderlands you've been dreaming about — Disney World, Epcot Center, and Sea World. Indicate your interest on the coupon below and details will be sent to you.

'84 is the year. Tampa is the place. Go for it!



Delta Airlines have been appointed official carrier for PLCAA '84. Delta has regular scheduled flights from most major US cities to Tampa and will be offering discounted fares to all PLCAA '84 participants — up to 30% off regular prices.



YES! I'm interested in attending PLCAA's 5th Annual Conference and Trade Show. Send me more information on:

REGISTRATION EXHIBITING MEMBERSHIP POST CONFERENCE CRUISE

NAME _____

COMPANY _____

ADDRESS _____

CITY, STATE, ZIP _____

Mail to: Professional Lawn Care Association of America
1225 Johnson Ferry Road, NE • Suite B220 • Marietta, Georgia 30067

as rotators for both intake and exhaust valves. These cause valves to turn minutely as they actuate to help assure proper seating. Valve stem wipers are another desirable feature. These non-metallic "sleeves" prevent carbon buildup on the stems by wiping off traces of that element with every stroke. Excessive carbon on the stems can deteriorate their guides, allowing sump oil to escape into combustion chambers, reducing plug performance and increasing oil consumption.

High temperatures created by sustained operation could cause valve warpage were it not for the special high-carbon alloy steel (such as Stellite) used for the valves in some engines. Besides loss of compression, severe warpage can lead to a breakdown.

The power you count on is primarily maintained by compression rings pressure fitted on the pistons. Those engines that use high-grade steel in compression rings will deliver many thousands more hours of performance. The

difference in head gaskets can spell blow-by. A top quality product such as the metal-clad Graphoil gasket resists erosion from extreme combustion pressures, gasses and sustained high operating temperatures. The small extra cost of a quality product is negligible when compared with the downtime involved in replacing a head gasket.

Combustion chambers vary, too, with the more sophisticated head design almost always worth the investment. For example, crowned head chambers create a highly turbulent swirling action to optimize the air-fuel ratio. This adds up to more power from a smaller displacement — and more economical operation.

Intake and exhaust manifold design should be considered, too.

Other features to watch for include industrial-grade bearings, heavy-duty crankshaft, and a mechanically driven fan (turf machine speeds are too slow for ram air cooling common in automotive road speeds).

Vibration and sound

How the engine is mounted will influence performance, too. Isolated mounts, those that separate and cushion the engine from the frame, greatly reduce vibration, which is as important to operator comfort as it is to component life.

An industrial-grade muffler not only makes sense from a standpoint of longer wear and less resistance to exhaust pressures, its quieter operation is less stressful for the operator and better conforms to noise restrictions.


Cursory appraisals of turf equipment have no place in today's economic climate. Functionally, machines may look quite alike. Even performance specifications may be similar.

It's only when the buyer goes well beyond the obvious that the product designed for a decade or more of regular use begins to emerge. The engine should head an evaluation list, for measuring the productivity of the machine begins with its performance.

WT&T

Advances in Turfgrass Pathology

published by **HARCOURT BRACE JOVANOVIĆ PUBLICATIONS** in cooperation with **Dr. B. G. Joyner, Dr. P. O. Larsen and Chemlawn Corporation**



This extensive volume contains chapters on:

- turfgrass diseases
- cool v.s. warm season pythium blight and other related pythium problems
- snow molds of turfgrasses
- fairy rings
- leaf spot of Kentucky Bluegrass in Minnesota
- initial and field fungicide screening
- turfgrass disease resistance
- PLUS MUCH MORE!

\$27.95* (hardcover)

COPIES LIMITED — DON'T DELAY!

ADVANCES IN TURFGRASS PATHOLOGY is a compilation of more than 23 reports and discussions by the nation's leading turfgrass pathologists. Explore the diseases that attack turfgrass. Find out how to conquer the battle of turfgrass diseases.

KEEP CURRENT WITH NEW IDEAS ON HOW TO HANDLE TURFGRASS PROBLEMS WITH ADVANCES IN TURFGRASS PATHOLOGY.

Return this coupon to: Book Sales
Harcourt Brace Jovanovich Publications
One East First Street, Duluth, MN 55802

YES! Please send me _____ copy(ies) of ADVANCES IN TURFGRASS PATHOLOGY.

Name _____
Address _____
City _____ State _____ Zip _____
Phone _____

\$27.95* hardcover
Quantity rates available on request.


A check or money order for _____ is enclosed.

*Please add \$3.00 per order plus \$1.00 per additional copy for postage and handling.

Please charge to my Visa, Master Card, or American Express (circle one)
Account Number _____
Expiration Date _____
Please allow 6-8 weeks for delivery.

WTT 64

New Toro® controllers combine simplicity with total control



Toro's 8+1 and 12+1 electromechanical controllers handle drip irrigation, too!

Here's the latest advance in controllers, solving many of the problems you've reported from the field. For example, we think you'll appreciate that the key to our 8+1 and 12+1 controllers is *simplicity*, yet with all the control you need. They're simple to install, and that saves you time and money! Simple to *program*, too. And simple to *service*. All the advantages you've been looking for. The

" + 1" part of our name? That stands for the one extra, *independent* system that controls drip irrigation *simultaneously* with any of the 8 or 12 stations. As far as we know, these new Toro electromechanical controllers are the only ones of their kind available today. As always, a step ahead of the rest . . . TORO!



The Toro Company, Irrigation Division
Dept. WT-684, P.O. Box 489, Riverside, CA 92502

EXCELLENCE IN IRRIGATION

Circle No. 142 on Reader Inquiry Card

PROBLEM SOLVERS

by Balakrishna Rao, Ph.D., and Thomas P. Mog, Ph.D.

Summer fertilizer safety

Problem: Our lawns are dethatched, limed (50 lbs./1,000 sq. ft.) and fertilized (10-6-4, 50% fertilizer) each spring and fall. The lawns consist mainly of Kentucky bluegrass and creeping and tall fescue with very little ryegrass, and all the clippings are picked up after mowing. Our summer fertilizer is 20-4-10, 40%. If this is applied before July (mandatory) would it be too strong for summer? (New Jersey)

Solution: Not knowing the exact amount (lbs./1,000 sq. ft.) of different fertilizers being used during different times of the year, it is difficult to respond to your question. However, if you are using these fertilizer formulations to provide no more than 1 lb. of actual nitrogen per 1,000 sq. ft. per treatment, then it should be safe enough to use in summer.

Sandburr control in Texas

Problem: What will eliminate sandburrs in the east Texas area between Dallas and Shreveport? (Texas)

Solution: You can use the preemergent materials like diphenamid, trifluralin or EPTC for sandburr weed control on turf and ornamentals in Texas. Although results may vary, you can expect some level of control from the use of these products.

You can expect better results using arsenical materials such as DSMA or MSMA as postemergent herbicides. These arsenical materials, however, may have phototoxic effects and produce temporary discoloration of the treated area. Treat the area when weeds are young and actively growing which would be during late spring and early fall.

Needle drop on Douglas fir

Problem: One of my account's Douglas fir trees are turning color and dropping their needles. What's really strange is that two trees may be right next to each other; one will be healthy and the other appears to be dying. Do you know what is causing this? (New York)

Solution: There are two possibilities; 1) a needle disease and 2) natural shedding of the older needles. So-called "evergreens", like the pines, spruces and firs, shed their foliage just as the broadleaved or deciduous trees do. Most conifers hold their needles for two or more years, after which they are shed.

Needle drop usually occurs in the fall and is a natural process which normally takes several weeks. Sometimes adverse environmental conditions trigger color change and shedding of the older needles at times other than fall. When this happens, the trees' appearance may change from normal to abnormal in a period of days. In both instances the newest needles are still on the tree.

Several different fungi can cause somewhat similar symptoms. The fungi responsible for a diseased condition that results in premature shedding of the foliage are collectively known as needlecast fungi.

With needlecast, one tree may be diseased and a nearby tree of the same species can be free of symptoms. This phenomenon is often attributed to differences in genetic makeup and is called host resistance.

If all firs were showing similar patterns of injury involving only the older needles, then one would suspect natural or environmentally induced shedding of the foliage and not disease.


Wood ashes as fertilizer

Problem: What kind of nutrient value is there in wood ashes from fireplaces? Some of our clients are thinking of using these around trees. I would appreciate your comments in this regard. (New York)

Solution: Recently, there has been increased interest in the use of wood ashes produced from fireplace burning as a fertilizer source. The wood ashes from fireplaces contain about 5% k₂o (potassium) and 25% Ca (Calcium). Many people use these ashes as fertilizer in their gardens. Hardwood ashes are frequently used as fertilizer for tobacco growing. From this information, I would imagine that the ashes from fireplaces could be used as fertilizer to supply the above nutrients.

Fruit or shade, rates differ

Problem: I plan on using Benlate to control scab. The label says use four to six ounces per 100 gallons on apples. For shade trees the rate is one pound per 100



Balakrishna Rao is plant pathologist and Thomas Mog is pest management specialist for Davey Tree Expert Co., Kent, OH.

Questions should be mailed to Problem Solver, Weeds Trees & Turf, 7500 Old Oak Boulevard, Cleveland, Ohio 44130. Please allow 2-3 months for an answer to appear in the magazine.

Now ~~All~~ ~~8~~ of the top 10 golf courses in America have Toro[®] irrigation



There are some good reasons WHY!

Toro irrigates all of America's top 10 golf courses...17 of the top 20 ...34 of the top 50. That's because irrigation probably is the most important single factor in turfgrass management, and most of those responsible realize that Toro offers them a hard-to-beat combination of expert knowledge and advanced-design equipment. Whether you're planning a new course, replacing your existing system, or trying to solve specific problem areas, a good partner to team up with is the man from Toro!



Toro's VT3 central-satellite control system utilizes solid state electronics for accuracy and dependability. And now, ask about Toro's exciting new VT3 Video Central Controller, utilizing an IBM personal computer, easily operated with a light pen!

TORO[®]

Excellence in Irrigation™

The Toro Company, Irrigation Division
Dept. WT-684, P.O. Box 489, Riverside, CA 92502

Circle No. 143 on Reader Inquiry Card

gallons. The apple rate would be cheaper. Why the big difference? (Pennsylvania)

Solution: The "apple rate" is for apples as a food crop. Tolerance levels have been set for pesticide residues on food crops. At the shade tree rate the fungicide residue on apples might exceed permissible levels. Finally, most commercial fruit growers spray their trees weekly or bi-weekly thus reducing the need for a long-lasting residual.

Satisfactory control of scab on shade trees can be achieved with two or three applications of Benlate at the rate of one pound per 100 gallons. Follow the label instructions as to timing and possible use of wetting agents.

Multi-use spray tanks

Problem: If washed, can a spray tank which is used for weed control be used for fungicide spraying of trees and shrubs? (New Jersey)

Solution: It is difficult to remove some herbicides after they have been used in a sprayer. This is particularly true of the phenoxy herbicides that are used for broadleaf weed control.

The following suggestions are from the Ohio Cooperative Extension Service publication, Chemical Weed Control In Commercial Nursery and Landscape Plantings.

1. 2,4D - It is difficult to remove 2,4D compounds from sprayers. It is advised that sprayers used for

applying these materials NOT be used for any purpose other than applying herbicides.

The 2,4D type materials can be removed if the following recommendations are followed immediately after use (as soon as spraying is completed).

a. Flush out the entire system with water detergent solution immediately after the solution (1 gallon of household ammonia in 10 gallons of water) and let stand for 12 to 24 hours. Disassemble the nozzles and soak the caps, screen, etc. in the ammonia solution.

b. Rinse thoroughly with water and let circulate through the sprayer.

c. Test spray a few plants which you know to be susceptible four to five days before using sprayer on larger areas.

2. PRINCEP - Rinse thoroughly with a detergent immediately after use. Check screens for clogging and, if present, soak in ammonia solution for 12 to 24 hours.

3. OTHER HERBICIDES - Any sprayer used to apply herbicides should be cleaned immediately after use. Special instructions for particularly troublesome herbicides have been previously outlined. If the procedures for cleaning out other herbicides are not specifically outlined,

a. Flush the sprayer system with clean water.

b. Rinse thoroughly with a detergent solution.

c. Rinse again with clear water.

The herbicide label may carry cleaning instructions. If so, follow the manufacturer's recommendation.

A Warren's Turf PRO report...

"We can't compromise...Comiskey class professional operation every

The turf at Comiskey Park is Warren's Champion™, an A-34 BenSun® custom blend of Kentucky bluegrass. Warren's A-34 BenSun® sod or seed is now used in many of the nation's leading stadiums.



Jim Walsh

Roger Bossard

EVENTS

JUNE

Harvard University Design Workshop, apply June 22 through August 16. Harvard University, Cambridge, MA. Summer advanced continuing education in the Graduate School of Design. Topics include Advanced Farden Design, Microcomputers in Landscape Architecture, Street Trees, Fountains and Pools, and Starting and Maintaining Your Own Firm. Tuition ranges from \$400 to \$1,150 and housing is available. Contact Continuing Education, 48 Quincy St., Cambridge, MA 02138 or call 617-495-9340.

Athletic Field Maintenance Seminar and Field Day, June 28. From 9:00 a.m. to 4:00 p.m. at Terry Park in Fort Myers, FL. Sponsored by the University of Florida Institute of Food and Agricultural Sciences. Contact Charlie Lowery, Court Plaza, D-103, 2663 Airport Rd. South, Naples, FL 33962.

JULY

American Association of Nurserymen Annual Conference, July 14-18. San

Antonio, TX. The 109th Annual Conference will be addressed by Lady Bird Johnson. Includes meetings of National Landscape Association, Garden Centers of America, and Wholesale Nursery Growers. Contact AAN, 1250 I St., NW, Suite 500, Washington, D.C., 20005. (202) 789-2900.

Aquatic Plant Management Society Annual Meeting, July 15-18. Richmond Hyatt House, Richmond, VA. Contact William Rushing, P.O. Box 16, Vicksburg, MS 39180. (601) 634-3542.

American Sod Producers Association Summer Convention and Field Days, July 24-26. Olympia Spa and Resort, Oconomowoc, WI. Includes outdoor equipment demonstration. Contact Doug Fender, ASPA, 4415 W. Harrison St., Hillside, IL 60162. (312) 449-2844.

AUGUST

Landscape Technology Courses, Cal Poly University, August 14-16. Pomona campus offers three programs; Arboriculture/Urban Forestry, Nursery Operations, and Dry

Climate Landscaping. Contact Wayne Smith, Cal Poly, Pomona, (714) 835-5550.

International Society of Arboriculture Conference, Quebec City, August 18-23. The 60th annual conference of ISA will be held at the Le Chateau Frontenac, Quebec City, Quebec, Canada. There will be educational sessions and commercial exhibits for all types of arborists. Contact Cal Bundy, ISA, P.O. Box 71, Urbana, IL 61801. (217) 328-2032.

Wisconsin Turfgrass Association Field Day, August 27. Oconomowoc Country Club, Oconomowoc, WI. Complete, "hands-on" display of outdoor power equipment. Contact Ed Devinger, Reinders Brothers, Inc., P.O. Box 57, Elm Grove, WI 53122. (414) 786-3300.

To insure that your event is included, please forward it, 90 days in advance, to: WEEDS TREES & TURF Events, 7500 Old Oak Boulevard, Cleveland, OH 44130.

Park has to be a first day of the season."

"Sometimes when we need sod, we need it fast... and there's no margin for error in delivery or quality. A good example was the night the White Sox clinched the division title last year. Fans flooded onto the field looking for victory 'souvenirs.' Many of them took home huge chunks of our outfield turf.

"With another home game just three days away, we needed fast service if the outfield was going to be playable by game time.

"A call to Jim Walsh at Warren's Turf Chicago operation had Jim and his people out before dawn cutting hundreds

of yards of new replacement sod for us to put down early the next day.

"The sod we use at Comiskey Park has to stand up under heavy traffic, make maximum use of the fertilizers we put down, be aggressive in its growth so that it chokes out weeds, take up to 65% shade and be mowed to a half an inch on the infield. That's a tall order for any turf.

"Over the last ten years, we've tried dozens of turf and seed formulations. We have never found a bluegrass that comes anywhere close to Warren's A-34 for maximum root strength and best blade resilience."

"Our management demands top performance from all of us...and our suppliers have to be just as professional as we are." Roger Bossard, Head Groundskeeper for Comiskey Park—home of the Chicago White Sox—reports on the backup he gets from Jim Walsh and his team of Warren's Turf Professionals.

"Wherever you put down roots, go with the PROs."

Warren's

TURF PROFESSIONALS

7502 South Main Street • Crystal Lake, Illinois 60014


Warren's offers only professional turf products.



Circle No. 146 on Reader Inquiry Card

Call (800) 828-TURF (8873) for the name and address of your nearest Warren's Turf Professional or Warren's TerraBond Dealer.

**GET ON TOP
WITH THE
BEST.**



PROMARK AERIAL LIFT

**AVAILABLE IN
FOUR MODELS**

TOP OF THE LINE FEATURES WITH BOTTOM LINE ECONOMY.

For a quality-engineered aerial lift, including a wide range of features at a reasonable price, the PROMARK is unique. If you are buying a lift, research comparison will convince you that the PROMARK is truly in a class by itself.

COUNT THESE OPERATIONAL FEATURES:

- Continuous 360° rotation.
- Further side reach.
- Hydraulic outriggers.
- Quick-mount installation.
- Yoke-mounted, self-leveling bucket.
- Ground level entry.
- Easy lift removal.
- Forward lift mounting—retains large load capacity.
- Smooth hydraulic controls.

WRITE OR PHONE FOR MORE INFORMATION.



PROMARK PRODUCTS, INC.

306 9th Avenue
City of Industry, CA 91746
(818) 961-9783

Circle No. 131 on Reader Inquiry Card

LETTERS

Rose kills in Missouri

I was very interested in Michael Dirr's April article entitled Winter Damage.

In my area of southeast Missouri (zone 6) people are now observing widespread rose kills and plant damage. Although 1983 began as a wet spring, the rains stopped in May and did not resume until October. We apparently received adequate fall rains, but our roses did not face any cold weather until mid-December when temperatures dropped drastically to about -16° for a couple of weeks with wind chill factors of -60° occasionally.

I believe that my area of Missouri does not get cold enough in the fall to harden plants properly for the cold arctic blasts that come in December and January. However, since we get frequent fall and winter rains instead of snow, should roses be protected with heavy mulches of straw and sawdust which might be damaging by holding too much moisture? Would it be better to cut roses back about two-thirds and cover them when conditions moderate?

Also, I have observed several roses which are alive at the crown that were on their own roots. I am not sure if this means anything or not since I need more observations.

Larry Hanning

Area Entomologist
State of Missouri Agriculture
Department, RR1, Box 312,
Burfordville, MO 63739

Thank you for sharing your observations. We ask other readers to share their observations about winter kill as you have. Your letter is being forwarded to Michael Dirr for a reply.

Exotic or native, adaptable trees needed

Douglas Chapman's article on street trees in the April issue was informative, as always. I agree that an effort should be made to find adaptable species for urban growing conditions, but I think Chapman puts too much emphasis on the use of natives.

Many desirable native trees are unsuitable in street tree planting programs. In Missouri, natives accustomed to poor growing conditions such as post oak, hickories, sassafras and others are commercially available.

Almost half the trees mentioned in

E-Z-GO'S HAUL OF FAME.



Our gas powered GX-800, and our new electric X-500 light turf vehicles are perfectly suited for tough jobs when you need a utility vehicle but a truck is too much.

The GX-800 sips gas like a miser, and won't mar your manicured turf. And its hill climbing ability and sure handling in bad weather make it a step up from any pick-up truck.

Likewise, the new electric X-500 will give you a full day's work on one night's charge. And it too is perfect when you need a vehicle as strong as an ox yet as gentle as a lamb.

For heavier work, the GT-7 has you covered. With a large payload capacity and steel-strong durability to match any heavy-duty turf truck.

Plus, it's infinitely adaptable to the job at hand. A hydraulic dump, aerator, spreader, sprayer, top dresser, and many other options can be easily added to make your particular task less of a chore.

The gas powered GX-800, GT-7, and the new electric X-500. They're tough, strong and best of all they're from E-Z-GO. Which means there's thirty years of experience engineered into each one. The World's Finest Utility Vehicles For Sports and Industry. **E-Z-GO** **TEXTRON**

P.O. Box 388, Marvin-Griffin Road, Augusta, Georgia 30913-2699, (404) 798-4311

Circle No. 254 on Reader Inquiry Card

Chapman's list of species for planting on adverse sites are imports. None of these should be rejected just because they are exotics. Using the criteria of native vs. non-native to select trees needlessly narrows the number of choices and restricts diversity.

What difference does it make if ninety percent of the trees in an urban area are exotic if they are the best trees available for the job?

Tim Frevert
Landscape architect
Missouri Department of

Conservation, P.O. Box 180, Jefferson City, MO 65102-0180

University of Georgia swims upstream

I agree with Dave Pinkus in general (Trends, April, Colleges Care More About Grants Than Students), but some of us are swimming upstream. I enclose our program for the Ornamental Horticulture program at the

University of Georgia, Athens. It has a good smattering of management and economics. If you know of some firms who want to hire interns or graduates of our program, please let me know.

Jake Tinga
Professor of Ornamental Horticulture
University of Georgia, College of Agriculture, Athens, GA 30602

Thanks for the memories

The Man of the Year trophy adorns my living room and is greatly admired. So far, we haven't filled it with suds or bubbly.

Thank you for giving me the exposure in select company—it was a thrill.

Fred V. Grau
President
The Musser International Turfgrass Foundation
College Park, MD

Fred V. Grau was presented the third annual Man of the Year trophy during a meeting of the Board of Directors of the Musser International Turfgrass Foundation at the GCSAA Show in Las Vegas. The Board consists of hand-picked contributors to the turf industry. Editor.

Forsythia poor indicator of crabgrass germ

I have just read Landscape Log in February *Weeds Trees & Turf*. A common mistake made and that has been circulated is the idea that crabgrass germinates around Forsythia bloom. I have been watching this occurrence for the last three years in the Salt Lake City area and have noticed that crabgrass typically is much later than Forsythia bloom.

Forsythia can bloom four to six weeks before crabgrass germination. Crabgrass germination has been so regular in our area that it seems like it is tied into photoperiodic response in some way. Germination in our area has been around May 23 each year for the last three years in heavy as well as light soils!

If I were to tie it into some sort of landscape phenology, I would probably use Radiant crabapple bloom, apple or pear bloom.

In our area, I've suggested that pre-emergents should be applied no later than the first week of May.

Robert L. Morris
Ornamental Horticulture Specialist
Utah State University
Salt Lake City, UT



They do. It's their job to know things first and then pass that information on to you fast. Things like new turf management techniques, effective methods of insect and weed control, what really works... and what won't.

LOOK FAMILIAR?

They should. You've seen them often at turf shows, seminars, conventions, wherever industry news is in the making. They're the editorial and sales management team of **WEEDS TREES & TURF**. Please meet (seated l. to r.) Ron Kempner, Bruce Shank, Dick Gore, and Maureen Hrehocik; (back row l. to r.) Bob Mierow, Kevin Cooney, Joe Kosempa, and Bob Earley.

THERE'S NO GRASS GROWING UNDER THEIR FEET.

They apologize for not always being in when you call, but great stories are found in the field, not in the office. These pros know the only way to really cover the green industry is to get out and be a part of it. From turf test plots in Oregon to the corridors of our Nation's Capital, they're following leads, surveying markets and interviewing the

landscape professionals who make our industry hum. Then, they use their years of experience to pull it all together, analyze it and present it to you in a crisp, easy-to-read and easy-to-use style. That way, you have the information you need to do your job better, faster and more effectively.

The next time you want to know something in the green industry, give them a call. If you can't reach them at the office, don't worry. They'll reach you in the pages of **WEEDS TREES & TURF**.

WEEDS TREES & TURF ®BPA
®VII

HRJ A HARCOURT BRACE JOVANOVIICH PUBLICATION

3091 Maple Drive, Suite 312, Atlanta, GA 30305
(404) 233-1817

— A GREAT MEDIUM TO GROW PROFITS IN. —

Subdue. The most effective fungicide against Pythium blight and damping-off.

Pythium weather. High temperatures, high humidity and high anxiety. Once Pythium takes root, it can destroy turf within hours.

Unless you take a grass-roots approach to Pythium. With Subdue.*

Subdue works both on contact and systemically.

Subdue fights Pythium blight and damping-off—as well as downy mildew (yellow tuft)—in two ways. On contact, Subdue destroys

the fungi in the soil. Systemically, Subdue prevents disease from within grass plants. That's because Subdue is water soluble—easily absorbed by roots. So Pythium—and now, downy mildew—don't have a chance.

Subdue also controls costs.

Subdue's systemic action means longer, more effective residual

protection. Fewer applications. Lower chemical costs. And savings in maintenance and labor. And Subdue's low application rate—1 to 2 fluid oz. per 1,000 sq. ft. for 10 to 21 days on established turf—makes Subdue the most cost-efficient protection you can buy.

Before Pythium weather strikes, subdue it. Use Subdue in a preventive maintenance control program. And get a good night's sleep.

Ciba-Geigy, Ag Division, Box 18300, Greensboro, NC 27419.

CIBA-GEIGY

© 1983 Ciba-Geigy Corporation



**HOW TO AVOID SLEEPLESS NIGHTS
DURING PYTHIUM WEATHER.**

SUBDUE

Circle No. 274 on Reader Inquiry Card

Scientific Guide To Pest Control Operations

by Dr. L.C. Truman
Dr. G.W. Bennett and
Dr. W.L. Butts



Domestic: \$32.50* (hardcover)
Foreign \$37.50* (hardcover)

The SCIENTIFIC GUIDE TO PEST CONTROL OPERATIONS is designed to provide a sound basis for studying the scientific aspects of pest control and promote technical competence. It places emphasis on urban and industrial pest problems. This volume also covers the laws and regulations concerning the pest control industry.

The SCIENTIFIC GUIDE TO PEST CONTROL OPERATIONS is written for owners, supervisors, servicemen, salesmen, students, persons preparing for state certification under the EPA/state programs for commercial pesticide applicators, and people interested in structural pest control.

The SCIENTIFIC GUIDE TO PEST CONTROL OPERATIONS - a *must* for anyone involved in the field of pest control!

SURVEY from page 16

in Ontario, has generated some of the first data on turf expenditures in the Province. The data show the value of producing and maintaining turfgrass in Ontario ranks third considering all agricultural products, after corn and tobacco.

The value of expenditures on turfgrass maintenance by all user groups was conservatively estimated at \$275 million, with \$75 million spent on turf equipment and \$26 million on pesticides, sod, and seed. Commercial lawn service sales were pegged at \$45 million, although a few lawn care operators told *Weeds Trees and Turf* this figure is low.

Golf courses spend more on maintaining turf than any other user group, followed by residential lawn care and sod farms.

Government turf managers surveyed indicated extension specialists could not help them with their most serious problems of labor and equipment shortages. Private turf managers were more supportive of extension services in problem solving.

Copies of the survey results are available from the Ontario Turfgrass Research Foundation, 54 Hershaw Crescent, Etobicoke, Ontario, Canada, M9C 3M4.

Return this coupon to: Book Sales

Harcourt Brace Jovanovich Publications
One East First Street, Duluth, MN 55802

Domestic: \$32.50* (hardcover)
Foreign \$37.50* (hardcover)

YES! Please send me _____ copy(ies) of the SCIENTIFIC GUIDE TO PEST CONTROL OPERATIONS.

A check or money order for _____ is enclosed.

Please charge to my Visa, Master Card, or American Express (circle one)

Account Number _____ Expiration Date _____

*Please add \$3.00 per order plus \$1.00 per additional copy for postage and handling.

Name _____

Address _____

City _____ State _____ Zip _____

Signature _____ Date _____

Phone _____

WTT 64

Quantity rates available upon request.

Please allow 6-3 weeks for delivery

TURF MANAGERS' HANDBOOK

By Dr. William Daniel and
Dr. Ray Freeborg

\$23.95* **hardcover** \$18.95* **paperback**

This essential reference book covers:

- scope and organization
- grasses and grooming
- rootzones and water
- nutrition
- pests and controls
- uses of turf
- serving turf needs
- AND MORE

The TURF MANAGERS' HANDBOOK is an easy on-the-job reference to planning, purchasing, hiring, construction, and plant selection. These 424 pages contain 150 illustrations, 96 color photographs plus 240 tables and forms.



ORDER YOUR COPY NOW!!!

Ordering Information

Please send _____ copies of the hardback (\$23.95* ea.)

_____ copies of the paperback (\$18.95* ea.)

Quantity rates available upon request.

*Please add \$3.00 per order plus \$1.00 per additional copy for postage and handling.

Please charge to my Visa, Master Card, or American Express (circle one)

Account Number _____

Expiration Date _____

Please allow 6-8 weeks for delivery.

Signature _____

Name (print) _____

Address _____

City _____ State _____ Zip _____

Phone _____

Send to: Book Sales

Harcourt Brace Jovanovich Publications

One East First Street

Duluth, MN 55802

WTT 64

GOLF

China plans prime golf resort complex

Construction is expected to begin by the end of 1984 on a major golf resort complex in the city of Guilin, People's Republic of China.

The Ronald Fream Design Group, golf course architects in Santa Rosa, CA, has been commissioned to prepare the master plan. The Fream Group, along with Hong Kong-based architects and engineers, will prepare the plans for three eventual sites in the scenic area.

The Beijing Government intends to develop the Guilin region with international standard tourists facilities, of which the golf resorts will be a part.

The initial site will include 18 holes of championship calibre golf, golf school practice area, and clubhouse. A 60-room luxury lodge and an additional 18 holes are planned for the site, which has exotic limestone pinnacles, rolling land, and pine trees. A second site near the Lijiang River gorge will have 36 holes of international standard competition golf, a teaching and practice school, and several other recreational facilities.



BEFORE YOU BUY ANYTHING ELSE, CHECK THE IMPORTANT BENEFITS OF LESCO TURF-TYPE TALL FESCUES.

- Drought Tolerance
- Disease Resistance
- Year-Round Color
- Low Growth Habit
- Shade Performance
- Versatility
- Darker Green Color
- Increased Density



Shaded area indicates where Olympic, Falcon and Apache are adapted.

Olympic, Falcon and Apache. Three of the best turf-type tall fescues. All available from LESCO. Perfect for home lawns, sod production, athletic fields, commercial turf and golf courses. The benefits are tough to ignore.

Quantities of turf-type tall fescues may be limited.
Call Barb today to place your fall order.

(800) 321-5325

(800) 362-7413

Nationwide

In Ohio

LESCO



Lesco, Inc., 20005 Lake Road, Rocky River, Ohio 44116, (216) 333-9250

Circle No. 121 on Reader Inquiry Card



CLASSIFIEDS

RATES: 95 cents per word (minimum charge, \$20). Bold face words or words in all capital letters charged at \$1.20 per word. Boxed or display ads charged at 1x—\$80.00, 3x—\$78.00, 6x—\$75.00, 12x—\$70.00 (one inch minimum). Agency commissions will be given only when camera-ready art is provided by agency. For ads using blind box number, add \$5 to total cost of ad. Send ad copy with payment to Dawn Anderson, WEEDS TREES & TURF, 1 East First Street, Duluth, MN 55802.

BOX NUMBER REPLIES: Mail box number replies to: WEEDS TREES & TURF, Classified Ad Department, 120 W. 2nd St., Duluth, MN 55802. Please include box number in address.

BUSINESS OPPORTUNITIES

WANT TO BUY OR SELL a golf course? Exclusively golf course transactions and appraisals. Ask for our catalog, McKay Golf and Country Club Properties, 15553 N. East Street, Lansing, Michigan 48906. Phone (517) 484-7726.

TF

Wanted: Distributors for aluminum edging 1/8 x 4 and 3/16 x 4 sizes. Excellent product. Mid State Distributors, 5700 Kraft SE, Grand Rapids, MI 49508, (616) 698-7350.

6/84

FOR SALE

LAWN SEED. Wholesale. Full line of top quality grasses. Improved bluegrass varieties, fine fescues and fine bladed ryegrasses. We specialize in custom mixing. Oliger Seed Company, 2705 Wingate Avenue, Akron, OH 44314. Call collect (216) 753-2259.

TF

TREE FERTILIZATION GUN, tested on over a million square feet of trees and shrubs all over the U.S. Good to 500 psi, repairable, non-corrosive. Buy direct from manufacturer, \$92.50. Arbor-Nomics, Inc., 5634-A Buford Highway, Atlanta, Georgia 30071. (404) 447-6037.

TF

SKYWORKER AERIAL BUCKETS — Sales of both new and used. Finest service anywhere in the country. Overhauls and changeovers are our specialty. Best prices on parts. We deal in all types of tree trimming equipment. Ask about fleet discounts. Call or write American Hydraulics, Inc., Route #4, Hartwell, GA 30643. (404) 376-3192.

7/84

SALE--SALE--SALE--Hannay 12 volt base mount reel motors, \$100.00 (new). Regular \$160.00. FMC (John Bean) parts and pumps at discounted prices. **Hydro pumps** parts and accessories at 25% discount. Call: Strong Ent., Inc., Collect: (305) 264-5525.

TF

MEYER ZOYSIA — Pure Meyer Z-52 zoysia sod, big savings on water, bug killers, mowing and fungicide, great service. **BEAUTY LAWN ZOYSIA** (Cincinnati) (513) 424-2052.

11/84

HYDROSEEDERS - MULCH SPREADERS — When looking for a good quality used Hydro-Seeder or Mulch Spreader at an affordable price, call Finn Corporation and ask about our factory reconditioned and demonstrator models. Toll free 800/543-7166. In Ohio 513/871-2529.

6/84

200 gallon Century Skidmount Sprayer 5200 Series, Hydro twin piston pump. Mechanical agitation, 200 foot hose, Chemlawn spray gun. (814) 838-1736.

6/84

FOR SALE — Replacement parts that will fit **BOBCAT, BUNTON** walk-behind mowers' main drive belt, list \$25.00 our price \$19.50. **DUMP CATCHER**, basket only \$125.00. Shipping only \$3.00 on any size order. Dealers wanted. Call or write for catalog. Precision Power Equipment, Inc., 2400 Boston Rd., Wilbraham, MA 01095, (413) 596-2996.

7/84

FOR SALE — Established sod farm in beautiful Western Colorado. 170 acres of sod plus equipment and 4 homes. Call (303) 858-7464 or write to Jim at 1268-18 1/2 Road, Fruita, CO 81521. 7/84

For Sale: One 10-1/2 ft. Methyl Bromide Fumigator, complete with controls and tanks. And one tarp or plastic puller. Both \$4500. Call (904) 732-3014.

TF

BALL BARRIER NETTING: Made of olefin fibers. 6-1/2 feet and 25 feet high. Strong and tough. Will not rust. Easy to handle. For Driving Ranges and Golf Courses. Keep golf balls from straying off-course. J.A. Cissel Mfg. Co., P.O. Box 339, Farmingdale, NJ 07727. (800) 631-2234.

6/84

USED EQUIPMENT

HYDRO-MULCHERS AND STRAW BLOWERS New and used. **JAMES LINCOLN CORPORATION**, 3220 S. Jupiter Rd., Garland, TX 75041. (214)840-2440 (TX), (800)527-2304(except TX) TF

HI-RANGERS AERIAL BASKETS 65', 57', and 53'. Skyworkers aerial baskets 65', 50', 40'. Vermeer stump cutter 1560.6. Vermeer tree spade 66, TS 44. Asplundh bucket and brush chippers, Bean sprayer, 9 ton trailer. Parkway Tree Service, 12026 W. Cherry, Wauwatosa, Wisc. 53226. (414) 257-1555.

TF

For Sale — 1978 Princeton Sod Harvester, Model #4020, with John Deere diesel. Ready for work! Looks good — Runs Good! \$17,500. Grass Farm, Morgan Hill, CA. (408) 226-9775.

TF

For Sale: Two Bean 1010 Sprayers, two hose reels, 150 feet of hose each. One unit \$2,800.00, one unit \$2,000.00. Excellent condition. Call Bruce, 218-367-2033.

6/84

NEW and USED EQUIPMENT — Asplundh, Hi Ranger and Lift-all forestry bucket trucks, Chipmoo wood chippers. Mirk, Inc., (216) 669-3567, (216) 669-3562, 7629 Chippewa Road, Orrville, Ohio 44667.

TF

1977 **TORO PARKMASTER** nine gang with Perkins Diesel. Low Hours - Excellent condition, \$17,900 or best offer. Call 203-966-2145 days or 203-966-0223 evenings.

6/84

TREE SPRAYER for sale. Bean Rotomist 100E. 100 gal., 27,000 c.f.m., excellent condition; trailer mounted. \$4,500. Kaiser Tree Preservation Co., 401-294-6397.

7/84

1980 **MACK TANDEM**, Model R686 — 316 H.P. 6 speed Maxidyne Transmission. 12000 front, 38000 rear. Power steering. Air Dryer. National cushionair seat. Used only one season — 51,000 mi. Heated barn storage. Absolutely new condition. \$33,500.00 U.S. Near Toronto (519) 893-1350.

7/84

1973 **BROUWER SOD HARVESTER**. Only 3600 hrs. Sod operation closed out. One owner. Excellent Condition. Parts & inventory & manual included. \$12,500.00 U.S. or best offer. Near Toronto (519) 893-1350.

7/84

Need **storage** for excess chemicals, fertilizer, equipment? **Used ocean cargo containers reconditioned and repainted** to an attractive state. Saves you money on construction costs and insurance rates, keeps you from relocating or trying to construct a building on leased property. Store your chemicals or flammables in a **safe, secure, watertight environment**. Steel frame, hardwood floors. 40' L x 8' W x 8-1/2' H, \$2,490 plus shipping. 20' L, \$1,990 plus shipping. (317) 873-5382 George. AS, P.O. Box 373, Zionsville, IN 46077.

7/84

FOR SALE BY OWNER: Princeton Piggyback Forklift, with Diesel Eng., Excellent Cond. Asking only \$12,000. George Wilcox Contracting Co., 501-945-5005 or 501-275-3518.

6/84

6' Lely tiller with seed box, 950 John Deere Turf Tractor with Loader, (WANT TO BUY — used 48' - 57' Asplundh Bucket Truck with chipper box and used brush chipper). 701-224-0100.

6/84

FOR SALE: 1983 Harley Rock Picker, 12' Rock Windrower in Phoenix, AZ. \$19,500 both units. Gundersons, Inc., Rapid City, SD, (605) 342-4610.

6/84

Large All Terrain Tree Mover. Terex 72-41 3 yd. Wheel Loader with Vermeer TS-66 Hydraulic Tree Spade. Quick Coupler Hydraulics, Bucket Included. Low Hour Machine with Cab, Heater, Lights and Wipers. **\$26,500**. Also Have Vermeer TS-44 on 4x4 Case 119 H.P. (Model 1200) Farm Tractor with 4 Wheel Steering. Excellent All Terrain Unit. Also Vermeer TS-30 Tree Mover on Vermeer M-470 4x4 Tractor with BH-700 Backhoe on Front and 6 Way Hydraulic Dozer. **\$10,000**. **Adam Fritz Company, Inc.**, P.O. Box 578, 33 Old McHenry Road, Lake Zurich, Ill. 60047. **Call Anytime 312-438-5101**.

6/84

1980 Hydro 180 3 gang spindle hydraulic mower — A1 — will trade. Meyers Turf Farms, Stilwell, Kansas. 913-681-2781.

6/84

Never used 1000 gallon trailer mounted Super Seeder (hydromulcher) with hose reel, excellent condition. Cost over \$10,000.00, will sacrifice for \$8,000.00. 317-447-9204.

7/84

Brouwer A3A Harvester 1981 slab machine on Ford. 254 hours. Like new. \$24,000. Bob Lassen, Grand Island, NE. 308-382-6532.

6/84

Toro Park Master lawn mower. 7 gang — 30' rough mowers with hydraulic lift. Gas motor rebuilt one year ago. High flotation rubber. Very good condition. (519) 846-5874. Elora, Ontario.

For Sale — Used Reinco Hydroseeder. 1000 gal. Good condition. Will sell or trade for straw blower. L.A.S. Landscaping, Pulaski, VA, (703) 980-7539.

6/84

**CLASSIFIEDS
GET RESULTS**

HELP WANTED

Tree and shrub care. Branch Managers and Sales Managers. America's fastest growing tree and shrub company seeking experienced, aggressive personnel for existing and expansion markets. Experience, degree and motivation key ingredients. Tremendous advancement opportunities. Excellent salary, benefits. All inquiries will be strictly confidential. Send resume, salary history to: **William Scott Carr, Tru Green Tree and Shrub Care, 2875 Northwind Drive, Suite 205, East Lansing, MI 48823.** Expand your potential with an industry leader. 6/84

Pest Management Manager. Aggressive Chicago area tree service has an opening for someone to take charge and manage its chemical applications program. Must be strong in entomology and pathology in both education and experience. Year 'round employment possible with full benefits. Supervisory experience helpful. Send resume and salary requirements to: WTT Box 341. E.E.O. 6/84

HELP WANTED — Top-rated resort/real estate property seeks knowledgeable retailer to manage nursery center. Degree in horticulture or comparable practical experience and supervisory skills required. Sea Island Company, located on Georgia coast between Jacksonville and Savannah, offers excellent salary plus extensive, non-contributory benefits including employee and dependent health/dental/life group insurance, sick and disability benefits, paid vacations/holidays and much more. Send resume in confidence to: Personnel, P.O. Box 1027, Sea Island, GA 31561. (912) 638-3611, ext. 191. 6/84

Arborist — Growing Midwestern tree service has several positions open: Crew leader/Foreman for private tree care: Must be experienced and proficient with rope and saddle and dealing with people. Good advancement opportunity. Arborist trainee: Must have horticulture, arboriculture, or Forestry training or experience. Possible advancement to Sales positions for aggressive self starters. Year 'round employment with full benefits. Please send resume and salary experience to: WTT Box 342. E.E.O. 6/84

WANTED

WANTED: Large Lindig and Royer Shredders. Lewis Equipment, 320 Third Street S.W., Winter Haven, FL 33880. (813)294-5893. 6/84

Wanted to buy HYDRO SEEDER. Please send photo. Johnson Hydro Seeding Corp., 13751 Travilah Road, Rockville, MD 20850, 301/340-0805. TF

Wanted: Large Royer-lindig-W&W-Shredders, or Large Hammer Mill with capacity 75-100 Y.P.H. 405-842-7177. 6/84

POSITION WANTED

Chemical Lawn and Tree Care Manager — Seeks a challenging career position for the same, or related industry. Seven years experience. Resume upon request. Write WTT Box 343. 6/84

Use Our Classified Blind Box Service for Confidential Results!

MISCELLANEOUS

DIESEL HI-RANGER TOWERS—48-100 Feet working heights. Daily, weekly, monthly rentals with or without operator. **MATLOCK LEASING,** Pottstown, PA (215) 326-7711 or (800) 345-7711. 12/84

KELWAY® SOIL ACIDITY TESTER, used by **PROFESSIONALS** nationwide. Direct reading, portable, serviceable. Model HB-2 reads moisture, too. Available from distributors. Brochure from **KEL INSTRUMENTS CO., INC.,** P.O. Box 1869, Clifton, N.J. 07015, (201) 471-3954. 10/84

ADVERTISER INDEX

NO.	ADVERTISER	PAGE	NO.	ADVERTISER	PAGE	NO.	ADVERTISER	PAGE
101	Barebo	26	122	Lofts Seed	CV 4	131	Pro Mark Products	76
102	Beauty Lawn		149	Lofts Seed	42A-42F	132	Rhone-Poulenc	41
	Zoysia	43	123	Mathews Equipment		133	Royal Coach/ Buckner	CV 3
103	E.F. Burlingham & Sons	63	124	J.J. Mauget Co.	65	134	SDS Biotech/Ag Chem Business	30,31
104	Chevron Chemical Co.	45		Mitsubishi (regional)	27	135	Solo	40
274	Ciba-Geigy Corp.	79	126	Mobay Chemical	36,37	136	Source Technology Biologicals	33
106	Ciba-Geigy Corp.	42		Mobay Chemical (regional)	15	137	Tee-2-Green	9
107	Dedoes Industries	28		Mobay Chemical (regional)	69	138	Tennant Co.	43
108	John Deere & Co.	4,5	129	Monsanto Co.	10,11	139	Terracare Products	50
109	Dow Chemical U.S.A.	CV2,p.1	130	Nor-Am Chemical Co.	7		Terrain Vehicle (regional)	27
110	Estech	34				141	Toro Co.	21
254	E-Z-Go/Div. of Textron	77				142	Toro Irrigation Div.	71
112	Finn Corp.	61				143	Toro Irrigation	73
113	Full Circle	55				144	Trebor Corp.	51
114	Glenmac	50				145	Turf-Seed	53
115	Holland Hitch Western Ltd.	51				146	Warren's Turf	74,75
116	Int'l Seeds	13					Whitmire Research Laboratories (regional)	78
253	Irri-Trol Mfg.	3				148	Woods/Div. of Hesston	35
118	Jacklin Seed Co.	49						
119	Jacklin Seed Co.	57						
120	Lesco	29						
121	Lesco	81						



**Nursery
Marketing
Council**
 Using the "good times"
 to build buying habits
 consumers will remember
 if "bad times" come

Thirsting for answers

There is a very real threat to the Green Industry looming on the horizon.

It's water. Not only lack of it, but quality as well. Everyone seems to know a problem exists, but somehow the threat hasn't been real enough yet and the bite in the pocketbook hasn't been big enough to build any kind of a groundswell of concern.



When the editors of WT&T decided to do a story on this critical issue, we didn't realize just how wide the floodgates were that we were opening.

For two and a half months I became immersed (sorry, no pun intended) in the subject of water. With every phone call I made, with every interview I did, with every bit of water information that flooded my "in" basket, this germ of a story idea had a rippling effect -- growing wider and wider with each new piece of information.

It was amazing that a resource so common and so taken for granted has such a profound effect on so many parts of our lives and livelihoods. From golf course superintendent to landscape architect to city water department official, from California to Florida, the sophistication level of knowledge and understanding of this very imminent threat to the health and well-being of an industry was indeed impressive.

What's being done industry-wide to ebb the tide of a shrinking resource is even more impressive. The Green Industry has very few ostriches with their heads buried in the sand waiting for the problem to go away. There is an awareness that while the U.S. essentially has enough water right now, that picture could change rapidly and drastically with continued misuse and fickle Mother Nature.

Next month and in August, a good many of the pages of this magazine will be devoted to a two-part series on water use, conservation, quality and quantity as it effects the Green Industry.

Part I will take a particularly hard look at problems in Florida, California and Texas because of the types of water problems those states contend with and the quantity of water they need to survive. In Part II in August, we will concentrate on solutions to some water problems through low water use turfgrass and ornamentals research, industry association involvement, and how the pivotal industry -- irrigation -- is responding to the increasing challenges.

The picture, surprisingly, while serious, is not bleak. The water situation in this country, does, however, demand continued and diligent attention by an industry who will feel any tightening of control of water the hardest.

Somewhere in this upcoming series you will see yourself and your industry. Hopefully, it will be as part of the solution, not the problem.

Maureen Hrehocik, managing editor

WT&T Editorial Advisory Board



Al Turgeon
Vice President
Tru Green
E. Lansing, MI



Warren Bidwell
Olympia Fields CC
Olympia Fields, IL



Douglas Chapman
Horticulturist
Dow Gardens
Midland, Michigan



Kent Kurtz
Professor
Horticulture
Cal Poly - Pomona



Harry Niemczyk
Professor, Turfgrass
Entomology
Ohio State University
Wooster, Ohio



Roger Funk
Vice President
Davey Tree
Expert Co.
Kent, Ohio

a Touch of class... Buckner Service.



No matter where you are, a Buckner distributor is near.
All are well trained and dedicated to serve you better than ever before.
Check the map and our distributor list. As you can see we're just about everywhere. So call Buckner for the name of the distributor nearest you.
You deserve a touch of class with fast and efficient Buckner Service.

A W F INC.
AQUA FLO INC.
AQUA MASTER
AUTO RAIN INC.
BADGER TURF & IRRIGATION
PAUL BLAKENEY CO.
BUILDERS PLUMBING & SUPPLY
BURDICK & BURDICK INC.
BURR CONTROLS
CAPITAL TURF & EQUIPMENT CO.
CARLSON'S UTILITY SUPPLY INC.
CENTURY RAIN-AID
COAST IRRIGATION SUPPLY INC.
COMMERCIAL IRRIGATION SERVICES INC.
COMPLETE IRRIGATION
CONELY CO.
CONTRACTORS SUPPLY
DALLAS PIPING PRODUCTS INC.
EARTH IRRIGATION & SUPPLY
EAST COAST SPRINKLER SUPPLY CO.
EVERGREEN IRRIGATION
EWING IRRIGATION PRODUCTS
H.D. FOWLER CO. INC.
FUL-FLO INDUSTRIES LTD.
GRAINGER IRRIGATION
GRASSROOTS TURF PRODUCTS INC.
ERNEST HARDISON SEED CO.
HAWAII IRRIGATION SUPPLY CO.
HILL IRRIGATION

HOME IRRIGATION CENTER
HYDRO SCAPE PRODUCTS
I & E SUPPLY CO.
INDUSTRIAL EXTINGUISHER & SUPPLY
INDUSTRIAL SALES
INDUSTRIES HARNOIS INC.
IRRIGATION & SPRINKLER SUPPLY
IRRIGATION SUPPLY CO. INC.
K & P SUPPLY
KENNEWICK INDUSTRIAL SUPPLY CO.
KERN TURF SUPPLY CO.
KEYSTONE BUILDERS SUPPLY
L & B PIPE SUPPLY
BOB LADD INC.
LAS VEGAS FERTILIZER CO.
LONG ISLAND IRRIGATION
M & A IRRIGATION
MAJOR IRRIGATION
MARSAN TURF & IRRIGATION
MELROSE SUPPLY & SALES CORP.
MORRIS IRRIGATION
S.V. MOFFETT COMPANY
MOUNTAIN SUPPLY COMPANY
NEUMARK COMPANY
ONTARIO TURF EQUIPMENT COMPANY
OLIVER AGRICULTURAL SUPPLY
PAWNEE IRRIGATION SUPPLY
PIPECO
PLASTIC WHOLESALE PLUMBING INC.

PORTER BROTHERS INC.
H.D. QUAD
DOC SAVAGE SUPPLY
SHEMIN NURSERIES INC.
SILVER CREEK IRRIGATION CO.
SISCO TURF
SMITH IRRIGATION & PLUMBING SUPPLY
SMITH SUPPLY
TEXAS TURF SUPPLY
TIECO INC.
TURF IRRIGATION & WATERWORKS
SUPPLY
UNITED SUPPLY CO.
VALLEY TURF & IRRIGATION
THE WARNER CO.
WATSON DISTRIBUTING CO.
WEST COAST IRRIGATION
WESTERN IRRIGATION
WESTERN NEVADA SUPPLY

Bringing the best.

Buckner[®]
Royal Coach Sprinklers, Inc.

4381 N. Brawley Ave./Fresno, CA 93711
(209) 275-0500/TWX 910 362 1167



Seeding REBEL in late September



Only seven weeks after seeding, REBEL provides a rich carpet of dense, fine-leaved turf on Washington D.C.'s Congressional Country Club, Bill Black, Greens and Grounds Chairman.

Proven performance makes **Rebel** number one.

Rebel
TURF TYPE TALL FESCUE

University tests across the country have proven most tall fescues have some good qualities; but in overall performance Rebel beats them all.

1982 Turfgrass Quality Ratings for Tall Fescue Varieties Tested at Eleven Locations in the United States

Variety	University Test Location, Quality Ratings 1-9; 9 = Best											
	NE	KS	MO	IL	GA	KY	MD	MS	NC	NJ	VA	AVERAGE
REBEL	6.7	7.7	7.0	5.6	8.2	7.5	6.5	4.8	7.0	7.6	6.8	6.9
Falcon	6.3	7.6	6.8	5.3	7.7	6.9	6.6	4.8	6.2	7.3	6.4	6.5
Olympic	6.3	7.3	—	5.0	8.1	6.5	6.3	4.2	5.8	6.6	6.4	6.3
Clemfine	—	7.2	—	—	7.3	6.3	5.9	4.1	6.3	4.6	5.9	6.0
Galway	—	7.0	—	5.1	7.5	6.0	6.4	4.8	6.1	5.1	5.5	5.9
KY-31	5.1	7.2	5.5	5.3	6.8	6.3	5.2	4.1	5.9	3.6	5.4	5.5

In these trials, turfgrass varieties were evaluated for appearance, establishment and quality of stand, resistance to drought and disease, and mowing qualities. REBEL took top honors in all-around attractiveness and performance:

- Resists heat, drought and disease
- Fine-textured (30% finer-leaved than Kentucky 31)
- Dense stand (188% denser than Kentucky 31)
- Excellent shade tolerance
- Rich color
- Fast establishment

Circle No. 122 on Reader Inquiry Card

For an all-around, top-performing tall fescue, try REBEL. Test it for yourself.



Lofts Inc.
Bound Brook, NJ 08805
(201) 356-8700 • (800) 526-3890

Lofts/Maryland
Beltsville, MD 20705
(301) 937-9292

Lofts/New England
Arlington, MA 02174
(617) 648-7550

Lofts/Western
Albany, OR 97321
(800) 547-4063

Sunbelt Seeds Inc.
Norcross, GA 30094
(404) 448-9932