



**Dollar spot**, shown at left, damages bentgrass, annual and perennial bluegrass, bermudagrass and zoysiagrass. In the South it is common in the spring and fall. In the north it is active anytime temperatures are between 60 and 85 degrees.



**Brown patch**, at right, is the most common turf disease in the Southeast. It affects primarily St. Augustine and zoysiagrass in the South and bentgrass and perennial ryegrasses in the North. It is prevalent during warm, moist days with cool nights.

# Diseases of Southern Turf

by Don Blasingame, extension plant pathologist, Mississippi State University

**M**anagers of southern turf not only fight diseases of six different warm season turf species, but also diseases of overseeded cool season turf species.

Although most of the diseases of southern turf are caused by fungi, other agents such as bacteria, viruses, and nematodes, can cause serious problems to certain grasses.

Turf managers in the South can't depend solely on fungicides for disease control. Good variety selection, proper fertilization, and appropriate cultural practices are very important in disease control. No amount of fungicide will compensate for poor fertility and cultural practices.

## Brown patch

Brown patch is the most common turf disease occurring in the southeastern United States. Although St. Augustine and zoysiagrass are the most susceptible species, even the more tolerant centipede, bermudagrass and ryegrass are frequently damaged by this fungus.

Brown patch is favored by warm, moist weather combined with cool nighttime temperatures. Therefore, in certain areas of the South, brown patch can and does occur any month of the year.

In the upper regions of the South the most favorable conditions for brown patch development occur from

# TURF DISEASE GUIDE

late April through mid-October.

Symptoms of brown patch on warm season grasses are different than the symptoms of the disease on cool season turf. Even though the grass is usually killed in a circular pattern, many times the smoke ring is not seen on southern turf. Also, under certain environmental conditions the fungus may cause a gradual thinning of the turf over a rather large area instead of killing in a circular pattern.

There are several factors that tend to make the grass more susceptible to brown patch. One is the excessive application of nitrogen fertilizer. The resulting lush growth is readily attacked.

Another is watering late in the afternoon and allowing the grass to remain wet for long periods of time. Excessive accumulation of thatch also creates a favorable environment for development of brown patch and many other diseases.

Fungicides are best used on a preventative schedule. Once symptoms develop control can be difficult.

## Dollar spot

Dollar spot is common on many species of grasses, including bermudagrass, zoysiagrass, and annual and perennial bluegrasses.

Symptoms of dollar spot are different on certain warm season grasses than those noted on cool season grasses.

On the finer textured grasses, such as bermudagrass and zoysiagrass, the disease kills grass in small patches two to three inches in diameter. Under severe conditions, these patches may coalesce so that the turf has a mottled appearance. Blades of grass at the outer edges of the infected area develop tan spots with reddish brown margins.

On the coarser warm season grasses, the turf is killed in larger patches ranging up to a foot in diameter.

Dollar spot is prevalent during periods of mild weather during the spring and fall.

Unlike brown patch, dollar spot is retarded by high levels of nitrogen. Still, turf managers should consider the impact of high nitrogen on brown patch and other diseases.

Watering should be performed only in the early morning so the foliage can dry quickly. Fungicides can be used to help bring the disease under control once it gets established.

## Leaf spots

There are a number of fungi that cause leaf spots on many southern

grasses. Regardless of the causal agent, leaf spots and their control on southern grasses are similar.

**Helminthosporium leaf spots** (*Bipolaris* spp.)—Bermudagrass and ryegrass are most severely affected by helminthosporium infections, although the fungus can survive on centipedegrass and St. Augustine.

Infection can occur over a wide range of temperature, but usually is more severe at 70 to 95 degrees F. Milder temperatures in the spring and

---

**The excessive accumulation of thatch creates a favorable environment for development of brown patch and many other diseases that are caused by fungi.**

---

fall are more favorable for infection.

Helminthosporium causes small, dark-colored spots or flecks on the leaves and sheaths. Leaf spots are usually more numerous near the collar of the leaf blades. Severely affected leaves wither and die and the turf frequently becomes brown and thin.

Symptoms on overseeded ryegrass are altogether different. Although leaf spots may occur, this same helminthosporium can cause severe crown rot. This causes a yellowing and discoloration of the grass and a general thinning of the turf.

Fertilize with adequate levels of nitrogen and potassium if helminthosporium diseases become a problem. With careful management, apply fungicides recommended for helminthosporium blight control.

**Gray leaf spots**—St. Augustine is the primary host for gray leaf spot. The disease occurs throughout the lower South during warm, humid weather.

Spots on the leaf blades are the most visible, but sheath and stem lesions also occur. Leaf spots begin as olive green to brown, water-soaked spots as small as a pin head. These enlarge rapidly and form a circular to elongate lesion that is brown to ash colored with purple margins.

The disease occurs during moderate to warm weather accompanied by high relative humidity. Severity of the disease is enhanced by applications of nitrogen fertilizer. It is more a problem in shaded areas where the grass remains wet from dew.

Treatment with a fungicide may become necessary if the disease outbreak is severe and accompanied by prolonged periods of wet favorable weather.

## Rust

Rust or *Puccinia* species infect a number of grasses grown in the South, including ryegrass, zoysiagrass, bluegrass, fescue, bermudagrass, and St. Augustine. Zoysiagrass and bluegrass are the most often infected grasses.

Susceptibility depends on the variety. Fungus infection is favored by minimum and maximum temperatures of 50 to 70 degrees F. respectively. For this reason, the disease does not usually cause severe damage over an extended period. It is likely to be more severe in shaded areas during rainy, humid weather. Affected turf will appear unthrifty and begin to thin.

The disease is characterized by the presence of pustules on the leaf blades. These pustules range from bright orange to cinnamon-brown in color depending upon the species of fungus present.

Certain varieties of ryegrass are extremely susceptible to rust and sometimes severe damage can occur. Zoysiagrass, especially Meyer and Emerald, are most severely infected by rust.

Fertilize to stimulate grass growth, mow on a four to five day schedule, and catch clippings. If necessary, a fungicide may be applied to help reduce the amount of disease present.

## Spring dead spot

Spring dead spot is a serious disease of bermudagrass in certain parts of the upper Sunbelt. It is found generally on bermudagrass or zoysiagrass under high maintenance.

Damage to the turf apparently occurs during the dormant season. When greenup occurs in the spring, there are areas a few inches to several feet in diameter where the sod is completely dead.

The causal agent for spring dead spot has not been identified. The only control procedures recommended are good cultural practices and limiting the use of nitrogen fertilizer, especially late in the growing season.

Research has shown that



# TURF DISEASE GUIDE

fungicides can limit the damage. However, at the present time only two fungicides are labelled and these may be limited to use in certain states.

## St. Augustine decline

St. Augustine decline (SAD) is caused by a virus. The symptoms are a mosaic type chlorosis of the leaf blades that resemble nutrient deficiency or mite feeding. Evidently there are several strains of the virus since there is a great range in damage to St. Augustine.

To this point, the disease has only been recorded in Arkansas, Texas, Louisiana, and Mississippi. There are no chemicals available for the control of SAD.

There are, however, several varieties of St. Augustine resistant to the virus. These can be planted in areas where the disease is a potential problem.

Floritam was the first variety release that has resistance to SAD. It is also resistant to chinch bugs. It has

poor cold tolerance and should be used only in the lower South.

Seville is resistant to SAD and is more shade tolerant than common St. Augustine. Raleigh has both SAD resistance and good winter hardiness.

## Downy mildew

Downy mildew of St. Augustine was first described on common St. Augustine in Texas in 1969. Since then the disease has spread and has been identified in Arkansas, Louisiana, and Mississippi.

Downy mildew appears as white, raised, linear streaks that develop parallel to the mid-veins of the leaf. Streaks appear in the spring and remain throughout the summer, giving the leaves a yellow appearance with some death toward the tips.

Severe disease occurs in grass grown in flood plains or poorly drained areas.

The white-streak symptom is easily confused with the virus disease, St. Augustine decline. However, the virus symptoms are more yellow in color and more mottled than striped.

Downy mildew has been difficult to control with most common turf fungicides. Good drainage is recommended for cultural control.

## Pythium blight

Pythium blight can be a devastating disease on overseeded ryegrasses. Bermudagrass and other warm season grasses can be affected to a lesser degree.

An abundance of moisture is required for pythium blight development. In addition, the disease is favored by warm temperatures.

Affected grass is killed rapidly in spots two to four inches in diameter. These spots may develop into streaks so that large areas of turf are damaged.

During early stages of development, the affected grass appears wilted and greasy. At times the affected turf spots may have a cottony appearance due to the abundant growth of the fungus. For this reason, the disease is frequently referred to as cottony blight.

Certain species of pythium can also cause root rot on turfgrasses. Due to the restricted root function the plants become chlorotic and the turf begins to thin.

On overseeded grasses the disease can be limited by using treated seed and delaying the overseeding until as late as possible during the fall. Water as little as possible during periods of favorable activity. The perennial

## The Mulching Solution.



## Bale Chopper

The Goossen Solution for fast, economical mulching.

### Mulch in seconds.

The Bale Chopper shreds and blows an entire bale of straw in less than a minute, eliminating tedious hand mulching.

### Mobile mulching.

The Bale Chopper with PTO can be mounted quickly to your tractor or can be used on a trailer. The gas model operates independently and can be used from a truck bed. This lets you use the Bale Chopper both in the fields and at residential and commercial landscaping sites.

### Mulch where you want it.

With the Goossen Bale Chopper you can blow mulch evenly over large areas with the 180° rotating directional blower or the 30', 6" diameter flexible hose.

### Multi-use mulcher.

The Goossen Bale Chopper shreds and applies mulching material to newly seeded lawns, strawberry beds, bedding plants, around trees and shrubbery, or in rows. The Bale Chopper also shreds straw, paper, leaves, and grass.

Find out more about the Bale Chopper, call Goossen Industries today.

IN NEBRASKA CALL

**402/228-4226**

TOLL FREE OUTSIDE NEBRASKA

**800-228-6542**

**Goossen** INDUSTRIES

927 W. COURT • BOX 705 • BEATRICE NE 68310

# TURF DISEASE GUIDE

## Turf Fungicide Directory

Common Name	Brand Name	Company	Circle No.
anilazine	Dymec 50	Gordon	201
	Dyrene	Mobay	202
	Ortho Dyrene	Ortho/Chevron	203
	Proturf Fung. III	Scotts	204
benomyl	Proturf Fung. DSB	Scotts	205
	Tersan 1991	Du Pont	206
cadmium	Caddy	Cleary	207
	Cadtrete	Cleary	208
	Cadminate	Mallinckrodt	209
	Kromad	Mallinckrodt	210
chloroneb	Proturf Fung. II	Scotts	211
	Teremec SP	Gordon	212
	Terreneb SP	Kincaid	241
chlorothalonil	Daconil 2787	SDS Biotech	213
	Proturf 10IV	Scotts	214
cycloheximide	Acti-dione TGF	Nor-Am	215
ethazol	Koban	Mallinckrodt	216
	Terrazole	Olin	217
fenarimol	Rubigan	Elanco	218
iprodione	Chipco 26019	Rhone Poulenc	219
	Proturf Fung. 6	Scotts	220
mancozeb	Fore	Rohm and Haas	221
	Formec 80	Gordon	222
maneb	Tersan LSR	Du Pont	223
mercuries	Calo-Clor	Mallinckrodt	224
	Calo-Gran	Mallinckrodt	225
metalaxyl	Subdue	Ciba Geigy	226
	Ridomil	Ciba Geigy	241
	Apron	Gustafson	242
PCNB	Terraclor 75	Olin	227
PMA (PMAS)	PMA, PMAS	Cleary	228
PMA plus Thiram	Proturf Broad Spectrum Fung.	Scotts	229
propamocarb	Banol	Nor-Am	230
thiabendazole	Mertect	Merck	243
thiophanate	Cleary's 3336	Cleary	231
thiophanate methy	Fungo 50	Mallinckrodt	232
	Proturf Systemic	Scotts	233
thiram	Chipco Thiram 75	Rhone Poulenc	234
	Spotrete	Cleary	235
thiophanate plus thiram	Bromosan	Cleary	236
thiophanate-methyl plus maneb	Duosan	Mallinckrodt	237
triadimefon	Bayleton	Mobay	238
	Proturf Fung. 7	Scotts	239
vinclozolin	Vorlan	Mallinckrodt	240

ryegrasses are not as susceptible as annual. Under severe disease pressure chemical control may be required.

### Fairy rings

Fairy rings generally appear in lawns and other turf areas as circles or arcs of dark green, fast-growing grass during the spring and early summer. A ring of thin dead grass may develop on one or both sides of this circle.

The disease is caused by one of several soil-inhabiting fungi that commonly produce mushrooms. Mushrooms that sometimes appear in the ring are the fruiting bodies of these fungi. Stimulation of the grass is due to the release of nutrients from the organic breakdown of the thatch by the growing fungus.

It is difficult to control fairy ring. There are no chemicals labeled for the control of fairy ring. Two general approaches may be considered, removal and suppression. Although relatively impractical, removal of infected soil and grass to a depth of 12 inches or more in a band several feet on each side of the infected area and replacement with clean soil is one solution.

Another approach is to suppress the disease. For low maintenance grass areas, increase the water and fertilization program to stimulate the declining grass inside the ring. Symptoms of fairy ring can be masked by pumping large quantities of water into this area.

### Slime molds

Slime molds are a group of organisms that cover above-ground plant parts with a dusty gray-black or dirty yellow mass. When you look closely at this growth, you see small round balls scattered over the plant. If you rub these between your fingers a sooty powder emerges. This consists of spores of the fungus.

Slime molds normally live in soil where they feed on decaying organic matter. When the mold is ready to reproduce, it grows up on to the grass blades so that the spores may be spread greater distances.

Slime molds do not feed on living plants. They only use them to assist in the distribution of spores during reproduction.

Slime molds occur during wet weather throughout the spring, summer and fall. They disappear rapidly as soon as it becomes dry. Chemical control is usually not necessary.

WT&T