



Agricultural Experiment Station or U.S. Department of Agriculture. Some well illustrated circulars, bulletins, and books are also available.

**Mechanical Control.** Hand methods of brush control, although effective, are slow, costly, and laborious. They are practical on small areas or in scattered stands.

Hand methods include grubbing, cutting, girdling, and burning.

*Grubbing* consists of using a grubbing hoe, shovel, or similar tool to dig enough of the root system out of the soil to kill the plants. The operation is difficult and time consuming, but effective if properly done.

*Cutting* down brush with axes or saws is most effective on woody species that are killed when the top growth is removed, such as eastern redcedar and blueberry juniper. Species that resprout can be treated with herbicides to prevent resprouting.

*Girdling* is cutting a ring through the bark and cambium layer to prevent movement of water and nutrients to top growth. Girdling is practical in scattered stands of large trees 6-inches in diameter or greater. It is most effective during the summer months. Herbicides can be applied to the cut ring for improved kill.

Portable chain or power saws and girdlers are available for brush control. They reduce labor, time, and cost in brush removal but have limited use in dense stands or large areas.

*Dozing* is one of the more widely used methods of brush control. Much clearing is done with straight dozer blades; however, many modifications and attachments are available for specialized clearing jobs.

Ideally, dozing removes brush and large trees by pushing or pulling the plants out with as much of the roots intact as possible. Special attachments to the straight blade include teeth or U-shaped "stingers" to allow cutting the plant off below the ground line and lifting out the roots.

Dozing is most commonly practiced in open stands of large trees and brush or on rocky soils where other mechanical control methods are limited. It is not desirable in dense stands of brush that sprout from the roots after top removal.

Dozed trees can be windrowed or stacked so the brush can be burned or left to decay. The equipment and fuel required for dozing are costly. Dozing heavily damages the turf and grounds.

*Mowing and shredding* are temporary control methods for weeds and small brush in landscape management. *continued on page 68*

# Right-of-Way Weed Control

by R. W. Bovey, USDA-ARS, College Station, TX

**M**any areas around buildings, ditches, billboards, poles, factories, shops, golf courses, parking areas, parks, industrial plant sites, vacant lots, schools, airports, roadsides, fences, lawns, pipelines, and other public or non-cropland areas may harbor unwanted trees and brush.

If these undesirable woody plants resprout after top removal, they may be difficult to kill and remove and cause unsightly appearance or interfere with the intended use of the property. The landscape manager has many options at his disposal to control and remove undesirable woody vegetation and replace it with desirable cover.

Woody plants can be removed by two primary methods or combina-

tions of these methods which include mechanical or chemical means. The selection of the method depends upon

- 1 size of the area to be cleared;
- 2 size of the brush;
- 3 difficulty of control and kind of brush;
- 4 proximity to valuable vegetation, water sources, and population centers; and
- 5 use to be made of the area following brush removal.

## Control methods

It is essential that problem plants be accurately identified so the proper control methods can be used. If the species cannot be identified, consult the local county agent, extension specialists, or personnel from the State



ment. Repeated mowing, once or twice a year, is needed for maintenance on most weed-infested areas.

Mower types vary, but most consist of sharp rotary blades. Heavy duty shredders can be used on large brush and small trees, three- to four-inches in diameter.

**Disking.** Large disk plows or tandem disks will destroy stands of small brush. They may also destroy the grass stand. Disking is limited to tillable soils. Disking prepares a good seedbed, although compaction by a cultipacker, roller, or other implement may be desired. The operation is expensive.

Cost of mechanical treatments is usually closely correlated with degree of soil disturbance and size and density of brush to be removed. Dozing, disking, and grubbing are among the most effective mechanical brush control treatments, but are the most costly to perform, while mowing is less expensive. Hand methods, such as sawing, axing or grubbing, are sometimes effective, but are slow, costly, and laborious.

**Chemical Control.** Herbicides are an important means of weed and brush control. Compared to mechanical practices, herbicides are usually less expensive, less damaging to the environment, and often more effective.

Herbicide sprays, however, are subject to drift and may damage susceptible crops or valuable vegetation on nearby areas if improperly applied.

A variety of herbicides and herbicide combinations are commercially available. It is necessary to understand the properties and effects of herbicides in order to safely and effectively use them.

Individual herbicides and combinations of herbicides are used for weed and woody plant control. After manufacturing, technical (pure) herbicide must be formulated with other ingredients to prepare usable products for ease of handling and to obtain the desired effects.

Since very small amounts are sometimes required per unit of land, uniform application is essential. Herbicides are formulated as liquid con-

centrates, wettable powders, and granules or pellets. It is important to know the characteristics and precautions to be taken when using herbicides.

### Characteristics of Herbicides

Phenoxy herbicides, such as 2,4-D, 2,4-DB, dichlorprop, and MCPA have

grasslands or grass crops. Rates of 0.25 to 2 pounds per acre effectively control many broadleaf plants.

The phenoxy compounds are relatively inexpensive and easy to apply. They are usually marketed as liquid concentrates as salts or esters.

The ester formulations are often more effective as foliar sprays on trees and brush than the salts. Amine formulations commonly available include dimethylamine, triethylamine, diethanolamine, trimethylamines, triethanolamine, and others.

Other inorganic salts of the phenoxy that have been sold include the ammonium, sodium, potassium, and lithium salts. Salts are sprayed in water carriers.

Esters are classified as high volatile or low volatile, depending upon how readily they vaporize. Low volatile esters should be used in areas where sensitive crops or vegetation are grown.

The concentration of the active ingredient, the "acid equivalent," is indicated on the label as pound-per-gallon. If a herbicide concentrate has an acid equivalent of 4 pounds per gallon, then 1 gallon of the concentrate contains 4 pounds by weight of the parent acid, regardless of formulation. Usually the most concentrated formulations cost less per pound and are more economical to use than weaker concentrates.

The phenoxy compounds are readily absorbed by leaves and are translocated throughout the plant along with the products of photosynthesis. Oil soluble formulations, usually esters, applied in kerosene or diesel oil will penetrate the bark of most woody plants, and can be used as basal sprays or foliar sprays to individual plants.

Phenoxy herbicides, however, are more commonly applied broadcast to large areas containing dense stands of brush. These herbicides are sprayed on above-ground parts and foliage since they are not effective at economical rates as soil-applied herbicides.

**Organic arsenicals.** The organic arsenicals include DSMA, MSMA, and cacodylic acid and are available as liquid concentrates. These compounds

*continued on page 72*

## Calculating Rates and Quantities

### Granular Materials

Example—A landscaper needs to treat 4,000 sq. ft. of Juniper with 4% RONSTAR granules at 4 lbs. aia. How much RONSTAR is required?

Formula  
$$\text{lbs. aia} \times \frac{\text{sq. ft. to be treated}}{44,000 \text{ sq. ft./acre}} \times \frac{100}{\% \text{ granular}} = \text{lbs. required to treat area}$$

Calculation  
$$4 \text{ lbs. aia} \times \frac{4,000}{44,000} \times \frac{100}{4} = 9.1 \text{ lbs. 4\% granular RONSTAR}$$

### Wettable Powders

Example—A landscape firm needs to treat 4,000 sq. ft. with 50% DEVRINOL wettable powder at the rate of 10 lbs. aia.

Formula  
$$\text{lbs. aia} \times \frac{\text{sq. ft. to be treated}}{44,000 \text{ sq. ft./acre}} \times \frac{100}{\% \text{ powder}} = \text{lbs. required to treat area}$$

Calculation  
$$10 \text{ lbs. aia} \times \frac{4,000}{44,000} \times \frac{100}{50} = 1.8 \text{ lbs. 50\% DEVRINOL W.P. to be mixed in enough water to cover 4,000 sq. ft.}$$

### Liquid Concentrate

Formula  
$$\text{lbs. aia} \times \frac{\text{sq. ft. to be treated}}{44,000} \times \frac{1}{\text{lbs. active ingredient/gallon}} = \text{gallons required to treat area}$$

Calculation  
$$2 \text{ lbs. aia} \times \frac{22,000}{44,000} \times \frac{1}{4} = .25 \text{ or } \frac{1}{4} \text{ gal. TREFLAN to be mixed in enough water to cover } \frac{1}{2} \text{ acre}$$

been used for over 30 years and are effective for the control of many weed and brush species. They are used to produce changes and shifts in plant cover for crops, pastures, lawns, or wildlife habitat.

The phenoxy herbicides are not toxic to livestock or man at dosages labeled for weed control and disappear rapidly from the soil, vegetation, and water. They do not accumulate in the food chain.

Susceptible vegetation, especially broadleaf plants, may be damaged from spray drift or from volatilization. Following label instructions and making applications during favorable weather should prevent drift and volatilization problems.

The phenoxy herbicides selectively control broadleaf weeds in

have limited use for woody plant control as foliar sprays.

DSMA and MSMA are used for postemergence weed control in tolerant lawn grasses and noncrop areas. Cacodylic acid is used as a general desiccant and defoliant (contact) spray on many crop and non-crop areas.

The organic arsenicals can be used for quick dieback of woody species, but plants tend to recover. They can be injected into the trunk to kill unwanted trees. As foliar sprays, they are applied in water. Complete coverage by spray is required for best results. They are inactivated by soil contact.

Organic arsenicals may injure desirable plants. By directed spraying, desirable plants can be avoided. Organic arsenicals have a moderate to low mammalian toxicity but are highly toxic to insects such as honey bees. They are mildly corrosive to spray equipment.

*Bipyridyliums*. Diquat and paraquat are desiccant (drying agents) and defoliant herbicides used for general contact activity against weeds and brush. In some situations, they are used as selective herbicides.

Paraquat is water soluble and is inactivated by soil contact. Woody species will usually resprout from foliar sprays of diquat and paraquat. Paraquat may be more effective as an injection treatment against some undesirable trees.

Paraquat is highly toxic when ingested. Extreme care must be taken to avoid breathing the spray mist or getting the concentrate on the skin. Use

of protective clothing and respirators are recommended when making applications.

Get immediate medical care in the event of ingestion. If treated within 12 hours, chances of survival are excellent.

*Benzoics*. Dicamba is a selective translocated herbicide. It controls many broadleaf weeds in pasture crops and turf and some woody plants.

Dicamba is similar to the phenoxy herbicides in activity and use. It is absorbed through roots, as well as foliage. It may be applied by either ground or aerial sprays or as granules, depending upon the weeds to be controlled and their proximity to susceptible crops.

Dicamba may be applied in mixtures with 2,4-D to broaden the spectrum of weed species controlled. It has a low order of toxicity to wildlife, fish, livestock, and humans. It rapidly degrades and does not accumulate in the environment. Dicamba has a low corrosion hazard to spray equipment.

It is formulated as a liquid as the dimethylamine or sodium salt and is sprayed in water carrier. It is also formulated in granular form as the acid.

Dicamba is highly water soluble. Care must be taken to prevent its movement into water sources. Spray drift of dicamba to sensitive crops, conifers, and certain woody plants should be prevented. Granular formulation may be preferred where drift of sprays would present a hazard to crops.

Dichlobenil is also a benzoic. It is a selective herbicide for control of an-

nual and perennial grasses and weeds in non-crop areas. Dichlobenil inhibits weed seed germination and plant cell division at the growing points.

Because of its relatively high volatility, application when the temperature exceeds 70 F. must be followed by a quarter-inch irrigation or rainfall soon after. Downward movement of active ingredient does not exceed two inches, so deeper-rooted woody ornamentals will not be effected. The control is achieved from a chemical barrier in the top one to two inches of soil.

*Ureas and uracils*. These compounds include bromacil, diuron, fenuron, fenuron-TCA, monuron, monuron-TCA, tebuthiuron, and hexazinone.

Ureas and uracil-type herbicides can be selective at low rates and non-selective at high rates. They are usually formulated as wettable powders for water sprays or as granules or pellets for dry application.

Bromacil (a uracil) will control a wide variety of woody species. If rates above 5 pounds per acre are used, it will also kill many desirable grasses and forbs on grazing lands.

Fenuron is no longer produced commercially, but a fenuron-TCA combination is available and is used to control certain woody plants and weeds on non-crop areas. Monuron and monuron-TCA combinations are commercially available, but have limited use for brush control since high rates are required for effectiveness. The monuron-TCA or fenuron-TCA combinations are generally used for non-selective, temporary sterilization in non-crop areas.


Tebuthiuron provides excellent control of a variety of undesirable woody plants. This herbicide is formulated as pellets and contains 20 or 40 percent active ingredient. A new "brush bullet" containing 13.8% active ingredient is also available. Tebuthiuron is commercially available for grazing lands and weed control on non-crop areas.

Hexazinone is a relatively new compound showing promise for woody plant control and use on non-crop areas. Hexazinone is recommended for forestry site preparation and pine release where loblolly, long leaf and short leaf, slash, and Virginia pines are grown.

The pelletized product can be spread in a grid pattern for hardwood brush control. It also shows promise for total vegetation control including perennial grasses. It is highly water soluble.

The ureas and uracils mentioned

### The Effect of Delaying the Application of Preemergence Herbicides





## Herbicide Directory

Herbicide	Brand Name(s)	Company	Uses
amitrole	Amitrole-T	Union Carbide	Non-selective grass and broadleaf control in non-crop areas including ditches, roadsides, industrial areas and marshes.
arsenal	Arsenal	American Cyanamid	Non-selective grass and broad leaf control in non-crop areas.
ammonium sulphamate	Ammate	DuPont	Non-selective rights of way herbicide
asulam	Asulox	Rhone Poulenc	Postemergence grassy weed control for turf and ornamentals
atrazine	atrazine	Ciba Geigy	Non-selective control in non-crop areas.
benefin	Balan	Elanco	Preemergence control of annual grasses and broadleaf weeds in established turf.
bensulide	Betasan Pre-San Lescosan Betamec-4	Stauffer Mallinckrodt Lesco PBI Gordon	Preemergence control of annual grasses and broadleaf weeds in established turf and established flower gardens. Safe near tulip and daffodil bulbs.
bentazon	Basagran	BASF	Selective postemergence control of nutsedge in warm-season turf.
bromacil	Hyvar	duPont	Nonselective control of weeds and grasses in non-crop areas. usually mixed with diuron for roadsides and rights-of-way.
bromoxynil	Brominal Buctril	Union Carbide Rhone Poulenc	Postemergence control of broadleaf weeds in seedling turf, established turf and non-crop areas.
cacodylic acid	Phytar Rad-E-Cate	Vertac Vineland	Nonselective control for turf renovation, edging and in plant beds.
chloramben	Amiben	Union Carbide	Preemergence control in ornamentals.
chlorfurecol	Maintain	Uniroyal	Growth regulator. Also controls broadleaf weeds and vines.
copper	Citrine-Plus	Applied Biochemists	Control algae, chara and hydrilla in potable water.
dalapon	Dalapon 85 Dowpon M	SDS Biotech Dow	Selective control of perennial and annual grasses in non-crop areas and ditchbanks.
dazomet	Mylone	Hopkins Ag.	Preplant sterilant for turf and ornamental beds.

*continued on page 78*

are absorbed primarily through the roots of plants. They may be applied in spring or fall when weeds and brush are actively growing and when adequate rainfall leaches them into the soil.

Fall, winter, and early spring applications of tebuthiuron and hexazinone can be timed to reduce injury to forage plants and eliminate hazards of drift. These compounds may kill trees at a considerable distance from the point of application, depending upon the size of the root system and whether it extends into the treated areas. Therefore, extreme care should be taken near desirable trees.

Ureas and uracils are non-volatile and do not corrode equipment. Most of the urea and uracil herbicides can be injurious to some forage and turf species when applied broadcast, especially as sprays. Applying herbicides as pellets or balls to confine the herbicide to a few spots in the treated area reduces exposure to desirable forage or turf plants. Also, application of granules, pellets, or sprays to the soil surface or sprays applied subsurface in rows or bands spaced 6 to 10 feet apart may reduce injury to forage or turf and may increase kill of some woody plants.

Most of the ureas and uracils persist in the soil for several months at rates used for brush control. They are low in toxicity to warm-blooded animals.

### Other Organic Herbicides

*Arsenal* (AC 252,925) is a new broad spectrum herbicide with activity against herbaceous and woody plants. It will control sedges, grasses and broadleaved weeds applied either preemergence or postemergence.

*Arsenal* is readily absorbed through foliage and roots. Complete kill may take several weeks.

Most herbaceous weeds are susceptible to *Arsenal* at rates of 0.4 to 1 pound per acre whereas woody species may require higher rates of 0.75 to 3 pounds per acre depending upon species to be controlled.

*Arsenal* is being tried experimentally for conifer release and weed control in bermudagrass. *Arsenal* should be applied postemergence to perennial weeds when they are actively growing and the herbicide is readily translocated.

It has a low order of toxicity to warm-blooded animals. It persists for 3 to 12 months in the soil depending upon rate applied and soil moisture content.

*Picloram* is a selective, translocated herbicide that effectively controls many weed and brush species in

*continued on page 78*

## Herbicide Directory *continued*

Herbicide	Brand Name(s)	Company	Uses
DCPA	Dacthal	SDS Biotech	Preemergence control of annual grasses and broadleaf weeds in turf and ornamental beds.
dicamba	Banvel	Velsicol	Selective postemergence control of broadleaf weeds in turf and for noncrop control of brush.
dichlobenil	Dyclomec	PBI Gordon	Selective weed control in ornamental beds and for total weed control on roadsides, fencerows, etc.
dichloprop	2,4-DP	SDS Biotech	Brush control and aquatic weed control.
diphenamid	Enide	TUCO, Upjohn	Selective control of annual grasses and broadleaf weeds in bermudagrass, dichondra and around ornamentals.
DSMA	Methar 30 Ansar DSMA Liquid DSMA 81% Weed-E-Rad	W.A. Cleary SDS Biotech Drexel Vertac Vineland	Selective postemergence control of sedges and grasses in turf and ditchbanks and storage yard.
diuron	Karmex Dynex Diuron 80WP Urox	du Pont Vertac Drexel Hopkins	Generally used at high rates for nonselective total weed control in industrial sites.
diquat	Diquat	Ortho, Chevron	Aquatic weed control.
endothall	Aquathol K Endothall	Pennwalt Pennwalt	Aquatic weed control and turf herbicide and dessicant.
EPTC	Eptam	Stauffer	Selective control of annual grassy weeds, nutgrass, and perennial weeds.
Ethofumesate	Prograss	Nor-Am	For control of <i>Poa annua</i> and white clover in fairways.
fenac	Fenac	Union Carbide	Preemergence long-term control of weeds, usually mixed with other herbicides.
fosamine	Krenite	du Pont	Brush control.
fluazifop-butyl	Fusilade	ICI Americas	Selective postemergence control of grassy weeds in ornamentals.
floridone	Sonar	Elanco	Broad spectrum herbicide for submersed and emersed aquatic weeds.
Glyphosate	Rodeo	Monsanto	For control of emerged aquatic weeds and broad leaf weeds in or near aquatic sites, such as ditchbanks.
glyphosate	Roundup	Monsanto	Nonselective, short-term herbicide for turf renovation and total weed control along fences and plant beds.

grasslands. It can be applied to the soil or foliage and is effective as injection/cut-surface treatments on many undesirable trees.

Picloram can be applied in liquid sprays and as pellets to brush in the spring and fall, depending upon the species to be controlled. It is absorbed by both foliage and roots. Most perennial grasses are resistant. Its high activity against many woody plants at moderate rates makes it desirable for brush control. Care must be taken to prevent drift of picloram to desirable plants.

Picloram is relatively persistent in soil, especially in cooler climates. Since it is water soluble, care must be taken to prevent its movement into water used for irrigation. It should not be applied where it can be leached or moved to sensitive plants by rainfall.

Picloram has a low mammalian toxicity and is only slightly corrosive to spray equipment.

Triclopyr (Garlon) is a relatively new selective postemergence herbicide for use of rights-of-way, industrial and forestry sites, and experimentally for rangeland brush control. It can also be used to kill trees and brush by injection/cut-surface treatments.

It is readily translocated in plants and is moderately toxic to warm-blooded animals. It degrades rapidly in the soil.

Ester formulations are available.

Glyphosate is a nonselective herbicide effective against both grasses and broadleaf plants.

Glyphosate is readily translocated from leaf and stem tissue to roots, resulting in a high degree of kill on many weeds. It is inactivated by contact with the soil and should not injure newly-seeded plants in treated soil.

Glyphosate is sprayed in water carrier. It is presently registered for use for noncrop and pre-till weed control and as a directed spray for orchards, plantations, Christmas trees, and many other crops. It is corrosive to some equipment.

Broadcast sprays over woody species will damage desirable forage plants. Applications should be made to individual plants on noncrop areas or areas to be renovated.

Glyphosate has a low order of mammalian toxicity.

Amitrole is effective against poison ivy and poison oak. If amitrole is accidentally sprayed on desirable plants, they are less likely to be severely injured than sprays of phenoxy, dicamba, or picloram.

Amitrole is available as a powder

Herbicide Directory *continued*

Herbicide	Brand Name(s)	Company	Uses
imidazoline	Arsenal	American Cyanamid	Broad spectrum systemic industrial herbicide
linuron	Lorox	du Pont	Short-term control of annual weeds in roadsides and fence rows.
mefluidide	Embark	3M	Growth regulator that suppresses <i>Poa annua</i> .
methyl bromide	Dowfume	Dow	Fumigant for pre-plant control. Also kills weed seed.
metham	Vepam	Stauffer	Preplant soil fumigant killing weeds, weed seed, insects and fungi.
metribuzen	Sencor	MoBay	Postemergence control of goosegrass in warm-season turf.
MCPP	MCPP Mecomec MCPP Chipco Turf Herbicide	SDS Biotech PBI Gordon WA Cleary Rhone Poulenc	Selective broadleaf weed control in turf. Often combined with other herbicides.
MSMA	Ansar Broadside Ansar Weed-Hoe	SDS Biotech Vertac Drexel Vineland	Postemergence selective control of crabgrass and broadleaf weeds in turf. Also, grassy weed control in ditchbanks, roadsides, industrial areas.
napropamide	Devinol	Stauffer	Selective control of weeds in ornamental beds and containers. Experimental combination with Betasan for season-long crabgrass control in turf.
oryzalin	Surflan	Elanco	Preemergence control of weeds in established ornamentals.
oxadiazon	Ronstar	Rhone Poulenc	Preemergence control of weeds in ornamentals and turf.
oxyfluofen	Goal	Rohm & Haas	Selective control of weeds in ornamentals.
paraquat	Paraquat	Ortho, Chevron	Nonselective control of weeds in rights-of-way, industrial areas and fencerows.
pendamethalin	Proturf Weedgrass Control	Scotts	Preemergence turf herbicide for control of grassy and broadleaf weeds.
picloram	Tordon	Dow	Systemic, long-term killer of woody plants and broadleaf weeds.
prometon	Pramitol	Ciba Geigy	Nonselective herbicide with long residual for industrial weed control.

*continued on page 86*

containing 50% active ingredient or as a liquid formulation. Another formulation, Amitrole-T, contains 2 pounds per gallon of ammonium thio-cyanate in addition to 2 pounds per gallon of amitrole.

Amitrole is effective through the roots and tops of plants. Amitrole cannot be used where there is any possibility of residues on food or feed crops.

Fosamine (Krenite) applied as foliar spray will kill or suppress several different woody species on non-crop land. If it is applied to trees or brush in late summer or early fall, susceptible species will fail to leaf out the following spring and subsequently die.

Rates of 1-1/2 to 3 gallons of fosamine are applied per acre during the two month period before fall leaf coloration. If rainfall occurs within 24 hours of application, effectiveness may be decreased. Use only in accordance with recommendations on the label. Fosamine cannot be used on cultivated land or in pastures.

**Inorganic Herbicides**

Ammonium sulfamate (AMS, Am-mate) is a nonselective herbicide. It is used extensively to kill all plants growing on rights-of-way.

AMS kills trees. It can be used as a foliage spray or in cuts through the bark. When applied to stumps, it prevents sprouting. AMS kills or suppresses herbaceous plants as readily as woody plants.

It is corrosive. Spray equipment should be cleaned immediately and coated with diesel oil or similar light oils after use.

**Oils** Diesel oil and kerosene are commonly used to control honey mesquite and huisache. One cup to 1 gallon of oil is used per tree, depending upon its size. The oil is applied around the base of the tree during dry weather when the soil is pulled away from the trunk. Application at this time enables the oil to penetrate to the lower buds on the stem.

Oils alone are not very effective herbicides when applied to the foliage of woody plants. However, diesel fuel is commonly used as a diluent and carrier for some oil soluble herbicides.

It is also used as a carrier in aircraft spraying, usually as a 1:4 oil-in-water emulsion. The oil reduces evaporation of the herbicide in arid areas and aids in penetrating stems and foliage of some woody plants. Water carrier, however, is equally as effective as oil carriers in some situations, such as use of 2,4-D on sagebrush.

*continued on page 82*