



TURF INSECT CONTROL

Despite research on alternatives and supplemental methods, insecticides remain the primary means of controlling insect damage to turf.

by Dr. Harry D. Niemczyk, OARDC; and Dr. Patricia Cobb, Auburn University

Many species of insects and other closely related arthropods live in the turfgrass environment. To control the relatively few we consider pests, it is wise to think of them in relation to the specific segment of the turfgrass environment in which they live.

Some, like mites and aphids, spend some or all of their lives on the grass blades; others such as billbug adults, cutworms, chinchbugs and sod webworms live in the thatch. Grubs usually live at the thatch-soil interface where they feed on thatch and living roots.

Focusing on the specific segment of the environment inhabited by a pest and delivering the control material or management practice so that it has the desired impact on the pest is known as the "target principle." Simple, but the amount of success in achieving control is directly related to the extent to which the target was reached.

Keys to control

Knowing the seasonal occurrence and damage of all life stages of each pest common to your area is a major step toward effective control. This information, combined with the characteristics of the particular turfgrass cultivar and the known length of residual of the proposed insecticide, makes insect control scientific rather than speculative.

Still, there will be twists to confuse any control program, such as soil type, heavy thatch, weather, and poor application uniformity.

Differences between the cool- and warm-season zones and among mountains, plains, and coastal areas, also result in variations in pest species and their seasonal occurrence.



Larger sod webworm (*Pedesia trisecta*) adult.



Grubs at a thatch/soil feeding, and a sod webworm in the thatch.

Knowledge of each pest's life cycle in your area is often as important as the choice of insecticide.

The purpose of this guide is to point out some major pests to watch out for in cool- and warm-season turfs in 1986, when their vulnerable stages occur, and some insecticides that may be used. No endorsement of named products is intended nor is criticism implied for those not mentioned.

Late winter (Mar.)

Chinchbugs and Billbugs—In northern zones chinchbugs and billbugs both overwinter as adults

in thatch or sheltered sites near buildings. They can become active during warm days in March. Infestations of hairy chinchbug and bluegrass billbug also occur in zoysia, Kentucky bluegrass and fine fescues.

In southern Florida, the southern chinchbug is active throughout the year. Most varieties of St. Augustinegrass and some bermudagrass are damaged by southern chinchbugs. Zoysia and bermudagrass are more likely to be infested by the hunting billbug.

When summer damage from chinchbugs and/or billbugs is expected, a preventative application of liquid or granular Dursban (chlorpyrifos-1 lb. AI/acre), or Oftanol (isofenphos-2 lb. AI/acre) may be made as soon as these insects begin to move about. Treatment at this time controls adults before eggs are laid. If spring is early, these applications may be needed as early as the second week of March. During a late spring, applications may need to be delayed until the last week of March.

Retreatment for chinchbugs in mid to late summer may be necessary if reinfestation from adjacent untreated



Vertical cross-section of black cutworms in their burrow.

areas occurs.

Preventative treatments may not be successful in southern Florida where the southern chinchbug has multiple generations and is resistant to most organophosphate insecticides in some areas. In southern Florida, where resistance is a problem, the insecticides Pydrin, Pounce, or Baygon have been substituted for organophosphates. Replacing susceptible turf varieties with Floratam St. Augustinegrass, a variety highly resistant to the southern chinchbug, will provide excellent natural control.

Grubs—The larvae of this group of pests normally overwinter six inches or deeper in the soil. If spring comes early, grub activity can be expected along with skunks and racoons who will tear up the turf searching for the grubs. Moles, which feed on grubs and earthworms, also become active at this time.

Application of Oftanol (2 lb. AI/acre) during March when frost is gone from the ground, provides control of overwintered grubs as they return to the surface. This treatment may not provide sufficient residual to control the late summer (July-August) infestation of grubs. Treatment at this time kills overwintering chinchbugs and billbugs and reduces infestations of these insects during the summer.

Mole crickets—Mole crickets have extended their range from Florida and eastern Georgia into southern Louisiana, eastern Texas and up the East Coast into the Carolinas. Timing of

treatments is critical and varies from one area to another.

The tawny and southern mole crickets are the primary pest species. Except for southern Florida, both have one generation per year. Mole crickets become active in March from north central Florida throughout their range in the Gulf States after overwintering deep in the ground as adults or nymphs. Tunnelling damage takes place at night in moist soil and increases as mole crickets become more active. Both mole cricket species begin spring mating flights in late March. In most areas March treatment is seldom required.

In years when feeding of overwintered mole crickets resumes earlier than normal, Oftanol (2 lb./AI/acre) has been used with some success. Generally, such applications are better made later in the year.

Black turfgrass ataenius—This golf course pest overwinters as an adult in the soil under debris in roughs or other protected areas. A few may be seen flying about on warm afternoons in early March. Usually this activity begins when crocus starts blooming and intensifies as the bloom of red bud appears.

While an application of Oftanol in March may be successful in preventing summer infestations of larvae, the probability of success is increased by waiting until April.

Greenbug—The only stage of the greenbug known to overwinter in northern states is the egg. Shiny black eggs deposited the previous fall may

be found adhering to grass blades fallen tree leaves, or other debris.

Treatment for the greenbug is not appropriate during the late winter.

Sod webworms—In cool-season areas, the most common sod webworm species overwinter as larvae in the thatch or upper inch of soil. Feeding does not resume until hibernation (dipause) is broken by early spring warmth.

Treatment for sod webworm is usually not appropriate during late winter.

Spring (April-May)

Chinchbugs and Billbugs—As warm days of spring approach, movement of chinchbug and billbug adults increases rapidly. Generally, egg laying begins the first week of April on warm-season turf and the first week of May on cool-season turf. Occasionally adult billbugs can be seen wandering about sidewalks on warm afternoons.

Generally, application of insecticides to prevent infestations of chinchbugs and billbugs should be completed by the first week in May in cool-season and mid-April in the South. Such applications are made before significant numbers of eggs are laid. This time may vary as much as a week or more depending upon the spring weather.

When the preventative approach is not used and southern chinchbugs are detected in May, diazinon (4 lb. AI/acre) provides control. In areas with three to five generations, two retreatments at six-week-intervals may be needed.

Grubs—Overwintered grubs return to the surface and begin feeding on turfgrass roots in April. Increased activity and damage from birds, moles, skunks, and racoons foraging on grubs can also be expected. Feeding by birds, mammals and grubs continues through May.


In cool-season areas, a single application of Oftanol (2 lb. AI/acre) made during April has been successful in controlling overwintered grubs with one year life cycles. This treatment and similar applications of Oftanol in May or June may not provide control of late summer infestations.

Infestations of such grubs can also be controlled during April (South) or May (North) by spot or general treatment with Turcam (bendiocarb, 2 lb. AI/acre), Peoxol (trichlorfon, 8 lb. AI/acre) or diazinon (5-6 lb. AI/acre). Golf course superintendents may also use ethoprop (Mocap, Scotts



INSECT CONTROL GUIDE

COMMON SOD WEBWORMS OF OHIO
WOOSTER, OHIO 1978-1982



Nematicide/Insecticide, 10 lb. AI/acre). Sevin (carbaryl, 2-4 lb. AI/acre) has been effective against larvae of the green June beetle.

Treatment should be delayed until grubs are in the top one inch of soil. Irrigation or rainfall should allow such applications to aid in moving the insecticides to the target grub as soon as possible.

Although milky spore disease products for control of Japanese beetle grubs may be applied anytime there is no frost in the soil, spring is a good time for such applications. The soil is open and frequent rains move the disease spores into the soil and thatch. It should be noted that only the Japanese beetle grub will be affected by milky spore.

Incidents of large grub infestations (larvae of June bugs) have been increasing in cool-season areas over the past three years. Locations of such infestations should be identified because reinfestation is likely every three years.

Controls such as Oftanol, diazinon, Proxol, or Turcam should be applied in August or September during years of when large numbers of adults are seen.

Eggs are laid in May and June, therefore treatment should be made in late summer, early fall of that year or early the next spring while the larvae are small. Later application against full-grown larvae have given inadequate control.

Mole crickets—Damage increases in April from north central Florida throughout the southern areas of the Gulf States. Mating and dispersal

flights continue as egg laying and hatching begin.

Spring treatment may be necessary in areas that were severely damaged last fall. Small damaged areas can be rolled or otherwise packed down so the turf roots are reconnected with the soil. To determine cricket presence, pour soapy water (2 oz. liquid dishwashing detergent in one gallon of water) on turf areas where infestation is suspected. Crickets will usually surface in 3 to 15 minutes (longer in cool weather).

Turcam (2 lb. AI/acre), diazinon (spray or granules, 5-6 lb. AI/acre, commercial turf only), or Oftanol (granular or liquid, 2 lb. AI/acre) can be used to control spring infestations.

In less critical areas, short residual treatment with Turcam (2 lb. AI/acre) or diazinon (5-6 lb. AI/acre) applied in late April or May may be adequate. Orthene, 755 Tree and Ornamental Spray (2 lb. AI/acre) applied late in the day and left unirrigated overnight may provide quick knockdown but little residual.

Critical turf areas may require greater residual control provided by early April insecticide applications. Mocap 10G (10 lb. AI/acre) provides up to four weeks control and Oftanol (2 lb. AI/acre) up to eight weeks control. Treatments should be made late in the day if possible and watered immediately.

Black turfgrass ateniensis—Adults of the black turfgrass ateniensis can be seen flying about in April and are often found in clipping catchers after early mowing of golf course greens. These adults begin laying eggs in

early May, or about the time Vanhoutte spirea first comes into bloom. Check with local extension entomologists for more precise time if needed.

Applications of Oftanol during April or May has successfully prevented larval infestations during the summer. Diazinon (5-6 lb. AI/acre) applied to fairways during egg laying kills adults and also prevents the development of summer larval infestations.

A word of caution—diazinon applications may be toxic to waterfowl such as geese feeding on the treated turf.

Sod webworms—Overwintered larvae of the sod webworm begin feeding as soon as the grass begins to grow. Usually damage is insignificant, but areas that do not green up may be infested. These areas frequently have probe holes from starlings feeding on the larvae.

In warm-season areas webworm larvae pupate during late March and early April. Moth flights begin in April in southernmost areas and during May in more northern areas.

Young larvae are usually present about two weeks after the spring moth flight peaks, so treatment of young larvae can be done in May in some areas.

Damage from the burrowing sod webworm may be evident in late May in the South. Rubbing a hand over turf suspected of being infested exposes larval burrows that are covered with a flap of duff and grass clippings.

When necessary, a wide range of insecticides including diazinon, Proxol, Aspon, Sevin, and others applied at labelled rates may be used to achieve control.

Cutworms—Moths of cutworms begin laying eggs on golf course greens and other turf areas in the spring. These eggs hatch producing larvae that feed on grass blades during the night. The black cutworm is the most common species on cool-season turf.

While visible damage in uncommon on home lawns, damage can be significant on golf course greens in late May.

Black, granulate, and variegated cutworm moths become active in March and April in the South. Larvae are present on turf, especially on golf greens and tees. Damage can become evident as early as mid-April. By May, the larvae are large enough to cause severe damage.

Generally the insecticides effective against sod webworm are also effective



Green June beetle grubs in their natural habitat—the turf.

fective against cutworms. The *target principle* of controlling these pests is to apply the insecticide late in the afternoon and allow night feeding cutworms to contact and feed on the treated foliage. Irrigation following liquid application is therefore not advisable unless specified on the product label.

Greenbug—Greenbug eggs begin hatching as early as April, but significant infestations do not develop until later in the year. Aphid numbers are too low to detect.

Winter grain mite—Damage from this mite is often first noted in April when turf areas are receiving spring fertilizer applications. By late May, the mites will have laid their eggs and died. Mites do not appear again until the eggs hatch in October.

If treatment is necessary, liquid diazinon or Dursban will provide control.

Clover mite—Incidents of visible damage to home lawns has been seen in April in several Ohio cities and Denver, Col. Usually a nuisance pest in and around homes, the clover mite appeared in large numbers (5,000 per sq. ft.) across entire lawns and on turf next to building foundations. Symptoms of injury were the same as the winter grain mite. Turf next to foundations was often killed.

The clover mite has a slightly pink body and eight pale colored legs. The first pair of legs are extremely long and protrude well out in front of the mite. The absence of bright red legs distinguishes the clover mite from the winter grain mite.

Treatment with liquid diazinon

(2.5 lb. AI/acre) or Dursban (1 lb. AI/acre) readily provides control.

Fire ants—Fire ants are spreading across much of the South causing serious and painful injury to man and animals. They begin establishing new mounds during warm, wet days of spring. During this time, ants are active near the surface of mounds and workers are foraging for food.

Mound treatments include diazinon granules or drenches; Orthene 755 dust, various Dursban formulations, Oftanol, or MC-96 (trichloroethane). Read the label for specific directions for mound treatment. Do not disturb the mound before or during treatment.

Where mound treatment is impractical, the turf can be treated with Amdro fire ant bait (no more than 1.5 lb. AI/acre). All the bait should be used within three days of opening. Retreatment during the fall is usually necessary.

Summer (June-August)

Chinchbugs—In northern cool-season turf, chinchbug eggs continue to hatch into June. Bright red nymphs appear. The number of chinchbugs increases rapidly in June and peaks in July and August when northern lawns can receive severe damage. This damage is often masked by summer dormancy of turf caused by drought. Hot, dry conditions are ideal for chinchbugs.

During August the nymphs molt into adults that mate, lay eggs, thus producing a second generation. Some northern areas have only one generation per year.

Southern chinchbugs are not usually a problem in well-irrigated turf or during summers when rainfall is plentiful. Southern chinchbug-damage first appears during the dry periods of June and July. Damage may continue throughout the summer and into the fall because of overlapping generations.

A wide range of insecticides may be used at label rates to control existing infestations. They include Dursban, diazinon, Aspon, and Sevin. Treatments should be made before injury is severe, otherwise, damaged areas may not recover.

Areas of southern Florida have pockets of southern chinchbugs resistant to these insecticides. Pydrin, Pounce, or Baygon may be substituted. Floratam St. Augustine, a chinchbug resistant variety, should be the primary turf variety grown in more southern coastal areas and Florida where southern chinchbug is a problem.

Billbugs—The bluegrass billbug larvae feed in grass stems during June and move to the plant crowns and roots and rhizomes during July. This feeding causes brown spots that frequently resemble the symptoms of some fungus diseases. Symptoms are also often masked when the turf is dormant from drought. The larvae usually move deeper into the soil under dry soil conditions. During late July and August the larvae burrow deeper into the soil to pupate and transform into adults.

Infestations discovered during this time may be treated at the same rates used for existing grub infestations with diazinon, Turcam, and Proxol.

Southern chinchbugs are not usually a problem in well-irrigated turf or during summers when rainfall is plentiful.

Irrigation or rain following application is helpful for optimal results. If larvae are feeding in the foot zone, control may be difficult to achieve. Insecticide applied during June should control feeding larvae.

Grubs—By June, in cool-season areas, grubs have stopped feeding and are in the pupal stage three to four inches in the soil. Beginning in mid-June and continuing through mid-July, the adults of various species emerge and burrow into the soil to lay eggs. Hatching and appearance of young larvae occur during July and August.

In warm-season areas, beetle flights continue and often peak in June, although the time flights occur varies from year to year. Japanese beetle flights occur mainly from mid to late May and June. Brown May or June beetle flights often follow heavy rains in late May and June. New generation grubs of most southern species can be found by mid-August.

Existing infestations of grubs found in July or August may be treated with Proxol, Turcam, Oftanol, diazinon, or Mocap (commercial turf only) at standard label rates. Sevin (2-4 lb. AI/acre) is effective against the green June beetle larvae.

Extreme heat and drought during the summer may cause some grubs to



move deeper in the soil. Under such conditions, irrigation several hours before treatment and a thorough soaking afterward is advisable.

Mole crickets—Egg laying diminishes in late June, and newly hatched nymphs of both species feed voraciously. Tunneling damage suddenly becomes obvious in July as the nymphs grow larger. Because of the potential for sudden damage at this time, turf areas should be inspected several times a week during this period.

Bait formulations have been effective in controlling mole cricket nymphs from June through August in the area from central Florida north and west through the Gulf States. Baits work best in eastern Georgia during spring and fall. Bait applications often must be repeated one or more times.

Bait formulations available include: 2% Baygon (1/2 lb./1,000 sq. ft.), 20% Sevin (5-10 lb. bait/acre), 5% Dursban (150 lb./acre or two applications of 75 lb./acre three weeks apart), and 2% malathion (100 lb./acre or two applications of 50 lb./acre three weeks apart).

Mole crickets are more active at night in moist soil. Turf should be irrigated several hours before baits are applied. Delay application until later in the day and do not irrigate for two-to-three days thereafter. Orthene 755 Tree and Ornamental spray (2 lb. AI/acre) can be applied late in the day and left unirrigated. However, re-treatment may be necessary due to short residual activity.

Residual control of mole crickets with Oftanol (2 lb. AI/acre) may vary with location and amount of rainfall. Applications of Oftanol have given up to 12 weeks control from the Florida panhandle along the Gulf Coast. Residual control was only six weeks in areas where August rainfall exceeded eight inches weekly. Oftanol works faster on mole cricket nymphs when watered in immediately.

Black turfgrass ateniensis—Eggs laid by beetles during May hatch in June and the larvae immediately begin feeding on the turf roots and thatch.

From late June to mid-July, symptoms of injury include wilting in spite of irrigation. In July, larvae move deep into the soil, pupate and emerge as adults. These adults lay eggs during August producing a second generation in states such as Ohio. The second generation larvae are capable of damaging turf.

If preventative applications of insecticide were not made, existing infestations may be spot or generally treated with Proxol, Turcam, diazinon, or Mocap at label rates.

Sod webworms—Damage from sod webworm larvae occurs occasionally in most of the cool-season turf region. Injury is more common in mid-western states and is usually seen in July and August. Older sod fields and heavily thatched turfs are good candidates for infestation. There are generally one or two generations per year, depending upon the species.

In warm season areas most sod webworms complete at least three generations a year with overlapping

During September, billbug adults that developed from summer larvae are often seen wandering about on paved surfaces.

generations toward the end of the season.

Damage is most severe from late June through August. In southern Florida where the tropical sod webworm is active throughout the year, damage is most severe in late summer and fall.

Hybrid bermudagrasses are favored by sod webworms, but damage occurs on other warm season grasses. Webworm damage to bermudagrass often superficially resembles symptoms of some diseases. Flushes of soapy water can be used to determine the presence of sod webworm larvae.

Insecticide applications should be made when larvae are present and/or one to two weeks after peak moth flight.

Formulations of Dursban, diazinon, Sevin, Proxol, or Aspon applied at labelled rates provide control. Retreatment may be necessary depending upon the location and number of generations.

Cutworms—Cutworm larvae continue to cause possible damage to golf course greens from June through August. These larvae pupate in the soil or thatch and emerge as moths that lay eggs for additional generations.

Cutworm larvae can be controlled with a wide range of insecticides label rates; including Dursban, Proxol, As-

pon, Sevin, and others. Irrigation following liquid applications are generally not advisable unless required on the label.

Fall armyworm—The fall armyworm is seldom a problem of cool-season turf.

But in the South, summer always means the arrival of the moths of this migratory pest. Although in mild winters fall armyworms may overwinter among the Gulf Coast, it is generally believed that the moths are blown in on winds from Central and South America. Several generations occur each season, one about every five weeks. Generations overlap in the fall.

Lush, green bermudagrasses are preferred. By late June, fall armyworm damage to turf has usually been reported along the Gulf Coast. Damage is seldom permanent, unless drought and/or heat stress follow.

Fall armyworms may feed anytime during the day but are most active in the early morning and late evening.

Treatment is most effective at these times. During hot, mid-day hours, larvae may retreat into the thatch.

Insecticides such as diazinon, Sevin, Dursban, and Proxol can be used at labelled rates to control fall armyworm.

Greenbug—Damaging populations of greenbug can occur from June through August. Populations and incidents of damage frequently varies from area to area, even within a city.

Symptoms of injury include turf under the dripline of trees and in open areas having a burnt orange color. When symptoms are seen, numerous aphids (40 or more) may be seen on a single grass blade. Close examination of damaged turf is necessary because the aphids are small. If left untreated, a heavy infestation can kill the turf.

Greenbug infestation may be controlled with liquid treatments of Dursban (1 lb. AI/acre), diazinon (2-5 lb. AI/acre), or Orthene at labelled rates. If reinfestation occurs following treatment with Dursban or diazinon, Orthene at labelled rates has been effective.

Fire ants—Fire ants are more difficult to control during hot, summer days because they are deeper in the soil. However, during rainy periods, they may become active and establish new mounds. Treatments during these months should be applied early in the morning before the heat of day.

Scale insects—Although Rhodgrass scale is present in Gulf Coast areas throughout the year, damage be-

continued on page 42



INSECT CONTROL GUIDE

comes most pronounced during the hot, dry days of summer. Bermudagrass and St. Augustinegrass are preferred hosts, but other grasses are also infested. Several treatments with diazinon and a wetting agent are required for control to be effective.

Ground pearls are scale insects that live in the soil throughout the year, sometimes 8-10 inches deep. In the spring eggs hatch producing nymphs that feed through out the summer by piercing turf roots and extracting plant fluids.

Chemical control for ground pearls has not been effective at any time of year. Damage is most severe during summer months when the turf is stressed from heat and drought.

Centipedegrass is especially susceptible to damage, particularly when weakened by overfertilization or drought. Proper fertilization, disease control, and adequate irrigation to maintain healthy turf is the best defense.

Summary of grub control tests in Ohio - 1971-84¹

Insecticide	Rate Lb AI/A	No. tests () ² and avg. % control
bendiocarb (Turcam) 76W	2.0	(9) 82%
ethoprop 5G (Mocap)	5.0	(5) 89%
ethoprop 5G (Mocap)	10.0	(5) 95%
ethoprop 10G (Mocap)	5.0	(3) 83%
ethoprop 10G (Mocap)	10.0	(2) 99%
trichlorfon (Proxol) 80S	8.0	(29) 73%
isazophos (Triumph) 4E	2.0	(8) 82%
isazophos 1G (Triumph)	2.0	(2) 99%
isofenphos (Oftanol) 2I	2.0	(12) 90%
isofenphos (Oftanol) 5G	2.0	(17) 86%
isofenphos (Oftanol) 1.5G	2.0	(7) 86%
diazinon 2G	5.5-6	(7) 66%
diazinon 5G	5.5-6	(2) 49%
diazinon 50WP	5.5	(1) 69%
diazinon 4EC	5.5-6	(20) 63%
diazinon 14G	5.5-6	(5) 29%
carbaryl 80S	8.0	(2) 63%
chlorpyrifos 0.5G (Dursban)	4.0	(8) 33%
chlorpyrifos 2EC (Dursban)	4.0	(5) 64%
chlorpyrifos 4EC (Dursban)	4.0	(6) 51%

¹ - Includes Japanese beetle, *Cyclocephala* spp., *Phyllophaga* spp., only.

² - Each test replicated 3 or 4X. Generally, treatments applied late summer or early spring. Irrigated (1/4-1/2 inch) after application, thatch - 1/2 inch, readings taken ca. 4-8 weeks after treatment. H. D. Niemczyk & K. T. Power.

Fall (Sept.-Oct.)

Chinchbugs—In the northern U.S. the second generation of chinchbug is at peak numbers in September. Nymphs complete their development to adults in late October. Most chinchbugs overwinter in the turf, but some move to protected areas before winter.

Generally, infestation levels at this time are not high enough to warrant the use of insecticides. Early fall rains and infection by a parasitic fungus (*Beauveria* spp.) usually provides sufficient control.

Damage by southern chinchbug may continue in untreated areas. Late summer applications of insecticide usually make fall treatment unnecessary.

Billbugs—During September billbug adults that developed from summer larvae are often seen wandering about on sidewalks, driveways, or other paved surfaces. Before winter, these adults seek shelter in thatch, along sidewalk edges, or near foundations and overwinter there. Many, if not most, overwinter in turf.

In some areas (Cincinnati, Ohio) a partial second generation is known to occur. Larvae of this generation have been known to cause visible damage in September and October. If necessary, diazinon, Turcam or Proxol may be used to control damage.


Grubs—Most species of grubs are in the third of their three stages of development and are feeding actively. When soil temperatures decrease in late October the larvae burrow deeper into the soil to overwinter. Severely cold winters have little effect on survival.

Treatments of existing grub infestations can be accomplished as late as early to mid-September, using standard grub insecticides. Treatment after this time may or may not kill the grubs before they move deeper into the soil to overwinter.

If the soil is dry, irrigation before treatment is advisable. Whenever treatment is applied, the grubs should be in the top one to two inches of soil.

Black turfgrass ateniensis—By Sep-

MARCO: A Lot of Sprayer in a Portable Unit!



D-8-100

Make applications of herbicides, pesticides and fertilizer quick and easy with a MARCO sprayer! MARCO's Prime Line of more than 20 models of spraying equipment features one-piece, hand-laid and baffled fiberglass tanks in custom sizes - each with a 20-year warranty!

Call or write for a brochure illustrating our complete line. We can custom build to fit your needs, too.

Contact MARCO today - and put our fully professional, fully affordable sprayers to work for you!

Here's a model that will meet the majority of your requirements:

Model D-8-100 Specifications

Pump Cap. 14 GPM
 Pressure 550 psi
 Power 8HP Briggs & Stratton Engine
 Tank Cap. 100 up to 400 gallons
 Standard Agitation Jet
 Pressure Control Unit ... 15 GPM
 Standard Drive Direct

Options

Turf Trailer Mounting
 Belt-Driven Mechanical Agitation
 Fill Well Basket Filter
 Canopy and Hose Reel to Hold 200' of 1/2" hose
 Dry Booms in Varying Lengths
 Boom Jet - Pistol
 Grip Gun
 52' or 72' Hand Boom and Drift Skirt



DIVISION OF LLO-MAR MFG. CO.

(800) 527-1574
 In Oklahoma, call (405) 362-3162

210 S. Main St.
 Newkirk, OK 74647

SEE MARCO'S NEW 1986 PRODUCTS AT BOOTH 940, GCSAA!

Circle No. 130 on Reader Inquiry Card

TURF VAC®

Only sweeper for both turf and pavement. Grass, glass and trash... wet or dry. Sweeping widths from 4 to 10 ft. Self-propelled and tow-type, both with ground dump and power lift dump.



Riding Model 80



Model FM-5 with hand intake hose

SANDANCER®

Versatile multi-purpose work vehicle. Hill-climbing 18 h.p. gas engine with a unique power train that eliminates stripped gears and burned-out clutches. Three models and a wide selection of options allow you to build a SANDANCER to match your work needs.



3 or 4 wheel models

Write for brochures and distributor in your area.



TURF VAC CORPORATION

P.O. Box 90129
 Long Beach, CA 90809
 (213) 426-9376

Circle No. 156 on Reader Inquiry Card



tember, adults of the current generation begin to fly into protected areas, such as golf course roughs, to overwinter. Larvae that have not completed development to adults before frost are killed.

Mole crickets—Mole crickets fly again in the fall, but no egg laying is known to occur at this time. The crickets are large and difficult to control at this time. Damage becomes obvious as turf growth slows.

Sprays of diazinon (5-6 lb. AI/acre) or Turcam (2 lb. AI/acre) may have to be repeated several times. Oftanol (2 lb. AI/acre) may work too slowly for adequate control of large crickets in October. Mocap 10G (10 lb. AI/acre, commercial turf only) is usually effective at this time providing up to four weeks residual control.

Sod webworm—Northern sod webworm larvae are small and cause little if any damage in the fall. Late in the fall the larvae construct a cocoon-like shelter in which they overwinter.

Except for the most southern areas where development is continuous, sod webworm larvae present in September will overwinter. Areas treated earlier in the season may be reinfested by this time. Treatment in September reduces the population for next season.

Fall armyworm—Fall attacks on newly established turf from mid-September through October may result in damage that will not recover with fall fertilization. This forces the turf to enter winter in a stressed condition. Such damage can contribute to winter turf mortality.

If needed, apply controls early in the morning or late in the day when fall armyworms are most active. Use diazinon, Sevin, Dursban, or Proxol.

Fire ants—Hot, dry periods in September and October may make fire ant control difficult. Once rain begins, fire ants become active and may be effectively controlled with mound treatments of diazinon, Dursban, Orthene, Amdro bait or MC-96. Larger infested areas where mound treatment is impractical can be treated with Amdro fire ant bait (1.5 lb./acre).

Greenbug—Severe infestations of greenbug have been known to occur as late as the first week of December. Areas having a history of infestation should be reexamined when mild temperatures extend late into the fall. Heavily infested turf will not survive through winter.

Late fall infestations may be controlled with the same insecticides used to control the pest in the summer. **WT&T**



WOODY ORNAMENTAL INSECT CONTROL

by Dr. D.G. Nielsen and Dr. J.R. Baker

Landscape trees and shrubs constitute a major investment and add significantly to the beauty and value of residential and commercial properties.

Landscape managers need to be familiar with woody plants, their requirements for survival and vitality and their pest problems to implement effective pest control programs. Most native trees and shrubs on undisturbed sites suffer only rarely from ravages of insect pests.

However, trees growing in landscapes are commonly stressed by construction damage, lack of water (or too much water if there is poor drainage), high temperatures, compacted soils, and other factors that reduce the tree's ability to either repel or withstand insect attacking without suffering decline.

Many major pests of trees and shrubs are probably opportunists that exploit hosts that have been weakened by physical factors.

This article is designed to help landscape managers develop strategies for controlling insect pests of woody plants.

The information is organized according to seasonal insect activity. The time or times an insect is vulnerable to a direct control tactic and up-to-date insect control recommendations are provided.

Control Strategies

In the past, pre-scheduled cover sprays were often used to control insect pests that may be present on the property, because the detrimental side effects of some pesticides were not yet known and landscape managers were not familiar enough with local pests to develop target spray programs.

Many times, all trees on a property were sprayed when only a few harbored a pest species. Today, conscientious tree care specialists use

Dr. D.G. Nielsen is a professor of entomology at Ohio State University, and Dr. J.R. Baker is an extension entomologist at North Carolina State University.



The life cycle of the black vine weevil (from left): egg, larva, pupa, adult. Four foliar sprays at three-to-four week intervals are required for control.

insecticidal sprays only after determining which pest(s) is present and whether or not it has reached a potentially damaging population level. They spot spray only infested plants.

This approach is cost effective in terms of material costs, landscape beauty and longevity, and environmental quality.

Dormant Period (November-March)

Some insect and mite pests are vulnerable to control tactics after trees and shrubs have become acclimated to winter temperatures. Landscape managers can capitalize on this vulnerability by performing pest control functions when other maintenance activities are not competing for their time.

Horticultural spray oils can be used safely on many woody plants to control overwintering eggs of spider mites and aphids and immature forms of adelgids and soft scales.

New information from Cornell University indicates that currently

available horticultural oils can be used throughout the year when the temperature is above freezing.

However, four cautions should be considered before using these products:

- Read the label to make sure the product is not phytotoxic to plants on which you intend to use it;
- Do not use oils at the dormant season rate in the fall before plants have become winter-hardy. In the South, woody ornamentals may not go dormant until late December;
- Oil sprays should not be used on plants under moisture stress or when temperatures are high with high humidity;
- Do not apply oil sprays to tender new growth in spring.

Spruce Gall and Pine Bark Adelgids

Adelgids are small, soft-bodied sucking insects that feed on the needle or bark of conifers.

On spruce they cause pineapple-like galls to form on the new growth.

Eastern spruce gall adelgid and Cooley spruce gall adelgid overwinter



as young nymphs on the undersides of branches on spruce or on the lower leaf surface of Douglas fir, the alternate host of the Cooley gall adelgid.

Pine bark adelgid overwinters as eggs, nymphs, and adults in bark cracks and crevices on white pine.

By late winter the trunks of infested pines may be snow-white from huge numbers of adelgids.

Adelgids can be controlled on all hosts any time after spruce galls open in late summer until just prior to bud break the following spring.

Horticultural oil may reduce the overwintering population, but oil may remove the waxy bloom from the needles of spruces and pines, causing them to become unsightly.

Sevin (carbaryl) or lindane can also be used in a thorough-coverage, hydraulic spray, making sure to cover the underside of spruce branches and Douglas fir needles. A wetting agent may be useful to help penetrate the fluffy wax covering that surrounds the insect, especially when treating the pine bark adelgid.

Mites

Spider mites, including spruce spider mite on coniferous evergreens and southern red mites on broadleaf evergreens, suck plant juices and deposit silk and waste material on their hosts, causing foliage to become dull and bronze-colored.

Mites tend to feed in spring and fall but populations usually crash in very hot or very cold weather. They overwinter in the egg stage which is susceptible to control with horticultural oil used at the dormant season rate.

False spider mites are tiny, flat mites which feed and reproduce slowly throughout the year. Most "winter injury" symptoms on Aucuba, azalea, and Japanese hollies are due to false spider mites.

Horticultural oils or other miticides give adequate control of these pests. Treatment is appropriate at any time of the year except in early spring when new, tender growth is emerging.

Aphids

Aphids are small, soft-bodied insects that insert their mouthparts into the phloem of leaves, stems, and roots to suck out sap.

Aphids excrete honeydew, a sweet liquid that coats heavily infested plants. Some species (melon aphid, apple aphid) feed on the most succulent part of the plant. Other species (giant willow aphid, giant bark aphid, *Cinara* aphids) feed on stems. High aphid populations can cause leaves to



Fuller rose beetles notch the leaves of shrubs in the South.

yellow and fall prematurely.

The honeydew they excrete serves as a substrate for a black sooty mold fungus that reduces the aesthetic appeal of the plant and reduces its ability to manufacture food.

Many aphids overwinter as exposed eggs on conifer needles or on stems and buds of other woody plants.

If a damaging aphid population was detected the previous summer or fall, an application of horticultural oil before bud break will reduce the spring aphid population, thereby giving the tree a chance to recover before aphids build up again during the spring and summer.

Soft Scales


Soft scales are another kind of sucking insect that drain a tree's energy and reduce its ability to manufacture food.

They suck sap from the phloem and produce honeydew. Some (e.g. the tulip tree scale) seriously weaken or kill their hosts.


Heavily infested trees and shrubs often become blackened with sooty molds. Some soft scales overwinter as immature forms (cottony maple, cottony maple leaf, magnolia, Pine tortise, and Fletcher scales) that are somewhat vulnerable to horticultural oils used at the dormant application rate.

Spring (April-late June)

Most insects become active in the



Mount Up, Ride Out.



When you're gunning for the pests who've been robbing your territory of good health and appearance, put Hannay Reels on your side. Or in the back. Or in a cabinet.

Mounted on trucks or trailers, versatile



SERIES 1500

Hannay Reels for pest control give you greater range, better control and neat, convenient hose storage.


The trail home is just as fast and easy, with direct crank rewind or chain and sprocket drive powered by an electric or compressed air motor.

Mount up with Hannay Reels, and ride with the best.

The Reels of Progress



CLIFFORD B. HANNAY & SON, INC. 600 EAST MAIN STREET WESTERLO, NEW YORK 12193 • TELEPHONE (518) 797-3791



Cinara aphids feeding on a juniper in the early spring.

spring, responding to warmer weather and resumption of plant growth and development.

Monitoring trees and shrubs for pests during spring is one of the most important tactics in a modern insect control program.

Newly expanding or expanded leaves should be checked for the presence of sucking insects, leafminers, and defoliators.

Tree limbs and trunks should be inspected to determine presence of active borer galleries as early detection will permit time for learning the identity of the pest and determining if a spray program is justified.

Defoliators

Eastern tent caterpillar, fall cankerworm, whitemarked tussock moth, and pine sawflies are among the first defoliators to begin feeding in spring.

The eastern tent caterpillar is obvious and readily detectable because it forms a silken tent in tree crotches, especially flowering fruit and nut trees.

During years of high caterpillar numbers, entire trees may be defoliated. These pests often reach maturity before defoliation is noticed.

The tent caterpillars and their relatives, including mimosa webworm, fall webworm, bagworm, and gypsy moth, are readily controlled with a number of conventional insecticides or with the bacterium, *Bacillus thuringiensis*, commonly known as B.t.

Pine sawflies are often difficult to detect since they blend-in with needles on their hosts. They often reach

maturity before defoliation is noticed.

Sawflies are related to bees and wasps and are highly susceptible to Sevin. Orthene (acephate) is also labeled for this use.

Elm leaf beetle larvae and adults consume foliage. There are two generations each summer. Sevin, Orthene, or Turcam/Dycarb (bendiocarb) can be used when trees leaf-out in spring. A second generation may require a second application in July.

Armored Scales

Armored scales, soft-bodied sucking insects that suck juices from leaves and stems but do not produce honeydew, are named armored scales because after the first stage molts, later stages are covered by cast skins and tough wax.

Consequently, they are vulnerable to contact insecticides only during the crawler and settled first nymph stages.

Armored scales overwinter as eggs (pine needle and oystershell scale), as mated females (euonymus and white peach scales), or in more than one stage (hemlock and tea scales). As indicated, all of them are most easily controlled with crawler sprays.

Armored scales that overwinter as eggs can usually be controlled with a single application of an insecticide, if thorough coverage is achieved. If application timing is not precise, a systemic insecticide like Metasystox-R (oxydemetonmethyl) or Orthene should be used.

Species like euonymus scale require more than one crawler spray, since the first hatching crawlers molt before the last spring genera-

tion eggs are laid.

Three thorough-coverage, hydraulic sprays at 10-to-14 day intervals are needed to provide an adequate level of control.

All armored scales that have more than one generation per year (pine needle, euonymus and white peach scales) should always be controlled during the spring crawler hatch because the hatching period is shorter at that time, so fewer sprays are required to provide control.

Horticultural oils may be effective for armored scale control. Use the summer rate after new plant growth has hardened-off.

Aphids

Aphid populations can explode in a short time, since a new generation can be produced every 10-to-15 days in the North and even faster in the South.

In the North, aphids are often at high population density during summer droughts, or just after a drought period, and should be controlled before they cause premature leaf drop.

In the South, aphid populations are often high in late winter and early spring before lady beetles and other predators become active. However, crape myrtle aphid populations often become damaging later in the summer.

Mistblower applications are excellent against free-living aphids.

Adelgids

Remember, overwintering forms can be controlled by using a hydraulic application of lindane or Sevin, stressing coverage to the underside of branches and leaves before bud break.

After bud break, adelgids on spruce are protected as their galls form. They become vulnerable again in fall after their galls open. Pine bark adelgids can be controlled in spring or summer as long as the pines are not under water stress and after the new growth has begun to harden off.

Leafminers

Birch, boxwood, and holly leafminers are highly specialized insects whose larval stages damage trees and shrubs by destroying tissues within the leaf.

Birch leafminer is a sawfly (closely related to bees and wasps) who emerge as adults in May.

Foliage can be protected by spraying when the adult sawflies are actively mating and feeding on birch trees.

Sevin, malathion, and lindane are effective before eggs are laid within leaf tissue. After egg laying begins or



INSECT CONTROL GUIDE

INSECTICIDE DIRECTORY

Common Name	Brand Name	Company
acephate	Orthene	Ortho/Chevron
<u>B. t.</u>	Thuricide	Zoecon
	Dipel	Abbott Labs
	SOK Bt	Nor-Am
bendiocarb	Turcam	Nor-Am
carbaryl	Sevin	Union Carbide
chlorpyrifos	Dursban	Dow
diazinon	Diazinon	Ciba Geigy
	Sarolex	Ciba Geigy
dicofol	Kelthane	Rohm & Haas
dicrotophos	Bidrin	Shell
dimethoate	Cygon	American Cyanamid
dioxathion	Deltic	Nor-Am
dymet	Dymet	Mallinckrodt
femobutain-oxide	Vendex	Shell
fenvalerate	Pydrin	Shell
fluvalinate	Mavrik	Zoecon
isofenphos	Oftanol	Mobay
malathion	Malathion	American Cyanamid
methoxychlor	Marlate	Kincaid
morestan	Imidan	Stauffer
oxydemetonmethyl	Metasystox	Mobay
phosmet	Imidan	Stauffer
trichlorfon	Dylox	Mobay
	Proxol	Nor-Am

mines begin to form, a systemic insecticide should be used. Metasystox-R, Orthene, and Cygon (dimethoate) are labeled for this use.

There are several generations per year, but the first two generations seem to be most destructive.

Holly leafminer is a true fly that has only one generation per year. Spring application of Metasystox-R or Orthene after the new plant growth has hardened-off is necessary to achieve control.

Boxwood leafminer is a gall midge which also has one generation per year. A mid-to late-spring application of Cygon will adequately control boxwood leafminer.

Spider Mites

Spider mites such as twospotted spider mite and tumid mite are common and damaging on plants under water stress and during long, hot droughts.

They complete many generations throughout the summer. Kelthane (dicofol), Mavrik (fluvalinate), or an-

other miticide should be used before mites cause foliage to turn bronze. A hydraulic sprayer must be used to maximize coverage, especially on plants with dense foliage, including foundation plantings, conifers, and other evergreens.

Two sprays must be used at a seven-to-10 day interval, since most miticides do not kill eggs. A single application will not be effective against spider mites.

Root Weevils

Root weevils (black vine, strawberry root), can be destructive in both adult and larval stages.

Adults chew notches in leaf margins. Larvae consume small roots and girdle larger roots, sometimes causing death of foundation plants, including rhododendron, azaleas, and yews (taxus).

Spray foliage with Orthene or Turcam/Dycarb in mid-June, followed by repeat applications at three-to-four week intervals until August.

Level of control is directly related

to the degree of coverage, so use a hydraulic sprayer to control root weevils. Drenching soil beneath host plants may help reduce larval populations.

Borers

Clearwing moth borers are common in lilac, ash, dogwood, rhododendron, oak, and flowering cherries. Flatheaded borers (adults are called metallic wood borers) are common in white-barked birches, oaks, and other stressed hardwoods.

Larvae do the damage by feeding beneath bark, disrupting movement of food and water, destroying the cambium (the growth layer of cells), and causing structural weakness. Clearwing presence and flight periods can be monitored with pheromone traps (see insert).

A single, thorough-coverage bark spray of Dursban (chlorpyrifos) or lindane, 10-to-14 days after first male moth capture, will provide season-long control of most clearwing moths.

Three applications of bark/foilage sprays with Turcam/Dycarb, Dursban, or lindane are required to control flatheaded borers.

Summer (July-Sept.)

Defoliators

Mimosa webworm, bagworm, fall webworm, Japanese beetle adults, and second generation elm leaf beetles sometimes become common in early summer.

All of these pests should be controlled when larvae are small to minimize damage and maximize effectiveness of the insecticide. Caterpillars can be controlled with one of the B.t. formulations. Sevin, Orthene, Turcam/Dycarb, and several other common insecticides will also control these pests.

Mistblower application may be cost-effective but may result in unacceptable drift of insecticidal sprays, especially in windy weather.

Japanese beetle adults defoliate many kinds of woody plants and roses in July and August. They are most easily controlled with weekly sprays of Sevin or Turcam/Dycarb.

Mavrik also gives long-term control of Japanese beetle adults. Japanese beetle traps can be used to capture large numbers of beetles, but they do not reduce defoliation or control the beetle population.

Area-wide grub control is the most effective way to reduce numbers of Japanese beetle adults and defoliation they cause.

Second generation elm leaf beetles



can cause significant defoliation if heavily infested trees are not sprayed. Birch leafminer can be controlled, if trees are sprayed when second or third generation adults are mating and ovipositing.

Scales

Crawlers of several soft scales (Fletcher, cottony maple, cottony maple leaf, pine tortoise, wax and tulip tree) hatch in late June or early July (earlier in the South).

They and settled nymphs are susceptible to scalicides (Sevin, Orthene, diazinon, Dursban, Turcam/Dycarb and Mavrik) in early July. A single, thorough-coverage, hydraulic spray should provide control.

Settled nymphs and other nymphal stages are vulnerable to contact insecticides, because they are not protected by cast skins and wax. Sprays to control soft scales should always be applied after all eggs have hatched to minimize the impact of pesticides on lady beetles and other predaceous insects, to minimize insecticide usage, and to maximize control.

Second generation pine needle scale, euonymus scale, and white peach scale crawlers hatch during July and early August. Two sprays at a 10-day interval may be required to control pine needle scale and white peach scale because crawlers hatch over a three week period.

Summer generation euonymus scale crawlers hatch over a longer period, so three applications at 10-day intervals are required.

Aphids

Aphid populations should be controlled before they secrete copious amounts of honeydew or do irreversible damage to leaves.

If aphids are allowed to build-up in high numbers, plant growth may be distorted and leaves may fall prematurely. Once honeydew and sooty mold are present they may persist long after aphids have been controlled by pesticides or natural enemies. Aphids are vulnerable to contact sprays whenever they are active.

Spider Mites

Spider mites can be controlled whenever they are active by spraying twice with a five-day (South) or 10-day (North) interval.

If trees are receiving repeated applications of Sevin to control other insects, be especially watchful for build-up of spider mites. Sevin selectively kills natural enemies of mites, thereby contributing to increases in spider mite populations.

Woody ornamentals—such as

some euonymus varieties, roses, and bedding plants—are frequently infested by twospotted spider mites and tumid spider mites. These mites are dispersed on air currents and may appear suddenly in large numbers in hot dry weather.

Root Weevils

The second and third applications of black vine weevil adulticides should be applied in July and August.

In the South, Japanese weevils and Fuller rose beetles can be controlled with Orthene as a spray and drench during July. A single spring application will not control black vine weevil or other weevils mentioned earlier.

Area-wide grub control is the most effective way to reduce numbers of Japanese beetle adults...

Borers

White-barked birches determined to be infested by bronze birch borer during the summer can be injected with Inject-A-Cide B (Bidrin) using microinjection procedures developed by the J. J. Mauget Company.

Injection must be done by a trained technician between early July and early August but should not be used as an annual, preventive tactic.

Infested trees should be watered weekly during summer and fall drought and fertilized in the fall after the first hard frost.

The following year, bark/foilage sprays should be implemented as indicated earlier.

The peachtree borer can be controlled with a single application of Dursban or lindane in early July (in the North) or late August (in the South). The second application for control of lesser peach tree borer should also be applied at this time to infested flowering cherries.

Fall (Sept.-Oct.)

Defoliators

Mimosa and fall webworms reach their highest population density and cause most defoliation during late summer and early fall. They should be controlled as soon as first generation larval webs are detected in early summer.

However, both pests are susceptible to larvicides in late August and early September. If B.t. is to be used, it must be applied when larvae are small to achieve an acceptable level of control.

Scales

Magnolia scale and tulip tree scale crawlers are produced in late August and early September. Infested magnolias should be sprayed when goldenrod is in full bloom (early September).

A single, thorough-coverage, hydraulic spray with Orthene or Sevin will provide excellent control. Magnolias and tulip trees may be severely stunted or even killed by heavy infestations of these scales.

Gall Adelgids

Galls on spruce caused by eastern and Cooley spruce gall adelgids turn brown and open in August and September. After galls open, adelgids are vulnerable to contact insecticides. Adelgids on spruce, Douglas fir, and pine remain vulnerable to insecticidal sprays until the following spring when new buds open.

Root Weevils

Attempts to control root weevil larvae should be made in early September and early October. Two drenches with Turcam/Dycarb have been effective against larvae established in soil surrounding roots of field plants.

A single drench with Turcam/Dycarb, Orthene, or Furadan (carbofuran) controls larvae infesting containerized plants. Recent evidence indicates that overwintered larvae may also be susceptible to drenches in early spring.

Closing Thought

We believe that the best way to minimize insect damage is to maintain healthy trees and shrubs. Trees should be matched to sites, watered during the first two years after planting and during summer and fall drought, pruned properly, fertilized in late fall, mulched, and aerated to promote root vitality.

Adherence to these basic horticultural practices will promote tree vitality and improve their ability to withstand attack by most insect pest species.

Throughout this article we have stressed the importance of pest identification, proper timing, and thorough coverage for achieving a high level of insect control.

Coverage and timing are often more important than the insecticide or miticide used. So, make sure of proper pest identification, determine when it is most vulnerable to control, and apply a pesticidal spray thoroughly to only infested trees following label directions. **WT&T**