

Drought tolerance of turfgrass

■ Landscapers, lawn care operators and golf course superintendents must often maintain acceptable turf with limited amounts of water. This is especially true in arid and semi-arid regions. Various approaches are used to conserve water, including the use of grasses that need only naturally available water and those that can be managed with reduced irrigation.

Many grasses can be grown in dry areas of the western United States without supplemental water. However, most of these are not well suited for finely-manicured areas because they grow too tall or produce a bunched and open turf.

In cool areas with good soils and at least 15 inches precipitation per year, drought-tolerant Kentucky bluegrass cultivars, once established, will usually persist, and produce acceptable turf in the spring and fall. Under dry conditions, bunch-type turfgrasses, including perennial ryegrass and hard fescue, often show exceptional short-term drought tolerance. However, with extended drought, these

Species selection is the key to healthy, low-water-use, attractive lawns and landscapes.

species tend to thin and do not provide acceptable turf quality.

Among the drought-tolerant grasses that can do well on golf course fairways and roughs are buffalograss and bermudagrass, both low-growing, sod-forming, warm-season species. For roughs and out-of-play areas, several additional grasses—including blue grama and inland saltgrass (warm-season species), and western and fairway wheatgrass, smooth brome grass and Russian wild ryegrass (cool-season species)—are fairly well adapted to limited supplies of water. Continued research may provide the refinements which will allow these grasses to become more widely used.

Another management approach is to

use limited irrigation. Watering in the summer to supply about 70 to 80 percent of potential evapotranspiration (ET) will normally provide acceptable turf from cool-season grasses.

The frequency of irrigation is also important. For example, irrigating Merion Kentucky bluegrass at two- and four-day intervals produces better turf than the same amount of water (50 to 75 percent of potential ET) applied at 14-day intervals.

continued on page 32

Pre-conditioning lawns

■ A lawn that is deprived of moisture for 45 days will typically suffer a 20 percent loss, and it will most likely turn a dormant straw color, but it probably isn't dead. After several days of moderate temperature and moisture, re-growth will begin, with visible results about two weeks after the dormancy is broken. It may take three to four weeks for severely drought-stricken lawns to fully recover, but they usually will, in time.

Here is how landscapers, golf course superintendents, lawn care operators and even homeowners can pre-condition their lawns for drought:

- Apply recommended amounts of turf fertilizer in the spring.
- Mow with sharp blades. Dull blades create more stress on the plant and cause it to require more moisture.
- Raise mowing height and reduce mowing frequency. Greater leaf surfaces allow the plant to store more water, and the leaves shade the rootzone from drying heat and evaporation.
- Water infrequently but deeply in the mornings or evenings to establish deeper rooting systems. To determine when to water, probe the soil to four to six inches deep; when the soil is dry or the probe is difficult to insert, it's time to water.
- Avoid excess traffic, which causes more stress.

—American Sod
Producers Association

Watering instructions

- Make sure your clients know how to save water while keeping their lawns looking their best. Here are some suggestions, whether in a hot, dry spell or not:
 - Water early in the morning to reduce evaporation.
 - Water the lawn separately from trees, shrubs and groundcovers, if possible.
 - Have thatch removed in spring if it's more than one-half inch thick. Thatch should not be removed in the heat of the summer.
 - Control weeds. They compete for water, light and nutrients.
 - Fertilize moderately, applying at the low end of recommended rates.
 - Keep your lawn at the right mowing height, a minimum of 1.5 inches for tall fescue, perennial rye and Kentucky bluegrass, a minimum of 0.5 inch for bermudagrass, zoysia-grass and St. Augustinegrass.
 - Have the lawn aerated as necessary to prevent compaction and to allow water to move more freely into the soil. Clay soils in particular need regular aeration.

—University of California

Drought-tolerant varieties

■ Select grasses according to the nature of the area to be covered, water availability and effect desired. For any of the lawn grasses to produce an aesthetically pleasing turf, supplemental water must be applied. Generally, grasses should be considered high water requirement plants, but drought tolerant turfgrasses are becoming increasingly available.

Here are National Turfgrass Evaluation Program (NTEP) ratings of various commercially-available varieties, grouped by cultivars. These are the highest ratings in each group of trials, with no statistical differences evident. Please note that most ratings are based on limited site evaluations. All are from NTEP 1992 progress reports.

BUFFALOGRASS (wilting)		Morning Star.....8.0	Merit.....4.0
315.....9.0		Navaho.....8.0	Opal.....4.0
609.....9.0		Quickstart.....8.0	Ram-1.....4.0
Bison.....9.0		Target.....8.0	
Buffalawn.....9.0		Topeka.....8.0	KENTUCKY BLUEGRASS
NE 84-378.....9.0		Prizm.....8.0	MEDIUM-HIGH MAINT.
Sharps Improved.....9.0			(dormancy)
BUFFALOGRASS (dormancy)		KENTUCKY BLUEGRASS MEDIUM-HIGH MAINT. (wilting)	
609.....8.2		Eagleton.....8.7	Barzan.....5.0
Texoka.....7.8		Barmax.....8.3	Glade.....5.0
315.....7.2		Monopoly.....8.3	Ronde.....5.0
Plains.....7.2		Silvia.....8.3	Indigo.....4.7
Bison.....7.0		A-34.....8.0	Marquis.....4.7
Sharps Improved.....7.0		Indigo.....8.0	Merit.....4.7
		Blacksburg.....7.7	Viva.....4.7
PERENNIAL RYE (dormancy)		Challenger.....7.7	
Nighthawk.....8.7		Classic.....7.7	KENTUCKY BLUEGRASS
Patriot II.....8.7		Freedom.....7.7	LOW MAINTENANCE
Pebble Beach.....8.7		Nustar.....7.7	(recovery)
Sherwood.....8.7		Preakness.....7.7	Monopoly.....6.3
Affinity.....8.3		Suffolk.....7.7	Banjo.....5.0
Envy.....8.3			Alene.....4.7
Stallion Select.....8.3		KENTUCKY BLUEGRASS LOW MAINTENANCE (dormancy)	
Accolade.....8.0		Fortuna.....5.0	S. Dakota Cert.....4.7
Achiever.....8.0		Voyager.....5.0	Barzan.....4.7
Advent.....8.0		Merion.....4.7	Kenblue.....4.3
Barrage.....8.0		Unique.....4.7	Nustar.....4.3
Caliente.....8.0		Banjo.....4.3	
Brightstar.....8.0		Amazon.....4.0	TALL FESCUE
Competitor.....8.0		Barzan.....4.0	(recovery)
Cutter.....8.0		Bronco.....4.0	Hubbard 87.....6.3
Gator.....8.0		Chelsea.....4.0	Phoenix.....6.0
Goalie.....8.0		Destiny.....4.0	Adventure.....5.7
			Chieftain.....5.7
			Guardian.....5.7
			Jaguar II.....5.7
			Monarch.....5.7
			Sundance.....5.7
			Willamette.....5.7
			Winchester.....5.7

Athletic field soil is a key to avoiding injuries

by Henry Indyk, Ph.D.

■ Consider athletic field soil conditions from two major perspectives: the best possible growth situation for the turf, and the basis for player safety.

For optimum turf growth, soil conditions must be suitable from both chemical and physical standpoints.

Chemical conditions include pH, nutrient status, level of salt concentration and contaminants.

Physical conditions include soil texture, infiltration and percolation, drainage, and susceptibility to compaction. From the standpoint of safety, add field grade and contour, evenness or levelness, existence of

Modified topsoil root zone mix. A 10-inch layer is spread over the subgrade showing one of the drainage lines.



depressions, and undesirable debris.


Assessing conditions—Know what's wrong before you take any action. Soil conditions must be assessed to analyze existing situations properly. This must be done by someone with knowledge and background in soils, someone who knows what to look

for and how to judge conditions accurately.

A great deal can be determined by visual observation during a site visit: physical conditions including soil texture, drainage, levelness, contour, grade, depressions, percentage of debris and existing turf condi-

continued on page 34

Roundup® points the way to savings - both in money and manpower.



When you treat an area with Roundup® herbicide, it takes less time than hand weeding or string trimming it. You also get complete control — not just burndown. Add in its low price and Roundup gives you a big hand in making the most of your money.

Roundup
Herbicide by Monsanto

Get your FREE kit, "Weeds Are No Longer Our Only Concern." Call 1-800-332-3111.

IND-47080 2/94

Circle No. 123 on Reader Inquiry Card

ATHLETIC SOIL from page 32

tions (density, evenness and vigor). Even hardness can be determined to some degree, though for best accuracy a physical measurement of hardness can be taken with an impact machine.

The site visit includes soil sampling for a laboratory quantitative assessment of chemical and physical characteristics. Take soil samples according to field variability, pulling some samples from each area with variations. From 10 to 20 samples should be taken. Gather them with a soil sample tool to a depth of at least six inches.

Combine and thoroughly mix the collected samples in a clean container. By testing the samples together, rather than testing each sample individually, results will reveal a composite of the entire field.

Send air-dried samples to a qualified, reputable laboratory for chemical testing and physical analysis.

Chemical testing should include levels for soil acidity (pH) and the major nutrients of phosphorus and potassium, and calcium and magnesium. If the field is in an area where micronutrients are normally deficient, test for these also.

Physical analysis should include mechanical analysis for sand, silt and clay proportions. This will help predict the ability of the soil to infiltrate and percolate water, its drainage characteristics, water-holding capacity, and its compaction susceptibility. Information gained from this analysis can be fortified with other laboratory determinations, such as sand

fractionation, bulk density and percolation rate.

Visual assessment and lab test results combined provide useful information on the existing status of the field.

Corrective procedures—It's vital to know what the soil profile conditions should be for the growth and support of a dense, vigorous and healthy natural turf sports playing surface in order to determine what needs to be done to get there.


Corrections in the chemical area will be based on the results of the laboratory analysis and are generally issued in prescription form by the lab. For example, use v amount of x material to alter the y level by z amount.

The proper pH levels for favorable turfgrass growth range between 6.0 and 7.0 with 6.5 being ideal. High pH—alkaline conditions—require acidifying; low pH—acidic conditions—must be neutralized. Sulphur compounds correct alkaline conditions; lime corrects acidity. The farther pH levels range from the ideal, the harder it will be to correct pH conditions for the proper establishment and maintenance of turfgrasses.

Lab prescriptions also spell out the amount and type of corrective materials to be applied according to specific nutrient deficiency levels.

In spite of pH and nutrient levels, high salts or other contaminants can affect turf growth. Contaminants may be due to misapplication of chemicals, or the result of leaks or spills. Excess salts may also come from contaminants. They may build up in coastal regions when areas are inundated with brackish water during flooding. In the Southwest, natural salt levels may be great enough that salt crusts appear on the soil surface.

Roundup® points the way to environmental stewardship.



The benefits of Roundup® herbicide extend beyond traditional trim and edge work. It's also being used to restore wildlife habitats. So when it comes to treating sensitive sites, Roundup gives you a big hand.

Roundup
Herbicide by Monsanto

Get your FREE kit, "Weeds Are No Longer Our Only Concern." Call 1-800-332-3111.

IND-47081 2/94

Circle No. 123 on Reader Inquiry Card

Roundup® points the way to application ease.



Learning how to properly use Roundup® herbicide is a snap. For even more convenience, try new Roundup Dry Pak. Either way, Roundup gives you a big hand in simplifying worker training.



Get your **FREE** kit, "Weeds Are No Longer Our Only Concern." Call 1-800-332-3111.

IND-47082 2/94

Circle No. 123 on Reader Inquiry Card

range (1.0 mm) down to the fine range (0.25 mm). Little—10 percent or less—of the sand should be in the range above very coarse or below very fine.

Depending on available sand sources, silica sand is preferred to calcareous sand, which normally carries a high pH.

Drainage problems—One can go through all the steps of rebuilding the soil, doing everything ideally, so that the soils are chemically and physically on target, grade the field, sod or seed, and have the field look beautiful until the first big rain. Then problems may appear: what happened with all of the expended energy and cost in rebuilding the soil?

Modification of the upper layer of soil, no matter how great the depth, is not the solution to a drainage problem unless the subsoil is such that it allows drainage, which is a rare solution. Essentially, what is above the subsoil, with good physical characteristics that allow percolation and infiltration, is then impeded by the subsoil barrier which causes water to back up.

An internal soil drainage system is needed to correct subsoil drainage problems, allowing the modified field media to function properly.

Rebuilding the soil requires "in depth" homework. Know what you have, where you want to go, and what you need to get there—and test every step of the way.

—*Dr. Henry Indyk is turfgrass agronomist for Turfcon, the professional consultation branch of The Greenway Group, Horsham, Pa. He was extension specialist with Rutgers for 30 years. Indyk is secretary of the national Sports Turf*

Salt concentration is measured easily with a conductivity test. Using water to leach out excess salts can be a long, time-consuming process. The water to be used also must be tested for salt levels. Gypsum can be used to accelerate the leaching process. In some cases where high salt levels exist, it may be necessary to replace existing soils. The new soils must be tested also, before changes are made.


Though a visual assessment gives a qualitative analysis of physical conditions (the sand:silt:clay ratio), it should always be backed up with a laboratory mechanical analysis to establish quantitative proportions. In addition to this mechanical analysis, a sieve analysis, also known as a sand fractionation analysis, should be conducted.

The sieve test separates and grades sand particles by specific sizes: very coarse, coarse, medium, fine and very fine. If the sand component is predominantly in the fine/very fine range, the small particles can fit together so closely that the sand functions more like silt and clay and their restriction on the rapid infiltration and percolation of water required for suitable athletic field soils.

Corrective procedures to improve filtration, percolation rates and compaction resistance require soil modification to change the physical make-up of the soils. To be effective, these modifications must incorporate the right amount of the right quality of sand. The critical factor is testing of the sand for particle size, pH, and salt index prior to use. There's no point bringing in sand that has problems of its own.

Stipulate sand fractions with 90 percent in the coarse

Roundup® points the way to beautiful results.



Used properly, Roundup® gives you a big hand in achieving long-lasting control that blends in with your surroundings.



Get your **FREE** kit, "Weeds Are No Longer Our Only Concern." Call 1-800-332-3111.

ALWAYS READ AND FOLLOW LABEL DIRECTIONS FOR ROUNDUP HERBICIDE. Roundup® is a registered trademark of Monsanto Company. © Monsanto Company 1994 IND-47083 2/94

Circle No. 123 on Reader Inquiry Card