

TABLE 2. CONT.

Common name	Trade Names	Uses
MSMA	Crabgrass Killer (Bonide); Daconate 6; Daconate Super (ISK Biosciences); Drexar 530 (Drexel); MSMA (Bonide); LESCO; MSMA Turf (Turfgo/United Horticultural Supply); 912 Herbicide; 120 Herbicide (Riverside/Terra International); Super Crabgrass Killer (Rockland); Weed Hoe (Monterey)	Selective post-emergence control of annual grasses.
MSMA+cacodylic acid	Broadside; Moncide (Monterey)	Selective post-emergence control of annual grasses.
sethoxydim	Vantage (BASF)	Selective post-emergence control of annual grasses in fine fescues.

cont. from page 21

perennial grasses, non-selective materials such as Roundup Pro or Finale can be used effectively. These products will remove both the unwanted weeds and any underlying turf. They should be made only during periods of the year when the weeds are actively growing and ample opportunity is available for renovating or re-establishing the turf.

General tips

Post-emergence herbicides should be applied when temperatures are moderate (less than 85 degrees F.) and growing conditions are good for both weed species and turf. Ide-

ally, early fall applications are most effective. This is a time when weed species are actively transporting materials to below-ground portions of the plant. Late spring or early summer applications can also be very effective. These applications should be made when soils are still moist and the weeds are actively growing. With the summer heat and possibly drought, the effectiveness of post-emergence herbicides will be much reduced.

Apply most post-emergence herbicides—and materials which contain phenoxy—judiciously around sensitive ornamental plantings. In spring and fall, many sensitive species can be injured through drift

or volatilization of these materials. Follow label instructions carefully to reduce the potential for non-target injury.

Cultural weed control programs can be developed to minimize the potential for weeds. Make herbicides one part of a comprehensive turf care program. Follow manufacturers' labeled instructions closely for best control and maximum safety with any selected material. □

—The author is an associate professor of turfgrass science at the University of Illinois Department of Natural Resources and Environmental Sciences.

Down South, control product tolerance a key

by TIM R. MURPHY, Ph.D. / University of Georgia

Proper turf nutrition and soil moisture will help you achieve maximum product efficacy.

Properly-maintained warm-season turfgrass is a good defense against strong weed competition.

Using correct fertility programs; following water requirements, mowing heights and schedules; and proper insect and disease control products all increase turfgrass vigor. They also improve the tolerance of warm-season turfgrasses to herbicides, and increase a weed control program's effectiveness.

Using herbicides without proper turfgrass management practices may control problem weeds, but will not produce high-quality turf.

Turfgrass managers in warm-season climates have a wide array of pre- and post-emergence herbicides that can be used to control weeds (Tables 1 and 2).

Tolerance factors

The single most important factor in se-

cont. on next page

cont. from previous page

lecting a herbicide is the tolerance of the turfgrass to the herbicide. Generally, most pre-emergence herbicides can be used on all established warm-season turfgrasses. There are exceptions. Ronstar is not labeled for use on centipedegrass or home lawns.

There is a dramatic difference in the tolerance of warm-season turfgrass species to post-emergence herbicides. Bermudagrass has good tolerance to MSMA and DSMA; however, carpetgrass, centipedegrass and St. Augustinegrass are severely injured by these herbicides.

Cultivars within a species may also respond differently to the same herbicide.

Post-emergence herbicides should be avoided when turfgrasses and weeds are stressed due to high air temperatures or drought. The tolerance of warm-season



Goosegrass germinates about two weeks later than crabgrass, at soil temperatures of 60-65 degrees F.



Southern crabgrass species germinate during February through April.

turfgrasses to post-emergence herbicides decreases at air temperatures greater than 90 degrees F., when turfgrasses are drought-stressed or when turfgrasses are growing under high soil moisture and high humidity.

Do not use post-emergence herbicides when turfgrasses and weeds are stressed by high air temperature or drought. Tolerance of warm-season turfgrasses to post-emergence herbicides decreases at air temperatures greater than 90 degrees F., when turf-

grasses are drought stressed, or when they are growing under high soil moisture and high relative humidity. Herbicides that contain 2,4-D, mecoprop, dichlorprop, imazaquin, MSMA and DSMA should not be applied at high air temperatures greater than 90 degrees F. because there is an increased risk of unacceptable turfgrass injury. Always follow the most restrictive warning that is shown on the label.

The tolerance of warm-season turfgrasses to post-emergence herbicides is generally lower during spring green-up than when the turfgrass is dormant or after full green-up. Research shows that the decline in turfgrass quality from the use of post-emergence herbicides during green-up is temporary and persists for two to six weeks after application. If a dense weed population dictates using a post-emergence herbicide during green-up, use only the lowest recommended rate or one-half the recommended rate to minimize herbicide injury to the turfgrass. If needed, the application can be repeated after green-up is complete.

Know the weed

Correct weed identification is a prerequisite for selecting an appropriate herbicide. After the weed has been identified, the herbicide label should be reviewed to determine if the herbicide will control the problem weed. Consult land grant university weed control guides for the effectiveness of herbicides in controlling weed species that are not listed on the herbicide label.

Application frequency

For some herbicides, particularly post-emergence products, a repeat application is necessary. For example, two applications of

TABLE 1. COMMON AND TRADE NAMES OF WARM-SEASON TURFGRASS PRE-EMERGENCE HERBICIDES¹

Common name	Trade name	Uses
atrazine	Aatrex, others	Annual broadleaf weeds.
benefin	Balan, others	Annual grass; some annual broadleaf weeds.
benefin+oryzalin	XL	Annual grass; some annual broadleaf weeds.
benefin+trifluralin	Team, others	Annual grass, some annual broadleaf weeds.
bensulide	Bensumec, Betasan, others	Primarily controls annual grasses.
bensulide+oxadiazon	Goosegrass/ Crabgrass Control	Annual grass control.
DCPA	Dacthal, others	Annual grass; some annual broadleaf weeds.
dithiopyr	Dimension	Annual grass; some annual broadleaf weeds.
ethofumesate	Prograss	Annual bluegrass control in bermudagrass and overseeded perennial ryegrass.
fenarimol	Rubigan	Annual bluegrass control in bermudagrass-overseeded cool-season turfgrasses.
isoxaben	Gallery	Annual broadleaf weeds. Does not control established perennials; provides residual control of some species that reproduce by seed.
metolachlor	Pennant	Controls yellow nutsedge and annual sedge; some annual grasses.
napropamide	Devrinol	Annual grass; some annual broadleaf weeds.
oryzalin	Surflan	Annual grass and some annual broadleaf weeds.
oxadiazon	Ronstar	Annual grass and some annual broadleaf weeds.
oxadiazon+benefin	Regalstar	Primarily controls annual grasses.
pendimethalin	Pre-M, Pendulum, others	Annual grass; some annual broadleaf weeds.
proflaminate	Barricade, Regalkade	Annual grass; some annual broadleaf weeds.
pronamide	Kerb	Winter annual weed control. May be used 90 days prior to overseeding bermudagrass with a cool-season turfgrass to control annual bluegrass.
simazine	Princep, others	Winter annual broadleaf weeds.

¹ Refer to the herbicide label for a complete listing of tolerant turfgrasses and labeled application sites.

SOURCE: DR. MURPHY

cont. on page 28



Use Image to control purple nutsedge, but not in bahiagrass or carpetgrass.

MSMA, at a 7- to 10-day interval, are needed to control crabgrass in bermudagrass.

In contrast, one application of Vantage will usually control crabgrass in centipede-grass. Pre-emergence herbicides may be applied either as full rate single applications, or as sequential repeated applications. With the sequential application program, one-half the maximum labeled rate is initially applied, with the remaining half applied 60 days later. With most pre-emergence herbicides, sequential applications tend to improve crabgrass and goosegrass control over the control achieved with a single application.

Application timing

Most pre-emergence herbicides control susceptible weeds during germination of weed seeds. Additionally, most pre-emergence herbicides require about one-half inch of rainfall or irrigation water to move the herbicide into the upper one to two inches of the soil profile.

A pre-emergence herbicide can undergo volatility losses and photodegradation the longer it remains on turfgrass foliage or thatch. Irrigation is advised unless a rainfall is anticipated within four to seven days of application.

Most species of crabgrass initiate germination when soil temperatures at the four-inch depth reach 53-58 degrees F. Depending on the geographical location, this will occur during February through April in the southern U.S.

Goosegrass germinates at a soil temperature of 60 to 65 degrees F., or approximately two to eight weeks later than crabgrass. On warm-season turfgrasses that are not fall-overseeded, pre-emergence herbi-

TABLE 2. NAMES OF WARM-SEASON TURF POST-EMERGENCE HERBICIDES¹

Common name	Trade name	Uses
asulam	Asulox	Grass weed control in St. Augustinegrass.
atrazine	Aatrex, others	Broadleaf, grass weed control.
bentazon	Basagran T/O	Primarily used for yellow nutsedge control.
bentazon+atrazine	Prompt	Yellow nutsedge and broadleaf weed control in centipede-grass, St. Augustinegrass, zoysiagrass.
bromoxynil	Buctril	Broadleaf weed control on non-residential turf.
2,4-D	Numerous formulations are available	Broadleaf weed control.
2,4-D+dicamba	Eight-One	Broadleaf weed control.
2,4-D+dichlorprop	Weedone DPC Amine, Weedone DPC Ester	Broadleaf weed control.
2,4-D+mecoprop +dicamba	Trimec Classic Trimec 992, Three-Way	Broadleaf weed control.
2,4-D+mecoprop +dichlorprop	Weedestroy Triamine, Weedestroy Tri-Ester	Broadleaf weed control.
dicamba	Vanquish	Broadleaf weed control.
diclofop-methyl ²	Illoxan	Goosegrass control in golf course bermudagrass.
diquat	Reward	Winter annual weed control in dormant bermuda.
DSMA	Numerous formulations are available	Grass weed control in bermudagrass and zoysiagrass.
ethofumesate	Prograss	Pre- and early post-emergence annual bluegrass control in overseeded bermudagrass; common bermudagrass suppression in St. Augustinegrass.
fenoxaprop	Acclaim	Annual grass control and suppression of bermudagrass in zoysiagrass.
fluazifop	Fusilade II	Bermudagrass control in zoysiagrass.
glyphosate	Roundup Pro	Winter annual weed control in dormant bermudagrass and bahiagrass.
halosulfuron	Manage	Controls yellow and purple nutsedge
imazaquin	Image	Purple nutsedge and wild garlic control in warm-season turfgrasses (except bahiagrass); controls certain annual broadleaf weeds.
mecoprop	Mecomec, Lescopex	Broadleaf weed control.
mecoprop+2,4-D +dicamba	Southern Trimec, Trimec Bent	Broadleaf weed control.
MCPA+mecoprop +dicamba	Trimec Encore, Encore DSC	Broadleaf weed control.
MCPA+mecoprop +dichlorprop	Weedestroy Triamine II, Weedestroy Tri-Ester II	Broadleaf weed control.
metribuzin	Sencor Turf	Goosegrass control in bermuda; prostrate spurge and numerous winter annual broadleaf weeds.
MSMA	Numerous formulations	Grass weed control in bermudagrass and zoysia. are available
MSMA+2,4-D+ mecoprop+dicamba	Trimec Plus	Grass and broadleaf weed control in bermudagrass and zoysiagrass.
pronamide	Kerb T/O	Annual bluegrass control in bermudagrass.
sethoxydim	Vantage	Annual grass control and suppression of bahiagrass in centipede-grass.
triclopyr+clopyralid	Confront	Broadleaf weed control in bermudagrass, centipede-grass and zoysiagrass.

¹ Refer to the herbicide label for a complete listing of tolerant turfgrasses and labeled application sites
² Diclofop-methyl has a state label for use in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee and Texas.

cides control annual bluegrass and certain annual broadleaf weeds. Annual bluegrass germinates at soil temperatures of around 70 degrees F. Apply the pre-emergence herbicide early in fall.

Post-emergence herbicides should be applied to small, actively-growing weeds. Perennial and annual weeds that grow

under good soil moisture conditions at moderate air temperatures are easier to control than weeds that are stressed due to adverse environmental conditions. **LM**

—The author is an agronomist in weed science at the University of Georgia Cooperative Extension Service.

SOURCE: DR. MURPHY