

Inside-out disease control

You'll get the best results when you use the right fungicide for the problem. New products give you even more options

BY MICHAEL L. AGNEW, PH.D.

Lawn disease problems are getting more complicated and selecting the proper fungicide to use is not a simple process, especially if it is to be used on residential lawns. The Food Quality Protection Act evaluation process has resulted in the voluntary removal of several fungicides from use on residential lawns. If you treat turf diseases, you may have to consider new solutions.

The ideal fungicide for use on residential lawns should be reasonably priced and possess curative properties, long term disease control and a positive human safety profile. The best way to select a fungicide is to understand key diseases and how a fungicide works on or in the plant.

How fungicides deliver control

"Mode of activity" is a phrase that refers to how the fungicide's active ingredient delivers disease control on or in a plant. You can use the mode of activity to give a general classification of the length of disease control provided by a particular fungicide, and to determine if the fungicide will provide any curative activity once the pathogen has infected the plant. There are three mode of activity types:

- ▶ contact,
- ▶ penetrant, and
- ▶ mesostemic fungicides.

Contact fungicides act only on the plant surface. They provide effective preventive

disease control on those fungi that are present on the outside of the plant, but little to no effect on a fungi present inside the plant. Because contact fungicides act on the surface, you must get thorough coverage with the fungicide to protect the plant. Contact fungicides are constantly being exposed to degradation through weathering and exposure to light (photodecomposition); thus, a typical contact fungicide will only provide protection for a period of seven to 14 days.

Examples of contact fungicides include chlorothalonil (Daconil), mancozeb (Fore) and quintozene (PCNB). Chlorothalonil produced after 1999 will no longer carry a label for residential lawn use.

Penetrating activity

Penetrant fungicides (sometimes referred to as systemic) act on the plant's surface, but they also stop the spread of fungi by penetrating the plant in quantities significant enough to be toxic to fungi inside. There are three subclassifications within penetrant fungicides: localized, acropetal and systemic.

Localized penetrants move into the plant tissue and remain at the point of entry. They protect only the immediate area where they were sprayed. There is little or no translocation within the plant, making it necessary to reapply every 14 to 21 days.

Examples of localized penetrant fungi-

continued on page 56



Necrotic ring spot symptoms



Necrotic ring spot being controlled


MODE OF ACTIVITY COMPARISON

Examples Characteristics:	Compass Mesostemic	Sterol Inhibitors Acropetal Penetrants	Chlorothalonil Protectant
acts on the plant surface			
absorbed by waxy layer of the plant			
redistribution by water			
redistribution by vapor movement			
penetrates plant tissue			
translaminar movement			
transported in the vascular system			

Compare costs & rates

As a lawn care operator or turf manager, you are not without various options. Newer fungicides may appear more expensive at first glance, but be aware that the application rates are a lot less than the rates for older standard fungicides. For example, the rate for Compass (0.1 to 0.25 oz/1,000ft²) is approximately 90% less than that of Daconil Ultrex (1.8 to 7.8 oz/1,000ft²). Reduced rate fungicides equate to less active ingredients placed into the environment, less exposure to an active ingredient by the applicator and less storage space.

MESOSTEMIC ACTIVITY



FUNGICIDE COST COMPARISON

Product	Rate	Spray interval (days)	Cost per application (\$/1000 ft ²)	Cost per treatment-day (\$/1000 ft ²)
Daconil Ultrex 82.5WDG	3.67 oz	14	2.06	0.21
Compass 50WDG	0.2 oz	21-28	3.38	0.12-0.16
Banner MAXX 1.24MEC	2.0 fl oz	28	3.82	0.13
Banner MAXX 1.24MEC + Compass 50WDG	1.0 oz + 0.15 oz	21-28	3.60	0.13-0.17
Heritage 50WG	0.4 oz	28	8.62	0.31
Cleary's 3336 50WP	6.0 oz	28	7.44	0.27

TABLE 1. FUNGICIDES LABELED FOR RESIDENTIAL LAWN

Pathogen	Fungicides	Rates (oz/1,000 ft ²)
Brown Patch	Azoxystrobin (Heritage 50WG)	0.2 — 0.4
	Flutolanil (ProStar 70WP)	1.5 — 3.0
	Thiophanate-methyl (Cleary's 3336 50WP)	2.0
	Trifloxystrobin (Compass 50WG)	0.1 — 0.25
Dollar Spot	Fenarimol (Rubigan 1AS)	0.75
	Myclobutanil (Eagle 40WP)	0.5 — 1.2
	Propiconazole (Banner MAXX 1.3 MEC)	1.0 - 2.0
	Thiophanate-methyl (Cleary's 3336 50WP)	2.0
	Triadimefon (Bayleton 50WP)	0.5
Gray Leaf Spot	Azoxystrobin (Heritage 50WG)	0.2 — 0.4
	Thiophanate-methyl (Cleary's 3336 50WP)	4.0 — 8.0
	Trifloxystrobin (Compass 50WG)	0.15 — 0.25
Leaf Spot (Melting out)	Azoxystrobin (Heritage 50WG)	0.2 — 0.4
	Thiophanate-methyl (Cleary's 3336 50WP)	4.0 — 8.0
	Trifloxystrobin (Compass 50WG)	0.1 — 0.2
Necrotic Ring Spot	Azoxystrobin (Heritage 50WG)	0.4
	Fenarimol (Rubigan 1AS)	4.0 — 8.0
	Myclobutanil (Eagle 40WP)	1.2
	Thiophanate-methyl (Cleary's 3336 50WP)	4.0 — 8.0
	Propiconazole (Banner MAXX 1.3 MEC)	4.0
Red Thread	Azoxystrobin (Heritage 50WG)	0.2 — 0.4
	Fenarimol (Rubigan 1AS)	8.0
	Myclobutanil (Eagle 40WP)	0.6 — 1.2
	Triadimefon (Bayleton 50WP)	0.5 — 1.0
	Thiophanate-methyl (Cleary's 3336 50WP)	2.0
	Propiconazole (Banner MAXX 1.3 MEC)	1.0 — 2.0
	Trifloxystrobin (Compass 50WG)	0.1 — 0.2
Summer Patch	Azoxystrobin (Heritage 50WG)	0.4
	Fenarimol (Rubigan 1AS)	4.0 — 8.0
	Myclobutanil (Eagle 40WP)	1.2
	Triadimefon (Bayleton 50WP)	2.0
	Thiophanate-methyl (Cleary's 3336 50WP)	4.0 — 8.0
	Propiconazole (Banner MAXX 1.3 MEC)	4.0
	Trifloxystrobin (Compass 50WG)	0.2 — 0.25

continued from page 54

cides include vinclozolin (Vorlan, Curalan) and iprodione (Chipco 26GT). Iprodione produced after 1999 will no longer carry a label for residential lawn use.

Acropetal penetrants enter the plant tissue and move translaminarily, from the top of the leaf surface through the leaf tissue and to the bottom of the leaf. They also move upward in the xylem, the water-conducting tissue of the plant's vascular system. This protects the plant tissue from the point of entry into the plant and upward. These fungicides have limited or no downward movement, so you must use enough water carrier to get the fungicide to the crown of the plant, typically 2 gal. of water per 1,000 ft². Most acropetal penetrants can provide up to 28 days of disease control, depending on rate and target fungi.

Some acropetal penetrants include propiconazole (Banner MAXX), triadimefon (Bayleton), azoxystrobin (Heritage) and mefenoxam (Subdue MAXX).

Systemic penetrants are similar to acropetal penetrants, but they also move through the phloem (food-conducting tissue) as well as the xylem. When applied, systemic penetrants can move both upward and downward in the plant tissue.

The only systemic penetrant used in turf is fosetyl Al (Aliette). This is limited to the control of Pythium blight and labeled to provide protection up to 21 days.

New activity mode

A new class of fungicide with a unique mode of activity called "mesostemic" was introduced in 1999. Trifloxystrobin (Compass) is the first turfgrass fungicide with a mesostemic mode of activity, combining some attributes of both contacts and penetrants to provide additional disease protection. This type of fungicide has similar activity to penetrant fungicides in that both

continued on page 58

What makes a mesostemic fungicide unique is that it forms a weather-resistant deposit on the surface in the waxy layer of the plant.

continued from page 56

penetrate the plant tissue and translocate translaminarily to the bottom of the leaf surface.

What makes a mesostemic fungicide unique is that it forms a weather-resistant deposit in the waxy surface layer of the plant. This deposit is a reservoir for continual penetration that replaces active ingredients lost to metabolism.

Mesostemic and contact fungicides both act on the plant surface to provide protection from external fungi. In addition, both types of fungicides can be redistributed on the leaf tissue.

Unlike a contact fungicide, however, mesostemic fungicides are rainfast and not

appreciably affected by weathering and photodecomposition.

A mesostemic fungicide also has the unique ability to redistribute at the plant surface by localized vapor movement despite lacking volatility and, consequently, the ability to move off-site. The vapor phase activity is seen over short distances of 2 in. to 3 in. within the plant canopy, and movement is greatest within the first 7 to 14 days after the application. This is when the free portion of the fungicide is available for redistribution. It also appears that the fungicide will move off of clippings to provide enhanced disease protection for the same time period.

The different ways a mesostemic fungi-

cide can provide plant protection adds up to 21 to 28 days of disease control.

Designing a program

Once you've decided which fungicide to use, use it properly. Consider these three factors:

- ▶ timing the fungicide application,
- ▶ water carrier volume, and
- ▶ turfgrass growth rate.

Timing. Most fungicide applications work best if applied on a preventive basis. You not only use less fungicide, but there is no plant damage. If you are treating for brown patch or gray leaf spot, get the first application on before weather conditions

continued on page 60

For 37 Years - the Foremost in Experience & Knowledge

LANDSCAPE LIGHTING

Acclaimed Distributor of the Industry Leaders:

<i>Nightscaping</i>		<i>Kim Lighting</i>
<i>B-K Lighting</i>		<i>Lumière Design</i>
<i>Kichler</i>		<i>Focus</i>
<i>Greenlee</i>		<i>Rockscapes</i>
<i>Hadco Lighting</i>		<i>Hanover Lantern</i>

Call TOLL FREE for Catalogs & Price Lists
1-800-457-0710

Go online to our website
View our catalogs, place an order at
www.callite.com

Unequaled Stock
CALIFORNIA SERIES transformers • Complete selection of lamps
All gauges of low voltage cable • Fixture accessories • Installation materials



California Landscape Lighting

A Division of Kina Enterprises, Inc.
31260 Cedar Valley Dr.
Westlake Village, CA 91362
Phone: 1-800-457-0710
FAX: 1-800-457-0730
Email: callite@pacbell.net

INFORM ♦ INTRODUCE ♦ INFLUENCE ♦ INSTRUCT

REPRINTS

Reprints of *Landscape Management* articles, advertisements, news items or special announcements are available through Advanstar Marketing Services. Customized to meet your specific needs, reprints are highly effective when you use them to:

- ◆ Develop direct-mail campaigns
- ◆ Provide product/service literature
- ◆ Create trade show distribution materials
- ◆ Present information at conferences and seminars

Extend your coverage to your website. Custom reprint packages include an E-Print of the same article to post on your website.

Landscape MANAGEMENT

ARTICLES
NEWS ITEMS
ADVERTISEMENTS

ADVANSTAR MARKETING SERVICES
MARCIE NAGY
1-800-225-4569 ext. 2744 • 440-891-2744
Fax: 440-891-2740
Email: mnagy@advanstar.com

Circle 135



Brown patch: individual patch on tall fescue

is rarely enough to provide 100% control of brown patch. Plan for two applications spaced 21 to 28 days apart.

A typical application for leaf spot or red thread control is to apply the fungicide at the first sign of disease. Since these are primarily foliage diseases, a one-time application may be enough to provide disease control. If weather conditions persist, a second application may be necessary 21 to 28 days later.

Water. Always use enough water to provide thorough coverage. Skimping on water volume can greatly reduce the length of disease control. If you don't use enough water when applying a fungicide, coverage is not good and the fungicide can be mowed off quickly. Control of foliar dis-

ease is best achieved when using water carrier volume of 2 gal. per 1,000 ft². If treating for soil-borne pathogens such as necrotic ring spot, water volumes of 4 gal. per 1,000 ft² are best.

Applying fungicides in too much water can also affect disease control — the fungicide may be moved beyond the turf and into the soil where it may not be available for uptake.

Growth rate. Finally, the efficacy of a fungicide is greatly affected by the growth rate of a turfgrass. Mowing removes fungicides, and if the turfgrass is growing too fast, more of the fungicide is removed with the increased clipping. **LM**

—Michael Agnew is research and development manager, Novartis Crop Protection, Kennett Square, PA.

continued from page 58

that are conducive to disease development occur — otherwise, the application should be considered as the post-infection kind. Both brown patch and gray leaf spot are active within the turf stand long before you see symptoms of damage. One application



BYPASS HAND PRUNER

The new Corona BP 6200 series bypass pruner takes pruning performance, comfort, and durability to new professional levels.

For starters, it won't slip from your hand — because the co-molded Corobond™ handles have been designed for a secure comfortable grip, greatly reducing hand and wrist fatigue. A larger patented bearingsurface, internal grease relief, and self-aligning pivot bolt with patented, hi-torque all-steel locking nut provides superior stability and a greatly reduced force to cut.

The BP 6200 features Coronium™ steel forged blade and hook for strength and durability.

The BP 6200 Bypass Pruner is the professional help you need. Take the Corona 2 week test drive and see for yourself.



The professional edge.

© 2000 Corona Clipper. 1540 East Sixth Street, Corona, California 92879 800-847-7863 or 909-737-6515 FAX 909-737-8657 www.coronaclipper.com

A Bellota Company



Series includes BP 6210, BP 6220, BP 6240