

Really know your pest enemies

Get to know — really know — the pests that commonly damage the turfgrass and ornamental plants under your care. You will dramatically increase your chances of controlling them.

It will take some training, but your customers, not to mention your company's bottom line, will benefit from your knowledge.

A limitless supply

The number of organisms that inhabit nearly all turfgrass and ornamental plants is almost limitless. These include disease pathogens, insects, weeds, micro-fauna and micro-flora, earthworms and vertebrate animals. Some cause no measurable damage and require no control. Some are beneficial because they help aerify the soil, decompose thatch or control other pest species.

Surprisingly, compared to the millions of insect species on earth, less than 100 plant-feeding species cause measurable damage to turfgrass and ornamental plants. In many instances (and to the untrained eye), pest and non-pest insects, including beneficials, look the same. That's why you must be able to accurately distinguish between them. In other words, before you

Knowledge is the key to dramatically increasing your chances of controlling turf/ornamental pests

BY DR. R. CHRIS WILLIAMSON



You have to get down and look carefully if you want to make an accurate pest ID.

can control a pest, you have to recognize what kind of pest it is and learn as much as you can about its biology. That knowledge must encompass behavior and habits, life cycle, plant-damaging life stages, and the life stage when it's most vulnerable to Integrated Pest Management (IPM) control.

suggests, it specifically targets the pest(s) causing the damage. Therefore, any information related to the pest in question is critical to successfully managing the pest.

Essentially, there are four steps for successful pest management:

Control with IPM

The concept or idea of understanding a pest's biology is fundamental to the concept of IPM, which is a synthesis of all possible control options available to you. IPM is a management approach and, contrary to what some people think, it is not exclusively biological or "organic."

IPM provides you with a variety of pest control options, strategies and techniques. As its title

continued on page 76

continued from page 74

► **Precise pest ID.** Spiders, mites, nematodes and other animals inhabit turfgrass and ornamentals. You must be able to tell non-pests and beneficial species from the pests. Often, you've already developed good control strategies for a particular pest because of previous experience. If not, you'll have to rely on resources such as university fact sheets, bulletins, insect flash cards, Web sites or insect collections. Other valuable resources include county extension agents, colleagues, consultants and agricultural chemical technical managers.

► **Behavior and habits.** After conclusively identifying the target pest, study its behavior and habits. You have to know where, when and how the target pest functions. In a sense, you have to be like a football coach who reviews game films of the opponent before building his team's game plan.

For example, consider the biology of black cutworm caterpillars. Since these caterpillars feed on the foliage of turfgrass plants at night, the most effective control strategy is to apply an insecticide that's been proven effective in black cutworm caterpillar control in the late afternoon or early evening, and to withhold irrigation for 12 hours. This gives the caterpillars time to contact and consume the insecticide. If you apply the product in the morning, you may decrease its effectiveness due to factors such as photodegradation (breakdown by sunlight) as well as volatilization (natural dispersion into the atmosphere). All pest insects have different control strategies.

► **Life cycle.** Knowledge of an insect's life cycle allows you to sample and monitor target pests. Monitoring is key to anticipating subsequent damage and implementing timely controls. Some insect species only have one generation per year, while others have multiple generations per year. As a result, control strategies for respective insect pests with only one generation per year are



often different than those for insect pests that have several generations per year.

Insects that have multiple generations per year typically require repeated control treatments. When the best IPM control strategy is the use of an insecticide, however, there's greater risk of promoting pest resistance, especially if similar insecticide chemistries (i.e. modes of action) are used. Technicians who understand the life cycles of insect pests are best prepared to develop control strategies that exploit the most susceptible or vulnerable life stages of the pests.

► **Damaging life stage.** You must be able to recognize signs that point to pest damage before it reaches a certain threshold. Again, frequent sampling and monitoring of plant material reveals early signs of insects and the damage they cause. This knowledge also allows you to solve the insect pest problem when it's easiest to solve — when the pest is at its most vulnerable stage.

Insects are more vulnerable to control measures or strategies at certain life stages. Typically, it's easier to control younger and smaller insect pests. For example, Japanese beetle grubs are considerably more vulnerable to certain insecticide treatments when they're young (newly hatched through first-instar grubs) compared to older or more mature (late-second and third-instar grubs).

This information enables you to time their treatments. Applying a curative insecticide treatment of a grub control product in the spring when the grubs are nearly fully mature would likely not be a good idea. This grub control treatment would be more effective if it was done in August

when Japanese beetle grubs are younger and more vulnerable. (Because dates vary due to geography and climate, check with your local extension service for best dates in your area.)

Be smart for a sound plan

Establishing an IPM program requires a sound understanding of growth habits and cultural requirements of your turf and ornamentals. You also need to understand any insect pests that threaten the turf and ornamentals you maintain. This includes knowledge of their behavior, life cycle, damaging life stages and vulnerable life stages.

Remember, there are three critical steps in successfully managing an insect pest. First, accurately identify the specific pest you're attempting to manage or control. Next, gain a comprehensive understanding of the organism's biology including but not limited to behavior and habits, life cycle, damaging life stage and vulnerable life stage. Finally, implement an IPM control strategy that provides the most effect control while taking into consideration the economic cost, potential environmental impact, and public perception. **LM**

— The author is the Turfgrass and Ornamental Specialist of the Department of Entomology, University of Wisconsin-Madison. He can be contacted at rcwillie@entomology.wisc.edu.

Beetle- mania



Adult Japanese beetles inflict characteristic feeding damage called skeletonization.

The Japanese beetle is a destructive urban landscape pest in the eastern United States. This metallic colored beetle was accidentally introduced to a nursery in

Riverton, NJ in 1916. It's likely that the beetle larvae (white grubs) arrived in the soil ball of nursery stock. The beetle is currently present in all states east of the Mississippi River except Florida.

Japanese beetles have one generation per year. The adults are present from June to August in Kentucky, when they'll actively feed and mate on susceptible plants, often defoliating them.

The females leave the plants to seek out turfgrass and lay eggs, which will hatch into grubs. The grubs feed on roots and organic matter in the soil and continue to grow throughout the summer. In the fall, as soil temperatures cool, they move down in the soil below the frostline for winter. In the spring, they move back up into the root zone and begin feeding. During May, they pupate in the soil in preparation for adult emergence in June. (Check with your local extension service for a more precise timing of the Japanese beetle life cycle in your region.)

Research and careful observation dispel some long-standing beliefs regarding Japanese beetles and their control

BY DAVID HELD

The damage Japanese beetles cause to trees, ornamentals and turfgrass is extensive. More than a half billion dollars a year is spent trying to control them. Since its introduction into the United States, the problem of managing this beetle has resulted in an array of management recommendations such as companion planting and host plant resistance. Let's take a careful look at some of the folklore surrounding Japanese beetles and try to balance those anecdotes with recommendations based on field research.

Trapping folklore

The Japanese beetle is strongly attracted to blends of plant odors, particularly those that are floral and fruity. Based on this, commercial traps were developed and proposed as a possible management tool for the adults and larvae. The use of traps in commercial and residential landscapes has been touted for reducing or eliminating de-

foliation by the adults and reducing the presence of grubs in the local turf. When evaluated in replicated experiments, however, the presence of traps didn't reduce the amount of damage to nearby plants. In fact, traps increased defoliation of nearby plants while having no effect on grub populations in surrounding turf.

Milky disease

Milky disease is a bacterial infection of grubs that results after it ingests spores of *paenibacillus popilliae* while it feeds. An infected grub has a distinctive milky-white appearance relative to a healthy grub. In the field, this disease is often present when grub populations are high. Milky disease is considered one of many natural buffers in the soil that suppress heavy populations of grubs.

Commercial products containing the bacterial spores have been developed for use as a microbial insecticide for Japanese beetles. Users apply powders containing lit-

erally millions of bacterial spores to infested turf for grub control. These products, however, didn't fulfill the claim as a "natural" insecticide. Also, the application of commercial powders to turf didn't increase the occurrence of the disease or result in reduced grub numbers. A contributing factor to the products' failure was their purity. Analysis showed significant contamination of milky disease powders with other non-infective bacterial spores.

Companion planting

Companion planting is a horticultural synergism between a crop plant and one or more plant species that results in the protection of the crop plant from pests. These garden companions are commonly aromatic herbs or other fragrant plants grown to mask the smell of a favorite garden ornamental or vegetable plant.

There are several recommendations regarding Japanese beetles and roses, one of their favorite foods, in organic gardening books and magazines. For example, interplanting members of the onion family, *allium* sp., anise or fennel *foeniculum vulgare*, and rue, *ruta graveolens*, with roses are said to protect the plants from Japanese beetle attacks. Likewise, certain plants like four o'clocks, mirabilis jalapa, and zonal gera-



A milky disease infected grub on the left, compared with a healthy grub on the right. Milky disease is common among concentrated infestations in the field but has been shown to be ineffective as a commercial microbial insecticide.



This infestation of grubs has completely consumed the roots of this turf. Turf damaged by white grubs like Japanese beetles is easy to diagnose. It will roll back like a loose carpet.

nium, *pelargonium x hortorum*, are considered effective trap crops because they may intercept beetles as they're flying to roses to feed.

When tested in replicated small garden plots, however, there was no reduction in the number of beetles on roses interplanted with rue, garlic chives or zonal geranium. Roses interplanted with geraniums generally had more beetles than roses planted alone.

Plant selection

Japanese beetle adults feed on over 300 species of plants in 79 plant families, but there are plants that are truly resistant to feeding. This resistance occurs across species and among cultivars of the same species. In field trials with different cultivars, resistance is quite evident when one cultivar is completely defoliated and the other cultivar next to it is untouched by the beetles. While no apparent resistance to beetle feeding has been shown among rose cultivars, there's considerable variation among the crabapples, lindens, elms and birch.

Selection of a resistant species or cultivar can be an important strategy for managing beetle feeding damage. Some of the information regarding resistant and susceptible woody plants can be found at: <http://www.uky.edu/Agriculture/PAT/rec/crop/pdf/entfa409.pdf>.

Japanese beetle adults feed on over 300 species of plants in 79 plant families.

The flowers of certain ornamental plants such as cannas, hibiscus, hollyhock, dahlias and roses are like candy to Japanese beetles. If these plants are in bloom, beetles will land on the flowers and begin feeding, almost ignoring the leaves. Among roses, cultivars with white or yellow flowers are more attractive to Japanese beetles.

continued on page 81

Beetles become paralyzed after feeding on the petals of zonal geraniums.



Tantalizing but toxic

Along with the anecdotes about trap crops, some plants are believed to be toxic to the Japanese beetle. It was even suggested that extensive planting of these alleged toxic species would slow or eliminate the spread of the beetles across the eastern United States. Leaves of the castor bean, *ricinus communis*, and leaves and flowers of the garden larkspur, *delphinium sp.*, and bottle-brush buckeye, *aesculus parviflora*, are said to kill the beetle if consumed. The toxicity of castor bean leaves to the beetles was disproved in a series of experi-

ments published in scientific papers during the 1930s and 1940s. Likewise, experiments with larkspur and bottle-brush buckeye failed to show any negative effects to Japanese beetles that feed on the flowers or leaves.

The zonal geranium, *pelargonium x hortorum*, is one plant that has fulfilled the claims of toxicity to the beetle. Zonal geraniums are a preferred food plant for Japanese beetles, which proves to be a fatal attraction. In 1920, a USDA entomologist noticed large numbers of dead beetles under zonal geraniums growing in the full sun. Beetles can eat just half a petal and become temporarily paralyzed. The paralysis occurs only when beetles

feed on the flowers and not the leaves, although paralyzed beetles falling from the flowers are often seen on the foliage.

Paralysis from feeding also occurs with other ornamental geraniums such as ivy geraniums, *pelargonium peltatum*. Beetles that become paralyzed on the plants in the landscape lose about 30% of their body weight in just a few hours. This likely explains why beetles die on plants growing in the full sun but may recover on plants growing in the shade. If beetles do recover, they typically shake off their buzz and feed again on the flowers that just intoxicated them.

—DH

The "ORIGINAL"

Back Saver™

Tailgate Assist System

US Patent 6,126,223



Thanks to this revolutionary and patented design, your heavy tailgate can be raised and lowered with virtually no effort, keeping your back and your bank account healthy.

The BackSaver Tailgate Assist . . .

- Helps reduce injuries, worker's comp. claims & property damages.
- Takes 100% of the weight off most tailgates. (2-sided assist)
- Will not allow most tailgates to fall down. (2-sided assist)
- Is easy to install & will work on most utility trailers and tailgates.

Dealer Inquiries are Welcome!

PRICES START AT \$139⁹⁵

877.388.8895 – www.thebacksaver.com



A littleleaf linden tree in mid-summer completely defoliated by adult Japanese beetles. Notice the typical top-down feeding pattern common to trees defoliated by Japanese beetles.

continued from page 79

Varieties with larger blooms are also more appealing to the beetle. In experiments with red or yellow flowering rose bushes, Japanese beetles will land on the yellow flowering ones, sometimes ignoring even the most fragrant red varieties.

Biorational insecticides

Adult beetles are commonly managed with applications of short-residual insecticides, like carbaryl or synthetic pyrethroids. There has been increased interest in the use of other biorational insecticides, including various formulations of neem-based feeding deterrents. In laboratory tests, these products effectively reduce the amount of feeding. In field tests when beetle infestations are heavy, even repeated applications aren't sufficient to prevent damage.

Regardless of the control product, once a plant becomes damaged it can serve as a beacon to other beetles, attracting them with odors produced by the damaged

leaves. Therefore, preventing feeding damage early and keeping plants undamaged as long as possible reduces the attraction of more beetles to those plants. **LM**

— *The author is with the University of Kentucky Entomology Department, Lexington, KY. Contact him at dwheld@uky.edu.*

NEED A DECK THE SIZE OF AN AIRCRAFT CARRIER?

A range of heavy-duty deck sizes – up to 72" – is just one of the many options you can choose from on your Yazoo/Kees mower. Featuring an engine with up to 34 HP, zero turning radius, built-in comfort, and much more, every Yazoo/Kees is built to last. Need a mower to help your productivity take off? No problem! Yazoo/Kees.

YAZOO/KEES

Contact your local YAZOO/KEES dealer
or call 1-877-368-TURF
www.yazookees.com

Large 'scale' woes

When a customer called Jeff Hazlett and complained that her family couldn't use the front door due to swarms of bees and flies, he had his suspicions about what was attracting the insects. He inspected the customer's landscape plantings and confirmed that scales had infested three 9-ft. tall star magnolias in her front yard.

"Honeydew produced by the scale insects was dripping on twigs and stems near the front door, and insects were feeding on it," says Hazlett, owner of Ever Green Tree & Lawn Care in Lancaster, PA. "Sooty mold had formed over the honeydew, and whole branches had turned black. It wasn't a pretty sight."


Sure signs of scale

The presence of honeydew is a telltale sign of scale infestation. Honeydew is a sugary, liquid waste product that covers surrounding areas. This sticky, sweet liquid serves as a growth medium for sooty mold, a black fungus that eventually covers honeydew-laden areas. Another indication of scale is a greater-than-usual number of nuisance insects such as ants, bees, flies, hornets and yellow jackets, which are all attracted to honeydew.

In severe cases of scale infestation, foliage and fruit turn black from sooty mold and drop prematurely. In large enough numbers, scales remove enough sap from plants to starve limbs, branches and leaves.

Recognize a scale infestation and the type of scale before you begin treatment

BY DEBBIE CLAYTON



This soil application initiated scale control.

For his customer with scale-infested magnolia trees, Hazlett soil applied Merit Insecticide in early spring, and followed it with a treatment of dormant oil. Later in spring, he made a foliar application of Merit mixed with a contact insecticide, ensuring that coverage was complete. The problem vanished by mid July.

"The customer was really impressed that the combination controlled scale so quickly," Hazlett says. "I attribute the successful treatment to the systemic properties of Merit combined with the dormant oil spray."

Preaching IPM

With three full-time employees, Hazlett provides landscape maintenance for primarily residential customers in Lancaster

County. Though he's been in business for 15 years, he keeps his operation small, offering customized services such as fertilizing, spraying and mulching. He regularly monitors plantings, explaining to his customers the benefits of an integrated pest management (IPM) system.

In spring 2002, after two relatively mild winters in eastern Pennsylvania, an outbreak of cottony camellia scale on yews and hollies erupted in Hazlett's area. "Cottony camellia is a soft scale," he says. "If customers have a lot of evergreens and hollies, I'll make a preventive treatment of Merit for scale."

In addition to applying insecticides and dormant oils, Hazlett often prunes out damaged branches and foliage, and fertil-

izes beds or trees to restore vitality. Because pachysandra is easily affected by scale, he often applies dormant oil in spring to prevent infestations. **LM**

— The author is a freelance writer and frequently writes about the Green Industry from her home in Horsham, PA.



Only a mother could love these magnolia scales with crawlers emerging.

Two scales, two treatments

Not all scale species ingest sap and produce honeydew. Two general types of scales commonly appear on U.S. ornamental trees and shrubs: soft scale and armored scale. Learn to identify the two since control strategies are different for armored (hard) and soft scale.

Soft scale characteristics:

- ▶ Bodies are exposed and immature stages are "soft"; mature adult females have a hardened shell to protect the young
- ▶ Females are oval-shaped and usually conspicuous — about 1/10-in. long
- ▶ One generation per year
- ▶ Produce honeydew, attracting nuisance insects
- ▶ Overwinter as immature females
- ▶ Relatively prolific, producing more than 600 eggs at a time

Soft scales damage plants and make

them unsightly by sapping juices, making sooty mold grow on honeydew. This substance is difficult to remove from parked cars, houses, porches and sidewalks.

Armored scale characteristics:

- ▶ Produce detached wax shell or "armor" over their bodies
- ▶ Females are usually elongated or circular and about 1/10-in. long
- ▶ Don't produce honeydew
- ▶ Two or more generations per year
- ▶ Overwinter as eggs, immatures or adults
- ▶ Produce fewer than 50 eggs at a time

Armored scales also damage plants by inserting long, straw-like mouthparts into plant tissue. Damage may appear as chlorotic areas of the leaves or yellowing of entire leaves or needles, resulting in leaf drop or dieback.




Convertible Mowers

Walk the Hill with the Convertible mower. For steep hills, roadside ditches and low limb trees. The Convertible can be utilized as a hydro walk behind. Then as easy as flipping a lever, the Convertible transforms into a rugged dependable "L" so you can Ride the Distance!

New Foot Assist Deck Lift

Offered with the following attachments:



Distributors:

New England States	Beta Power Products	508-393-1924
PA-NJ-DE	Canns Bilco	800-811-0285
VA-WV-MD-NC	UTECO	888-368-8326
TX-OK-	Kretchmar	580-395-2606
NY	Stiefvater Distributor	315-853-5581
KS-MO- S.IL	AES	800-843-3546
GA-SC-AL-FL	Southeast Equipment	800-241-4384
TN	Southern Lawn	877-204-6873
IA-NE	Scott-Hourigan Co.	402-362-7711

Distributor inquiries welcome for open areas.

Rich Mfg., Inc. • P.O. Box 145 • Thorntown, Indiana 46071
www.convertiblemower.com • 765-436-2744