

Table of Control Measures for Major Shade Tree Pests

Pest	Material & Formulation	Amount per 100 gals. water	Remarks
Aphids	Thiodan 50% W.P.	1 lb.	Long residual action. Observations indicate that many foliar insects may be well controlled.
	or		
	Sevin 50% W.P.	2 lbs.	Effective against a broad range of insects.
	Lindane 25% W.P.	1 lb.	Good residual action.
Cankerworms	or		
	Malathion 50% W.P.	4 lbs.	Double strength in the fall for gall-producing-type aphids is especially effective.
	DDT 50% W.P.	2 lbs.	Usually, one application as the leaves are unfolding is sufficient for effective control.
	DDT 25% E.C.	1 qt.	
Lead Arsenate	3-4 lbs.		
Methoxychlor 50% W.P.	2 lbs.		
Mites	Tedior.® 25% W.P.	1 lb.	Kills eggs and young mites. Slow kill of adults. Extremely long residual action — 6 to 10 weeks.
	or		
	Aramite 15% W.P.	2 lbs.	Summer use for adults and young mites.
	Ovex 50% W.P.	Deciduous-1 lb. Evergreen-2 lbs.	Kills eggs primarily.
Gypsy Moth	or		
	Malathion 25% W.P.	4 lbs.	Use on protective schedule.
	Ethion .67 Superior 60 Oil E.C.	2-3 qts.	Apply when mites first appear.
	DDT 50% W.P.	2 lbs.	
Borer Caterpillar Type	or		
	Lead Arsenate	3-5 lbs.	Apply as close to egg hatching as possible.
	Sevin 50% W.P.	2 lbs.	
Grub Type	DDT 25% E.C.	1 gal.	Dates of application vary considerably. The insect species and the host to be treated are different. For detailed recommendations consult your pesticide supplier or technical advisor.
	or		
	DDT 50% W.P.	4 lbs.	
Leaf Miners	Lindane 20% E.C.	2 qts.	Materials used against adults may require second application 10 days later. Observation of mining within the leaves is required for efficient timing of larval sprays.
	or		
	DDT 50% W.P.	4 lbs.	
	Lindane 25% W.P.	1 lb.	
Scale Insects	or		
	Malathion 25% W.P.	4 lbs.	For Dormant Sprays. (Thorough coverage is necessary)
	Sevin 50% W.P.	2 lbs.	
	or		
	DDT 50% W.P.	2 lbs.	For Foliar Sprays — directed against the crawler stages.
	Superior 70 Second Oil	2-3 gals.	
Ethion Superior Oil	2-3 gals.		
Elm Leaf Beetle	or		
	Malathion 25% W.P. Plus DDT 50% W.P.	4 lbs. 2 lbs.	For Foliar Sprays — directed against the crawler stages.
	or		
	Sevin 50% W.P.	2 lbs.	Apply when eggs hatch or when summer crawlers first appear.
or			
Ethion .67 Superior 60 Oil E.C.	2-3 qts.		
European Elm Bark Beetle	DDT 25% E.C.	1 qt.	Where resistance to DDT is suspected, lead arsenate or Sevin are logical choices. Sevin 85W may be substituted where mist blowers are utilized.
	or		
	DDT 50% W.P.	2 lbs.	
	Lead Arsenate	5-6 lbs.	
Anthracnose	or		
	Sevin 50% W.P.	2 lbs.	Dormant** Hydraulic sprayers — 8-12 gallons of finished spray should be applied per tree.
	DDT 25% E.C.	8 gals.	
	or		
Methoxychlor 25% E.C.	8 gals.	Dormant For concentrate sprayers, use 46 gallons of either E.C. to 54 gallons of water. Spray at least 2-5 gallons of finished spray per medium to large tree.	
DDT 25% E.C.	12% emulsion		
Powdery Mildew	or		
	Methoxychlor 25% E.C.	12% emulsion	
Leaf Spot	** If summer applications are needed — DDT (1%) for hydraulics and DDT (6%) for mist blowers may be used.		
	Puratized Agricultural Spray	1 pt.	Spray in early spring before buds swell, and again before the leaves unfold. Follow dormant application with one or two applications at weekly intervals depending upon the rainfall.
European Elm Bark Beetle	Zineb	2 lbs.	Apply midsummer on lilac. Avoid spraying at high temperatures.
	or		
Anthracnose	Puratized Agricultural Spray	1 pt.	Helpful for leaf spot on walnut, elm, and maple.
	or		
Powdery Mildew	COCS	4 lbs.	Preventive treatments are suggested where severe in past seasons.
	or		
Leaf Spot	Ferbam	1-2 lbs.	

and lindane has been recommended for some of the grub types.

Leaf Miners

There are many species of leaf miners — elm, birch, and oak leaf miners, to name but a few. Injury resulting from these pests frequently appears as blotches or blisters on the foliage. Partially or completely mined leaves turn brown and are very noticeable.

Loss of leaves is one end result of attack by the birch leaf miner. This damage is usually confined to newly developing foliage. Beyond the second generation, birch miners generally invoke little damage except for such feeding as is necessary to maintain their population. Repeated leaf losses can cause death or weaken the tree, making it susceptible to attack by other insects or diseases. Hence, treatment to curb infestations before they get out of hand is vital.

In June, the adults deposit their eggs either on the foliage or in the leaf tissue. Once eggs hatch the tiny larvae begin feeding between the upper and lower surfaces of the leaves. Birch leaf

miners may have as many as four generations per year; oak leaf miner up to three. Adults of many leaf miners are tiny, two-winged, black midges or flies. The mature birch leaf miner, however, is a small, black, four-winged wasp-like sawfly.

Leaf miners are relatively difficult to curb. Timing is critical. lindane, applied when leaves are expanding and small blisters or mines are just becoming evident, has given a good degree of control.

Larval stages of leaf miners may be curbed by applications of lindane in the latter part of May for the birch leaf miner, and in early June for elm leaf miners. Treatment with malathion is also effective for control of birch leaf miners.

Scale Insects

Because of their small size and general appearance, scale insects are often overlooked until injury has reached advanced stages. This is often the point where it becomes too late to save part or all of the trees attacked.

Scale insects have a tremendous

potential for reproduction. Preventive control sprays are advisable in areas where these pests are known to be active from year to year. Treatment with appropriate chemicals should be made on a "planned" schedule, not merely once damage is evident.

Injury caused by scale insects can be extensive, manifesting itself in any of four ways: (1) reduction of growth; (2) abnormal growth; (3) death of the host plant; (4) secretion of an unsightly sticky honeydew substance.

Scale insects are members of a large and complex family. For purposes of discussion, they can be divided into three groups. The first is the covered or armored scale insects, which include such pests as oystershell, euonymus, San Jose, and others. At varying intervals, the young of this species crawl out from under the protection of the female's shell or scale. After a short period of activity, they find a favorable location on the host to insert their threadlike mouthparts and feed by sucking the sap. Soon they lose their skins, and, if female, their legs and antennae also. The females combine cast skins with exudations of wax and form them into coverings. After the fourth molt (growth stage), the males soon become minute two-winged insects.

The second type is the naked or soft scale insects. Included here are such common pests as the cottony maple scale and lecanium species. They are usually humped and resemble the shell of a turtle, are not protected by a separate waxy scale or shell, and perhaps most important, frequently move from place to place while feeding and may crawl back to the twigs in the fall of the year.

A third type of insect closely related to the scale group is the mealybug. Types are too numerous to list at length.

For best results in controlling scale insects it is important that correct identification of the species be made. As with many other pests, timing of spray applications for scale insects is critical. This is especially true with the covered scales where the young come out from under the protective female shells and crawl around. The crawler stage is the best time to

MALATHION



INSECTICIDE

Outdoors, Malathion protects shrubbery, flowers, fruits and vegetables from leafhoppers, beetles, many species of worms, and other destructive pests. In the greenhouse, Malathion controls aphids, mites and soft scale. A friend in the home too, this all-around insecticide kills flies, mosquitoes, silverfish, bedbugs, carpet beetles, crickets,

fleas, ticks, and resistant roaches.

Malathion has high residue tolerance. In fact, it is so safe that it can be applied directly to grains, seeds and household pets. Premium Grade Malathion is almost odorless. For successful growing and repeat sales, formulate with Malathion and sell Malathion formulations in your retail store.

PROLIN* RODENTICIDE -- ANOTHER MONEY-MAKER

Prolin bait, available under various brand names, is as safe as warfarin and more thorough. It contains warfarin plus a helper ingredient for devastating two-way action against mice and rats. Write Prentiss for dealer information.



Prentiss Drug & Chemical Co., Inc.

101 WEST 31st STREET, NEW YORK 1, N. Y. • 9 SOUTH CLINTON STREET, CHICAGO 6, ILL.

Atlanta • Detroit • San Francisco • Los Angeles • Toronto • Montreal • Fort Worth • Pittsburgh • Philadelphia

* TM OF WISCONSIN ALUMNI RESEARCH FOUNDATION

control these scales. Time of spray applications can vary from April through September, depending upon the species and number of generations per year. Recommended materials include Ethion Superior Oil or Superior 70 Second Oil for dormant sprays. Sevin, a malathion-DDT combination, or the Ethion.67 Superior 60 Oil E. C. are excellent choices for foliar sprays.

Elm Leaf, Elm Bark Beetles

In recent years many cases of the dreaded Dutch Elm Disease have been traced to the feeding habits of the European elm bark beetle, aided by the widespread and highly destructive elm leaf beetle. It has been found that severe attacks of elm leaf beetles upon the foliage weaken elm trees and make them attractive breeding grounds for elm bark beetles, which we now know are carriers of Dutch Elm Disease.

Large infestations of elm leaf beetles will cause leaves to yellow and fall prematurely. The adult insects are about $\frac{1}{4}$ inch long. Those found in spring and summer appear quite yellowish with dark

stripes along the sides, while in the fall and winter, they are a more uniform olive green.

European elm bark beetles are reddish brown in color with pitchy, red wings, and approximately $\frac{1}{8}$ inch long. The rear portion of their bodies appears to have been chopped off abruptly. These pests feed on small twigs, particularly in the crotches of healthy tree branches. The breeding galleries made by the adult female beetles beneath the bark of weakened or dead trees range from 1 to 2 inches long and contain 80 to 140 eggs per adult gallery.

For effective control of the bark beetle a thorough dormant application is required. As a supplementary measure, a spray in mid-July is also advisable.

In curbing leaf beetles, sprays should be applied when leaves are $\frac{3}{4}$ grown or as soon as feeding is noted (usually late May or early June). A second dosage in mid-summer may be needed, particularly if only some of the trees in the area have been sprayed previously.

DDT is suggested as a dormant spray for elm leaf and elm bark beetles. For late spring and summer sprays, DDT, lead arsenate, or Sevin can be used to halt both pests. New work by several research groups indicates that Thiodan[®] insecticide will probably be labelled for the control of both pests.

Significant Disease Pests

In addition to the many insect pests which threaten shade trees each spring and summer, there are fungus problems to beware as well. A few of the more significant disease pests are cited here to aid in recognition and control.

Anthracnose

Anthracnose is a fungus which overwinters in leaves and cankers and attacks the twigs or leaves of ash, linden, horse chestnut, maple, oak, sycamore, and certain other hardwoods. Oak and sycamore are both attacked by one particular species of this pest, while each of the other varieties is plagued by a different species. Thus, leaf injury to various types of trees differs

How to spot & control danger in turfgrasses - forests - ponds



Couch: DISEASES OF TURFGRASSES

by Houston B. Couch, Department of Plant Pathology, Pennsylvania State University.
1962. 304 pages. \$11.00
Consulting Editor: J. Ritchie Cowan

"Nearly 100 pages are given over to analysis of turfgrass disease chemicals, and grasses susceptible to ailments. Arranged both by common and technical names, these tables provide a handy guide which could be used by technical directors and servicemen alike. Couch's new text is more than a handbook, however. In the first 178 pages, the distinguished author delves insistently into the rudiments of both fungus and nematode-incited diseases, and gives valuable pointers on how to tell one disease from another. Another highlight of the volume are the illustrations, both full color and black-and white, which give vivid reinforcement to the details set forth in the text." — *Pest Control*



Graham: CONCEPTS OF FOREST ENTOMOLOGY

by Kenneth Graham, Professor of Forest Entomology, The University of British Columbia, Vancouver.
1963. 392 pages. \$9.50
Consulting Editor: Peter Gray, University of Pittsburgh.

"Practical uses of theoretical information about the detrimental effects of insects in forests is offered in the latest of Reinhold's biological books... Insects are not treated superficially; rather, in-depth discussion on the pest and the manner in which it damages trees, whether by boring, leaf mining, or root chewing, gives the student of forestry a sound basis upon which to determine timber loss. A table is included which sets forth the principal insects found in forests and what part of trees they damage. This makes damage detective work simpler... a refreshing, practical slant that pinpoints the study of entomology in the scheme of forest management." — *Pest Control*



Bennett: MANAGEMENT OF ARTIFICIAL LAKES AND PONDS

by George W. Bennett, Aquatic Biologist and Head of the Aquatic Biology Section, Illinois Natural History Survey, Urbana.
1962. 320 pages. \$8.00
Consulting Editor: Peter Gray, University of Pittsburgh.

"The author has included the most recent scientific advances and practical techniques in artificial lake and pond management. For those interested in a comprehensive and integrated presentation of the dynamics and management of warm-water fish populations in artificial lakes and ponds, this is just the book." — *Commercial Fisheries Review*

ORDER ON APPROVAL! Inspect the book of your choice for 30 days. See how it can serve you both as a stimulating refresher and handy reference. Then if you are not completely satisfied return it and pay nothing. Clip out and mail the convenient coupon to us today, we'll rush you your book.

REINHOLD BOOK DIVISION, 450 PARK AVENUE, NEW YORK 22, N. Y.

REINHOLD BOOK DIVISION, Dept. M-158
430 Park Avenue — New York 22, New York

Please send me a copy of the book(s) checked below for 30 days' approval (U.S.A. & Canada only) under the following terms:

Purchase price enclosed (Reinhold pays regular delivery charges)

Bill me (plus delivery charges)

Bennett: Management of Artificial Lakes and Ponds (\$8.00)

Couch: Diseases of Turfgrasses (\$11.00)

Graham: Concepts of Forest Entomology (\$9.50)

Name _____

Professional Affiliation _____

Address _____

City _____ Zone _____ State _____

Save money! Enclose full purchase price with your order and Reinhold pays regular delivery charges. Same return privilege guaranteed. Please include sales tax on Calif., Ohio, Penna., and NYC orders. Send check or money order only, do not enclose cash.

slightly in coloring, size, and extent.

Two principal types of injury are usually distinguishable. One is early leaf blight which results in the complete killing of young leaves during April or May. This type of injury varies from partial to complete defoliation of tender leaves and is similar to frost injury in appearance. It is most common on sycamore. The second type of injury shows up later in the season as irregular brown areas adjacent to and surrounding diseased spots on midribs and lateral veins of mature leaves. This type rarely causes serious defoliation, but the fungus may spread through the twigs into younger branches, producing cankers which in turn can cause twig and branch dieback.

As with many other tree pests, successful control by spraying depends on the frequency and the thoroughness of application. A suggested schedule follows:

- (1) Just before the buds start to swell in the early spring.
- (2) When the buds are breaking and before the leaves unfold.
- (3) One or two applications at weekly intervals depending upon the amount of rainfall.

The following treatment is suggested in curbing anthracnose: Puratized Agricultural Spray (a mercury compound spray) — 1 pint per 100 gallons of water applied in the dormant stage followed by 2 treatments of Zineb (2 pounds per 100 gallons of water).

Powdery Mildew

Extensively distributed and occurring on a wide variety of plants, the powdery mildews have distinguishing characteristics which enable specialists to identify them. The powdery mildews are important in this country on maple, oak, alder, elm, poplar, and willow.

The damage caused by this disease is usually slight except when young trees are attacked. The powdery mildews are usually confined to the upper or lower surfaces of leaves, forming a white superficial growth visible to the naked eye. Occasionally, when conditions are right, and the infestation is severe, the mildews will progress onto fruits and twigs of the host. In addition to covering the surface



Citywide programs for controlling shade tree pests are frequently handled by CAs. Spraying shown above is part of a Japanese beetle control program in Sacramento, Calif.

of the leaf, the mildews also produce a sucking organ which penetrates the cells of the host to withdraw food. The spores of the developing mildews are readily carried by the wind and spread the fungus widely during summer months. The fungus overwinters in the black fruiting bodies found on fallen leaves, ruptures in the spring, discharging the spores which are carried by air currents to infest new foliage.

The powdery mildews definitely create an unsightly condition on the leaves of the host trees, but they may be checked during the growing season by use of a sulfur dust or spray at weekly intervals. Burning the fallen leaves in autumn is also quite helpful in a small or confined area.

Leaf Spots

Leaf spots on trees can be caused by insects, toxic gases, bacteria, and fungi. Leaf diseases such as this can be important when defoliation results or when the infestation is so severe that most all the leaves are involved and cannot function normally. Defoliation for several successive years may bring about the death of hardwoods. Species of trees commonly infested by the leaf spot fungi include: elm, chestnut, oak, red maple, hickory, ash, locust, and sycamore.

Characteristic of leaf spot is the formation of dead areas in the leaf. Size and shape of the area can vary from small to large and round to irregular. The deadened tissue varies in color from yellowish to all shades of brown and

black. Some of the dead areas frequently fall out, leaving holes (sometimes called "shot hole").

The spread of the leaf spot fungi is basically the same with all species. They overwinter in the old fallen leaves, develop and spread to the newly developing leaves in the spring. Wet seasons favor increased infestations of leaf spot.

Recommendations for control include the burning of fallen leaves in the fall. Dead and cankered twigs should be pruned out before the buds break in the spring. If infestations are severe, spraying with ferban or COCS (copper oxychloride-sulfate) is helpful. Puratized Agricultural Spray (mercury compound) is also useful in the control of leaf spot of walnut, elm, and maple.

Subsoil Problems Blight Lawns

Dead spots on lawns or patches that yellow during dry weather may be an indication of subsoil problems, according to Vaughn H. Holyoke, assistant crops specialist with the Cooperative Extension Service, University of Maine, Orono.

Pockets in subsoil, 4" to 8" below the top, can collect water in fall and winter, causing winter-killing, Holyoke explains.







"These areas can be the starting point of a severe weed problem, as well," Holyoke adds. In established lawns, turf renovation is the only solution to subsoil problems, so be sure to watch out for it when establishing new lawns, Holyoke warns.



Stauffer's new liquid
SEVIN*
4 FLOWABLE

I N S E C T I C I D E

CONTROLS MANY SHADE TREE INSECTS

-  May be used in mist blowers, foggers and all other types of sprayers
-  Leaves a minimum of visible residue
-  Is easy to use safely, less toxic than DDT to fish and wildlife
-  Controls insects resistant to chlorinated hydrocarbons
-  Controls a wide variety of insects on deciduous and evergreen trees
-  Is an extremely fine dispersion of SEVIN in water

For further information and directions for use, write Stauffer Chemical Company, Agricultural Chemicals Division, 380 Madison Avenue, New York 17, New York

*Reg. T. M.



When Writing to Advertisers Please Mention WEEDS AND TURF

W-15



Sevin

Insecticide:

New Carbamate for Turf, Trees, and Ornamentals

By DR. J. B. HARRY

Marketing Manager, Agricultural Products
Union Carbide Chemicals Company
New York, New York

CONTRACT spraying of home lawns and gardens is growing by leaps and bounds.

The day is past when the average homeowner can keep most of his pests under control with two or three stock pesticides. Greater insect depredations, a buildup of insect resistance to some spray materials, the wide variety of sprays and dusts, and the recent concern about spray materials have all contributed to take pest control out of the hands of the homeowner and turn it over to the specialist — the contract applicator.

One of many insecticides which operators are now using for a variety of jobs is Sevin, a product of Union Carbide. Being a carbamate, it is unrelated to many older compounds. Chemically known as carbaryl (1-naphthyl N-methylcarbamate), Sevin has a broad spectrum of usefulness and a low degree of toxicity. Spraymen who must handle it are relatively safe from any injurious effects, researchers have reported.

Normal pesticidal precautions should be observed, however. When used according to directions, though, Sevin is not likely to bother customers' children or pets if they come into contact with it on foliage. If it drifts, it will not contaminate crops or harm domestic animals, small game, or fish, when used at the prescribed rates. It is said to be 200 times

safer to goldfish, than is DDT.

Formulations Available

Sevin insecticide is available as a 50% wettable powder, an 85% microfine wettable powder, several liquid formulations including a suspension of 4 pounds per gallon of active material (called Sevin 4 Flowable), 5 and 10% granular formulations, and dusts, each produced by various formulators.

There are some formulating details of which CAs should be informed. To maintain quality of spray preparations, formulations should be mixed accurately and with care. To a partly filled tank with agitator operating, Sevin can be added along with any other emulsifiable formulations desired. While spraying, the agitator should continue operating so as not to let the preparation settle. After spraying, flushing with clean water will eliminate residues.

Separable suspensions of Sevin insecticide should not sit in the tank overnight.

Sevin is compatible with commonly used insecticides such as DDT and other chlorinated hydrocarbons. Compatible organic phosphate compounds include Guthion, malathion, and parathion. Botanicals are also compatible with the product. Fungicides which may be applied along with Sevin include glyodin, organomercury compounds, fixed cop-

pers, dithiocarbamates, sulfur, captan, Cyprex, and Phaltan.

Sevin should not be mixed with any alkaline materials. Bordeaux, lime, lime sulfur, and casein-lime will lessen effectiveness of this new compound.

All common miticides are compatible with the chemical, which is an advantage because Sevin alone does not control certain kinds of common mites.

Sevin controls many insects that have built up resistance to chlorinated hydrocarbon and phosphate insecticides. According to Dr. Edward J. Duda, director of the Bartlett Tree Research Laboratories, Sevin has a good residual effect, and has given up to two and three weeks control of insects such as lace bugs on oaks and sycamores.

Bartlett experts have also had good results using Sevin on birch, oak, and boxwood leaf miners, Japanese beetles (which seem to be moving from gardens into rougher areas), chinch bugs, leaf tiers, tent caterpillars, cankerworms, and periodical cicadas (17-year locusts).

Cicadas were especially troublesome in sections along the East Coast in 1962, and are scheduled to appear over a broad band in the east-central United States in 1963.

Cicada Control

Cicadas are controlled by preparing a 2% suspension of wettable powder; approximately a gallon of this is applied to each 50-foot tree. Application should be delayed about 10 days after emergence from the ground; this is the time when female cicadas begin laying eggs in twigs of trees.

Other pests on which Sevin has been successful include bagworms, elm leaf beetles, rose aphids and slugs, lecanium scales, and pine leaf aphids.

Sevin is not only useful on lawns (for earwigs, armyworms, fleas, leafhoppers, millipedes, sod webworms, and mosquitoes, as well as chinch bugs), but against insects on home-grown tree fruits, most of the small fruits and nuts, vegetables, rose bushes, garden flowers, and shrubs at 1.0 to 2.0 lb. of 50% wettable powder per 100 gallons of water, or 1 qt. 4 Flowable per 100 gallons of water.

For best results against chinch bugs, lawns in infested areas



Experiments with Sevin insecticide on a variety of ornamentals and shade trees have demonstrated it to be effective for controlling most common tree pests.

should be sprayed about the time the young insects are hatching, when temperatures rise into the 70s. One and one-quarter pounds of 85W in 150 to 200 gallons of water will treat 5000 sq. ft. (or 2 lbs. of 50W in the same amount of carrier). This is the same as 1 lb. actual Sevin per 5000 sq. ft.

According to Dr. Louis Pyrenson of the Long Island Agricultural and Technical Institute in Farmingdale, N.Y., best results are obtained when the lawn is mowed and given a good wetting with a sprinkler before applying the insecticide. This brings the bugs to the lawn surface. After application, the turf should be wet down with clear water.

One application of Sevin is capable of controlling an entire generation of chinch bugs. Applications may have to be repeated after 2 or 3 weeks if young of a new generation begin to appear.


CAs who spray flower gardens for insect pests can use Sevin on a wide variety of nonwoody plants including asters, begonias, mums,

dahlias, gladiolas, iris, marigolds, orchids, roses, and zinnias. The shrubs include andromedas, azaleas, camelias, euonymus, hydrangeas, laurel, lilacs, rhododendrons, rose of Sharon, and yews, at 1 lb. active ingredient per 100 gallons of water.

On ornamental trees the list includes arbor vitae, ashes, birches, bamboos, cypress, dogwoods, elms, firs, hemlocks, hickories, junipers, magnolias, pines, sassafras, red-buds, and others. But avoid using Sevin on Boston or Japanese ivy, since it may injure these plants, experimenters discovered.

Sevin poses slight hazard to desirable birds and other animal species, according to James K. Keith, a research biologist with the U. S. Fish and Wildlife Service of the Department of Interior. "Our work to date with Sevin insecticide indicates that it has a low potential hazard to wildlife and that it is one of the more desirable pesticides for use where wildlife values are of consideration," Keith has stated.

HOARY CRESS
(*Cardaria draba*)



Aliases of hoary cress are: perennial peppergrass, hoary pepperwort, and white-top. This alien weed of the mustard family is a perennial which reproduces by seed and by sprouting of rootstocks. It is found in New England, and some mid-coastal states, through the Midwest to the Pacific. Hoary cress seems best adapted to cool climatic conditions such as those found in the Rocky Mountain region.

Stems (4) of hoary cress grow singly straight up, 1-2 ft. tall, branching only at the top. Stems and leaves are hairy. Lower leaves have obvious petioles (stalks) and are spatula shaped with slightly scalloped margins. Higher up on the stem, leaves lose their petioles and appear to clasp the stem (5).

Flowers on separate stalks are arranged in clusters at nearly the same height on the ends of all branches. Flowers are typical of mustards (family Cruciferae), with four tiny white petals in the shape of a cross. As branches grow, flowering at the ends continues; flowers still bloom after lower seeds have matured (1).

Seed pods (2) are heart-shaped, bulbous, attached to the flower stalk at the broad end of this heart-shaped pod. A pod holds two oval seeds, each $\frac{1}{16}$ inch long. Seeds are reddish brown and have a rough texture (3).

Roots of hoary cress are slender and many-branched, sometimes penetrating soil to a depth of 10 feet. It is the ability of roots to regenerate tops which makes this weed so difficult to control (6).

Other plants, much easier to control, which may be confused with hoary cress are members of its own family; all have white flowers. Peppergrass, *Lepidium virginicum*, is not a perennial, is easily pulled up, does not have clasping leaves, and has circular flattened seed pods. Field peppergrass, *L. campestre*, does have clasping leaves, but they are arrow-shaped (saggitate); seed pods are more boat-shaped.

Pennycress, *Thlaspi* sp., looks somewhat like hoary cress but has no hair on the leaves and has very flat and rounded pods. Shepherd's purse, *Capsella bursa-pastoris*, has saggitate, clasping leaves and seed pods are heart-shaped, but they are attached at the narrow end of the pod and are more flattened than those of hoary cress.

Repeated applications of 2,4-D will control hoary cress if applied at the late vegetative to early bud stage of growth. Combinations of leachable soil sterilants such as sodium chlorate, fenac, TBA, boron compounds, substituted urea compounds, and triazines may be used effectively. These herbicides also kill most other plants.

Where sterilization may cause undesirable erosion, planting over with vigorous grasses after application of a contact herbicide eventually controls hoary cress especially with well-timed 2,4-D treatments.

Prepared in cooperation with Crops Research Division, Agricultural Research Service, United States Department of Agriculture, Beltsville, Maryland.

(DRAWINGS FROM NORTH CENTRAL REGIONAL PUBLICATION NO. 36, USDA EXTENSION SERVICE)

"Industry Hour" to Highlight Fla. Turf Show, Aug. 27-29

New event added to the 11th Annual Florida Turf Conference, scheduled for the University of Florida, Gainesville, Aug. 27-29, will be an "Industry Hour."

Featured will be chemical and fertilizer products for turf. Each firm that exhibited these products at the Second Annual Trade Show (See *Weeds and Turf*, July 1963, p. 16), will be eligible to participate. Plans are now being made to continue this event each year, with a new theme being selected each time.

Although final program details and confirmation of speakers are not yet completed, general outline of the Conference will follow those of recent years. "Insects" will be theme of the general sessions, which open Tuesday afternoon, August 27.

Professional discussion sessions will be held Wednesday morning, with sections devoted to golf turf; playfields, parks, and cemeteries; retail dealers and garden suppliers; horticultural sprayers and lawn service agencies; and turf nurseries. Tour of turf plots and a barbecue, with the banquet that evening, will conclude Wednesday's sessions.

Conference will close Thursday morning, with a general session devoted to 1963 Turf Research Reports.

For more information on the program, or advance registration forms, write to Richard C. Wallace, Executive Secretary, Florida Turf-Grass Assn., P.O. Box 5284, Jacksonville, Fla.

Velsicol Has Bandane Booklet

Information on Bandane, a new pre-emergence crabgrass control from Velsicol Chemical Corp., is available to CAs in a new pamphlet from that firm.

Bandane has a mild selective action, and will not harm seeds, seedlings, or desirable grasses, according to the company. In addition, one application will eliminate ants and grubs from turf for several seasons, Velsicol claims.

For more information, write to the company at 330 East Grand Ave., Chicago 11, Ill.

Protect ornamentals and shade trees from mites and diseases

KELTHANE® Controls mites on ornamentals, nursery stock and shade trees.

DITHANE® Wettable powder fungicides control certain diseases on ornamentals and turf. For example; black spot, botrytis, anthracnose, leaf blight, leaf spot, stemphylium and various lawn diseases.

KARATHANE® Controls powdery mildew on ornamentals.

TRITON® B-1956 Spray additive that imparts maximum effectiveness to insecticides and fungicides.

For full information on how these Rohm & Haas products can be used to protect gardens and parks, fill in and send the attached coupon.

ROHM & HAAS
PHILADELPHIA 5, PA.



**Agricultural & Sanitary Chemicals Department
ROHM & HAAS COMPANY, PHILADELPHIA 5, PA.**

Please send me full information on Kelthane, Dithane, Karathane and Triton B-1956, their application and dosage schedules for use on ornamentals and shade trees.

NAME _____

COMPANY _____

ADDRESS _____



Get full protection... full vision...comfort... with Acme Gas Masks

There are a lot of things you'll like about an Acme Gas Mask. For example, you'll like the full protection that guards your face and eyes. And you'll like the Acme's Full-Vision Design. It lets you see more than any other mask. (Ask for Visionscope Proof.)

You'll breathe easier too. That's because our large Inhalation-Exhalation Valve is designed for minimum resistance. This adds to your comfort. So does the special soft-but-durable rubber we use for the face piece.

For "proof of the pudding," try an Acme Mask. Call your distributor. He'll be glad to demonstrate. Or . . .

MAIL THIS COUPON NOW. CHECK BULLETINS YOU WANT
Distributors: A few important territories available.

Acme Protection Equipment Co., 1225 Kalamazoo St., South Haven,
Please send Bulletins I've checked: Mich.

How to Select a Gas Mask New Protection Guide to 500 Gases New Facts about vision

Name _____

Company _____

Street _____

City _____ Zone _____ State _____





This quartet of tree experts paused to examine results of the severe midwestern winter while attending the International Shade Tree Conference Ohio Chapter meeting in Cleveland. Left to right are John Michalko, Cleveland Commissioner of Shade Trees; Dr. L. C. Chadwick, group secretary-treasurer; Dr. Paul E. Tilford, executive secretary of the National Arborists Association; and George Kase, Cincinnati Park Board, president of the Ohio organization.

Ohio Shade Tree Conference Laments Severe Winter and Hot, Dry Summer

"Because of a tough winter, which caused us to lose a lot of young trees, and the current drought in the Midwest, we're having a hard time of it," John Michalko, Commissioner of Shade Trees for the City of Cleveland, told delegates to the Ohio Chapter of the International Shade Tree Conference.

Nearly 80 arborists, tree company supervisors, and utility foresters gathered July 10 at the Cleveland Municipal Shade Tree Headquarters for the annual meeting, which featured tours of outstanding tree plantings in the Cleveland area, and a demonstration of some new tree maintenance equipment.

Michalko's remarks preceded a report from Dr. L. C. Chadwick, secretary of the Ohio group, who reported that the Buckeye tree organization, with a membership of 191, is now second in size only to the California chapter.

During the business session, chaired by president George Kase, of the Cincinnati Park Board, delegates voted to incorporate the Ohio chapter.

After the equipment show, the Ohio treemen moved to the Cleveland Museum of Art for additional talks on the role of tree personnel in urban redevelopment.

"There used to be a conflict between municipalities and electric companies over trees," Lee Howley, vice president and general counsel of the Cleveland Electric

Illuminating Company, told the assembly.

This conflict is being resolved, however, through such programs as CEI's Master Tree Plan, which helps municipalities plan tree plantings which will not interfere with utility lines, traffic, or sidewalks and sewers, and which will still enhance the esthetic character of a city's neighborhoods.

Next meeting of the Ohio Chapter will be in the last week of January in Columbus. Exact dates will be announced later, Dr. Chadwick told *Weeds and Turf*.

Meeting Dates

Alabama Nurserymen / Southern Nurserymen Annual Convention, Hotel Admiral Semmes, Mobile, Aug. 4-6.

International Shade Tree Conference, Royal York Hotel, Toronto, Ontario, Canada, Aug 4-9.

Rutgers Turfgrass Field Days, Rutgers University College of Agriculture, New Brunswick, N.J., Aug. 7-9.

11th Annual University of Florida Turfgrass Management Conference, University of Florida, Gainesville, Aug. 27-29.

Nurserymen and Growers Annual Short Course, University of Florida, Gainesville, August 28-30.

Midwest Turf Field Day, Purdue University, Lafayette, Ind., Sept. 9 (repeated Sept. 10).

70th Annual Farm Equipment Institute Convention, Roosevelt Hotel, New Orleans, La., Sept. 29-Oct. 2.

International Shade Tree Group Has Varied Convention Program

A program designed to satisfy tree specialists from all phases of the industry highlights the 39th Annual International Shade Tree Conference this month in Toronto, Ontario, Canada.

Nearly 800 delegates are expected to register during the 6-day tree seminar which runs from August 4 to 9 at Toronto's Royal York Hotel.

Much of the program is devoted to discussions of tree pest control, including talks on non-infectious tree diseases. Moderated by Dr. Spencer H. Davis, ISTC vice president, these tree ailment talks include "Ash Dieback in the Northeast," "Maple Decline in the Northeast," and "Sweet Gum Blight."

Role of tree surgery in modern arboriculture is the subject of an address by C. L. Wachtel, Wachtel Tree Science and Service Co., Wauwatosa, Wis.

Concepts of soil aeration and the relationship of this process to tree growth are being examined Monday by George Yelenosky from the Department of Botany, Duke University, Durham, N. C.

Split sessions are scheduled for Tuesday to better accommodate the varied interests of conference delegates. In the series of talks geared to members of the National Arborists Association, markets and public relations are covered.

"Shade Tree Spraying—the Tree, the Public, the Arborist" is the title of a talk by Henry F. Davis, Lowden Tree Specialists, Inc., Needham, Mass. "New Markets for Tree Service" is being presented by Robert Felix, Harder Tree Service, Inc., Hempstead, N. Y.

Also on Tuesday, those interested in municipal arboriculture may attend discussions chaired by John G. Michalko, chairman of the ISTC's Municipal Arborists Committee.

A significant talk on the real facts behind pesticide usage, presented by Dr. George C. Decker, distinguished entomologist of the Illinois Natural History Survey, Urbana, is in the offing Thursday.

A detailed report of the conference will appear in *W&T* next month.