

New Aspon™

gives you
chinch bug control
that's better
than the best!



Better than the best lawn chinch bug control you've been able to offer until now. That's what you give your customers when you use Stauffer's new ASPON insecticide.

Chinch bug destruction is a growing lawn and turf problem around the country. This insect is causing wide-spread damage not only in Florida, but along the Gulf Coast, through the Middle Atlantic states and into southern New England.

This steady spread has stimulated intensive research at Stauffer. Most effective of many materials developed in this work has been one which was code-named "ASP-51." This powerful new pesticide has been shown, in numerous tests and comparisons, to give control superior to that provided by any other material. ASP-51 is now known as "ASPON", and is now available to commercial spraymen.

Here are the advantages you get with ASPON:

- Highest percentage control—close to 100% within 48 hours after application
- Longest residual control—greater than that of any other material tested
- No pesticide resistance—control of all chinch bugs, including those resistant to other insecticides
- Low hazard—more safety in use and after use than possible with other leading chinch bug control materials

Remember, ASPON is now available to commercial spraymen. Offer the experienced application of the most effective chinch bug control available: ASPON. Get all the details—Fact Sheet, specimen label, summary of test data, chemical, physical and toxicological properties. Write or call the nearest Stauffer Chemical Company office: 380 MADISON AVENUE, NEW YORK 17, N. Y. • P. O. BOX 5388, TAMPA 5, FLA. • P. O. BOX 7222, HEIGHTS STATION, HOUSTON 8, TEX. • 1300 E. DAVENPORT ST., WESLACO, TEX. • P. O. BOX 509, NORTH LITTLE ROCK, ARK.

™ Stauffer's trademark for an insecticide

STRAIGHT?

IS your ad approach the shortest distance between sell and buy?

Is your sales promotion plan aimed straight at the multibillion dollar weed control and turf spraying market?

Or are you way off center, calling on the wrong people with your monthly ads —

Reaching a small segment of the bulk buyers —

Spreading yourself thin just to reach an occasional prospect?

You need a vertical book.

Weeds and Turf is a vertical book.

Straight as an arrow, your monthly sales message streaks towards target.

And what a target!

Weeds and Turf's 10,000 circulation reaches every facet of weed control, and lawn and ornamental spraying.

Contract applicators.

Highway officials.

Railway and utility rights-of-way supervisors.

State and local officers.

That is, if they're interested in how to control or enhance vegetation.

Otherwise, they read something else.

These straight-thinking men have one thing in common: they want to

learn more about outdoor pest control.

That's their job.

They buy lots of chemicals and equipment to help do it.

Want them to buy yours?

Then make sure your sales message goes straight to the men who do the buying.

Make sure your sales efforts don't stray off target, waste your time and money.

Weeds and Turf is a truly vertical book. It's as straight as they come.

Write us for a quiverful of facts: rate card, brochure, market bulletins.

Do it straightaway!


WEEDS and TURF
PEST CONTROL

Get dependable

insect control on lawns with

DIAZINON[®]

INSECTICIDE




With more and more PCO's expanding their services, insect control on lawns has become big business.

Indoors, Diazinon is the standard insecticide for control of roaches. Outdoors, this outstanding insecticide is the standard for control of lawn chinch bugs, lawn moths, brown dog ticks, Bermuda mites, Rhodes grass scale, fleas, chiggers, ants and many other pests. It is recommended by agricultural authorities and lawn and turf experts everywhere.

Diazinon gives quick kill of insects attacking lawns, and provides long residual protection, with safety in use, and economy in the finished spray. Follow recommended rates and directions on the label.

Don't take chances with call-backs—depend upon Diazinon for lawn insect control. Free literature will be sent upon request. Address Department PC-37.

CREATORS OF CHEMICALS FOR MODERN AGRICULTURE 

GEIGY AGRICULTURAL CHEMICALS • Division of Geigy Chemical Corporation • Saw Mill River Road, Ardsley, New York

VERTICALLY . . .



HORIZONTALLY . . .



**YOU ALWAYS GET
PERFECT PERFORMANCE . . .
WITH THE DEPENDABLE . . .**



**HARDIE AERO-MIST
SPRAYER DUSTER**

Here's the one unit that gives you just what you want . . . when you want it. The famous Hardie Aero-Mist Sprayer. It embodies the recommendations of State & Federal authorities; tree service organizations and foresters who sought in one unit a high-velocity ample air volume sprayer for any type job, along with low price.

With its high velocity, 150 mph, it is of special value in the treatment of trees infested with the bark beetle. Public Health officials also use the Hardie for mosquito control. With the attachment of the Hardie Duster, you have two machines in one. You can use either spray materials; dust or granular materials. Write for full details.

**HARDIE
SPRAYERS**

AMERICAN PULLEY CO. DEPT. PC
4200 WISSAHICKON AVE. PHILA., PA.

WEEDS and TURF

PEST CONTROL

A SECTION OF PEST CONTROL MAGAZINE

July, 1963

Features

How to Control Pests
of Ornamentals
By Dr. Donald Schuder W-8

Improved Management Techniques
Stressed at 2nd Fla. Turf Show
By Walter D. Anderson W-16

"Concepts of Forest Entomology"
Book Review W-17

Freeman Explains Federal Law
To Pesticide Use Investigators. W-21

Departments

Know Your Species W-18

Meeting Dates W-20

W & T Mailbox W-20

Literature You'll Want W-21

Trimnings W-22

Published Monthly by
TRADE MAGAZINES, INC.
1900 Euclid Avenue
Cleveland 15, Ohio

JAMES A. NELSON
Publisher

CHARLES D. WEBB
Editor

WALLIS W. WOOD
Editorial Assistant

DAVID E. SCHNEIDER
Staff Biologist

R. J. HOFFER
Circulation Manager

Advertising Representatives
National Headquarters
1900 Euclid Avenue
Cleveland 15, Ohio
Phone: Area Code: 216 + 771-4169

New York City
Billingslea & Ficke
420 Lexington Avenue
Phone: Area Code: 212 + LExington 2-3667

Single Copies: 35 cents

Annual Rate for
12 Monthly Issues: \$3.00

Contents of this Issue © Trade Magazines, Inc., 1963

The first year

This July issue marks the first anniversary of *Weeds and Turf*, a publication conceived as a service to America's contract applicators.

It would be false modesty to deny how successful the fledgling magazine has been. But it would be equally incorrect if we failed to take note of all the people who've helped spur on this successful year.

Readers have been most generous with their letters, most of which contain very constructive suggestions.

Suppliers have offered their counsel, and have given important support through our advertising pages, which of course make this kind of venture possible.

Researchers from universities, from government agencies, and from manufacturers have written many articles for us (or more accurately, for *you*); these same men have offered technical guidance, and have reviewed manuscripts by our research staff.

In the past year, several original staff-written articles were presented. Approved after many weeks of study, these papers on turf diseases, pre-emergent herbicides, soil sterilants, lawn insects, and the definitive study of the chinch bug, are all examples of the editorial policy which has gained the confidence of the 10,000 who receive *W&T* each month.

In the coming year, more of such articles are planned, along with features on legislation, business management, advertising, equipment maintenance, and dozens of technical articles on turf, trees, and weeds.

The widely praised weed identification box will be continued.

We hope that these future issues will find the same hearty welcome that marked last year. We'll be glad to get any suggestions about the content of *Weeds and Turf* as it heads for its *second* anniversary.

In the meantime, we want to say a sincere thank you to our readers, our advertisers, and to our authors for making the first year a success.

MALATHION

from

PENICK

in a variety of
formulations for

---insects on ornamentals
---outdoor fogging

write today for free catalog on

PENICK

MALATHION FORMULATIONS

Have You Tried

**NEW PREMIUM GRADE
MALATHION —**

Also Available from

PENICK

S. B. PENICK & COMPANY *Pest Control Supply Department*
100 CHURCH ST., NEW YORK 8 • 735 W. DIVISION ST., CHICAGO 10 • 4161 BECK AVE., ST. LOUIS 16
LOS ANGELES SAN FRANCISCO PORTLAND, ORE.

FOR FAST SERVICE ON . . .

Chlordane • DDT • Dicapthon • Dieldrin • Dethmor® Warfarin
Lindane • Malathion • Powdered Pyrethrum • Pyrethrum Formulations

CALL PENICK FIRST! ORDERS SHIPPED WITHIN 24 HOURS

Penick can supply most of the pesticides you use . . .
under a streamlined, personal-service system that saves you
time and bothersome paperwork.

We provide a handy check-off order form and price list
to simplify ordering. You get just one invoice to check.
And deliveries are always prompt . . . often overnight . . .
to minimize your inventory needs and eliminate irksome
follow-up calls.

You'll also find our monthly PCO Newsletter a big help
in keeping abreast of new developments.

Why not try us on your next order?

easy turf and weed control—
with a famous

FONTAN



**THE R.5 FONTAN FOR HEAVY DUTY
WEIGHT 37 LBS.**

A self-contained unit, the Fontan has jets to interchange for misting or spraying, another attachment to interchange for dusting. Designed for versatility, dependability and safety, the Fontan has metal frame and padded straps for comfortable operation.

FONTAN + MALATHION = ECONOMY

The Fontan adjusts to allow low-volume spraying with less dilute liquid and a higher concentration of Malathion or other chemical. Both Fontans offer complete portability, choice of droplet size, easy maintenance, safe fuel injection, corrosion-resistant plated parts.



**THE R.6 FONTAN FOR SMALLER JOBS
WEIGHT 24 LBS.**

A torch interchanged with the spray head and nozzle converts either Fontan into a flame thrower for weed killing, snow removal, burning off paint, heating tar for roofs and street repairs, and numerous similar tasks.

SOUTHERN MILL CREEK PRODUCTS COMPANY, Inc.
BOX 4297, 7C-3, TAMPA, FLORIDA

Please send further information on:

R.5 FONTAN R.6 FONTAN MALATHION

NAME _____

ADDRESS _____

CITY _____ ZONE _____ STATE _____

NATIONAL
DISTRIBUTORS OF
INSECTICIDES,
EQUIPMENT AND
WEED KILLERS



Southern MILL CREEK PRODUCTS COMPANY, Inc.

BOX 4297 • PHONE 253-0191 • 1906 N. ARMENIA AVE., TAMPA 7, FLORIDA

When Writing to Advertisers Please Mention WEEDS AND TURF



These white pine trees have been completely defoliated by bagworms, a common leaf-feeding pest of many trees and shrubs.

How To Control Pests of Ornamentals

By DR. DONALD L. SCHUDER

Associate Professor, Department of Entomology
Purdue University, Lafayette, Ind.

THE PRESENT trend of the American people for suburban living has opened a new field of service, that of insect control of lawns and ornamental plantings.

Several pest control firms in the East and Midwest are already established in this field, and are finding it an excellent source of income in association with their normal household pest control operations. Other companies should investigate the possibilities of expanding into this line of work, while the demand is still growing and there is room for more operators.

Before going into this new business, PCOs and CAs should give some thought to the potential in their areas and the ability of each firm and its employees to take on this new line of endeavor. A change would necessitate knowledge of a new group of insects, new and often larger equipment, and new methods of operations. Most of the actual work is done during the spring and summer months, which are already the busy season for the PCO. Many of the firms in this business have trained personnel who specialize in the care of lawns and ornamental plantings.

In order to understand the scope of this business and the possibilities of success in it, the operators should understand the insects and plants involved, insecticides and equipment needed, and numerous other problems. This article briefly

summarizes some of the pests, and lists methods for their control. The topic is divided into *Sucking Insects*, *Chewing Insects*, *Tree Boring Insects*, *Leafminers*, *Galls*, and *Mites*.

Sucking Insects

This group includes the various aphids, lacebugs, plant bugs, and leafhoppers which weaken plants by sucking out the plant sap. These insects frequently inject into the plant tissues secretions which injure or kill the plant cells.

Aphids or *Plant Lice* — Nearly all ornamental trees and shrubs are subject to injury by aphids. These insects are small, soft-bodied insects which may vary in color from green to shades of red, brown, or black. The insects may attack both aerial and subterranean portions of the plant, but are commonly seen clustered on the new growth or the underside of leaves.

When the population of aphids is heavy, they may cause the twigs and leaves to become curled and twisted, flower buds hardened, and flowers malformed. Aphids secrete a sweet, sticky substance called "honey-dew." The material is attractive to ants and flies and is a nuisance on cars, chairs, tables, and the like which are beneath infested plants. A black, sooty mold may grow on the honey-dew.

Aphids may be controlled during the summertime by applying nicotine sulphate and soap, lindane, malathion, thiodan, or diazinon. The highly poisonous parathion and tetraethyl pyrophosphate (TEPP) are also highly effective against aphids, but are not recommended for use by anyone unfamiliar with handling them. Lindane or BHC is preferred for woolly aphids.

A forceful stream of water from

Last December, *Weeds and Turf* presented its first article on ornamental pest control, written by Dr. L. C. Kuitert of the University of Florida. This article by midwesterner Schuder gives another slant on the important ornamentals market, and offers more control hints for successful contract applicators.

the hose will help to remove sticky honey-dew from plants. Since ants frequently are involved in transporting and tending aphids, it is sometimes helpful to kill the ant colony by treating their nesting areas with chlordane.

Lacebugs — Many trees and shrubs are attacked by species of lacebugs. The insects are about $\frac{1}{8}$ inch in length, dark in color, with transparent, lacy wings. Both the adults and nymphs feed on the underside of leaves. They extract so much plant juice that the leaves become pale and mottled with white splotches. The lower surface of the leaves are spotted with black and brownish dots.

About the last of May, apply lindane, malathion, DDT, or nicotine sulfate. The new systemic insecticide, dimethoate, is also effective. Be sure to treat the underneath side of the leaves. DDT may induce a mite problem and should be combined with a good miticide.

Leafhoppers — Leafhoppers are small, wedge-shaped, active insects which run, hop, or fly when disturbed. These sucking insects are usually yellow or greenish in color, while a few kinds are striped. Leafhoppers feed on the underside of the leaves, causing them to become light in color and marked with small, white dots on the upper surface. Often, the cast skins of the nymphs remain attached to the lower surface of the leaf and may serve as an aid in identifying the cause of the injury. Some species of leafhoppers are known to transmit tree diseases such as phloem necrosis on elm.

For control, apply DDT about the middle of June, or whenever the insects are discovered. Be sure to direct the spray to the underside of the foliage where the leafhoppers feed.

Plant bugs — These small, shield-shaped insects are active and run or fly freely when disturbed. Both the adult and nymphal stages have sucking mouthparts. Several species of this group inject toxins into the plant, causing the parts attacked to wither and die. Terminal buds, shoots, and blossoms are commonly affected.

DDT is the best insecticide for controlling plant bugs. The best



Dr. Donald L. Schuder, assistant professor of entomology at Purdue University, Lafayette, Ind., details the role CAs and PCOs can play in treating for pests of ornamentals.

results are usually obtained by spraying early in the morning, because the insects are less active at that time. Preventive measures are important, and include the removal of weeds, trash, and debris.

Scale Insects — Scale insects are serious pests of many ornamental plants, but unlike the other insects which suck sap from the plants, they are inactive during most of their lives. Scale insects are quite small, usually dark in color so that they are difficult to detect. Infested plants appear unhealthy, grow poorly, and the foliage may be small and chlorotic. For ease in identification, scale insects may be grouped into three divisions: (1) armored scales, (2) soft or unarmored scales, and (3) the mealybugs.

Armored Scales

Armored scales are the most common type of scale insect found on trees and shrubs. They secrete a hard, waxy covering over their bodies. The insect feeds and grows beneath this protective covering or scale. The scale may be circular, oblong, or pear-shaped. The armored scales are quite small, varying in size from $\frac{1}{2}$ to $\frac{1}{8}$ inch in length or diameter. A typical life cycle may be illustrated with the gray race of the oystershell scale. These insects overwinter in the egg stage, beneath the female scale of the previous season. These eggs hatch the latter part of May or the first of June. The young scale, called a nymph or crawler, moves about over the host plant until it finds a suitable place to

feed. It then inserts its needle-like mouthparts and draws its liquid food supply from the phloem tissues of the plant. At this time, the insect loses its mobility and goes through a series of five molts, casting off its skin before it reaches maximum size and sexual maturity. The males emerge as winged adults and fly about in search of the females. Eggs are laid underneath the female scale and the cycle is repeated the following season. Other species of scales have multiple generations and the cycle just described may be repeated several times each year. In some species, no males are known and reproduction is parthenogenic.

Species having one generation a year can be controlled by sprays in early June. This group includes Putnam scale, pit-making scale of sweet gum, scurfy scale, and the gray race of oystershell scale.

Species having two generations per year can be controlled by sprays in early June and again in late July. This group includes such common species as the brown race of oystershell scale, pine needle scale, and the juniper scale.

Multiple generation species frequently require sprays at monthly intervals for adequate control. Some of the species included in this group are the San Jose scale, euonymus scale, and the tea scale.

Unarmored Scales

Soft, unarmored, or lecanium scales differ from the armored scales and have their waxy secretion as an integral part of their body. The body shape may be flattened, slightly convex, or hemispherical. They are larger than the armored scales, varying from $\frac{1}{12}$ to $\frac{1}{2}$ inch in length. The eggs are laid beneath the female's body and hatch in one to three weeks. Development is slow and may require almost a year for a single generation. The nymphs of several of the soft scales feed on the leaves during the summer and move back to the twigs in the fall. Most species produce honey-dew.

Some of the unarmored scale species require spraying in June. This group includes: Terrapin scale, hickory lecanium, cottony maple scale, Fletcher's or the taxus lecanium, European elm scale, European fruit lecanium, and

spruce bud scale, among others.

Some of the other unarmored scale insects require spray applications in mid-August. Some common examples are the tulip tree scale, magnolia scale, and oak kermes.

Mealybugs


The mealybugs are soft-bodied insects and usually are covered with a powdery, cottony-waxy material. Unlike the scale insects already discussed, the mealybugs have legs and are able to move about throughout their entire life. They vary in size from about $\frac{1}{5}$ to $\frac{1}{3}$ inch in length when full-grown. The female mealybug deposits her eggs in a white, cottony sack. The species noticed on ornamental trees and shrubs produce two or more generations each year. Some common mealybugs are the grape or taxus mealybug and the juniper mealybug.

Control of Scale Insects

The control of scale insects is often difficult because of failure to obtain adequate spray coverage. To be effective, the spray material must come in contact with the insects. Care should be taken to see that the spray is evenly applied to all parts of the plant. The addition of a wetting agent to the spray mix will help wet the insect and insure penetration of the insecticide through the insect's protective covering.

Scale insects may be controlled during both the dormant period and the growing season. During the dormant period, apply dormant oil, lime sulfur, or the dinitro

Oystershell scales have infested these apple-tree twigs. Large insects are the mature females, while the smaller ones are the young, newly settled crawlers.



compounds. When the scale insects occur on oil-susceptible plants, use superior or summer oils. DN and liquid lime sulfur will discolor paint and stone and should be used with caution around home plantings.


Summer crawler sprays should be timed to coincide with egg-hatch. The most commonly available and effective material for summer application is malathion. The insecticide Sevin has been found to be very effective against many of the lecanium scales. Other insecticides which have been shown to be effective against the scale crawlers are Ethion, Trithion, phosphamidon, Guthion, demeton, phorate, diazinon, chlorthion, and Cygon.

Chewing or Leaf-Feeding Insects

Leaf feeding or chewing insects include those species which consume solid foliage tissue. Members of this group may skeletonize the foliage or entirely defoliate the plant.

The Bagworm — The bagworm is a general feeder and may be found attacking the foliage of both deciduous and evergreen trees and shrubs. The bagworm is so named because the caterpillar spends its life inside a silken, spindle-shaped bag. The bag is usually adorned with bits of twigs and leaves from the host plant and looks so much like a part of the tree that it usually goes unnoticed until extensive damage has been done. The bagworm overwinters in the egg stage inside the old bag. The eggs hatch in late May and early June. The young caterpillars immediately spin bags and continue to enlarge them as they grow. When the bagworm larvae mature in late August and early September, the bags are about 2 inches long. The bags are then attached firmly to twigs or branches and the worms pupate and change into adults. There is only one generation each year.

For bagworm control when only a few small trees are involved, bagworms can be controlled by hand-picking the bags and burning them. This may be done at any time when the bags are discovered, but is most easily done during the fall and winter, or early spring. Spraying is the most effective method when the trees are large



Yellow-necked caterpillars, which usually feed in colonies on pin oaks, will throw back their heads and curl up the tip of the abdomen as a protective action when they are disturbed.

or numerous. The sprays are most effective when applied in early June, as soon as possible after the eggs hatch. Several insecticides are effective. Malathion is the most widely accepted material. Other materials which have been recommended include diazinon, toxaphene, and lead arsenate. The new insecticide DDVP is suggested for trial by professional sprayers.

Blister Beetles — Blister beetles are active, black, gray, or striped insects which appear suddenly in enormous numbers and strip the foliage from some species of ornamental trees and vines. The beetles are long, cylindrical, and have an unusual appearance because of their long legs. This group of insects usually appears in large numbers during the summer. They are difficult to control. The best control has been obtained by applying 5% granular dieldrin to the soil beneath the trees or vines. The insects are killed when they return to the soil in the cooler parts of the evening and night. Sprays or dusts of cryolite, chlordane, and DDT have also been suggested.

Leaf-feeding Caterpillars — Caterpillars, the immature stage of butterflies and moths, exist in a wide variety of sizes and colors. Some may be covered with hairs and spines, while others are entirely naked. They may feed singly or in large colonies. All species have voracious appetites and consume large quantities of foliage. Most species are easily controlled with DDT sprays. In the case of some of the webworms, it is essential to use high pressure, so that the foliage inside of the web is covered with the insecticide.

Leaf Beetles — "Leaf beetles" is a term applied to several species