

"Technical help,"

says Don Brabec, Building and Grounds Superintendent, United Air Lines. "The Greenfield representative is a professional. He shows a sincere interest and if we have a problem, we can count on him to help us solve it."



"No costly call-backs,"

says landscaper Bob Rynberk. "That's what we've saved using Greenfield Measured Release Lawn Food. We can count on it giving superior results. Our crews can work, not answer complaints."

Why is there a growing move to Greenfield products for grounds maintenance?



"Unique products for special jobs,"

shown by results at the Rose Garden in Chicago's Grant Park. Greenfield pioneered products for roses, shrubs, flowers and evergreens that prevent weeds and feed at the same time. They save time, money and worry.

"Labor savings,"

says Herb Buesching, grounds superintendent at Concordia Sr. College, Ft. Wayne, Indiana. "Greenfield's weed and feed products for ornamental plantings have reduced our maintenance time by as much as 50%."

Greenfield products are different.

They are developed through conscientious research at the Greenfield (Ind.) Laboratories of Eli Lilly and Company to do more.

For example, the original triple action crab grass killer, the first weed and feed products for ornamentals, and the first combination pre- and post-emergence crab grass killer came from the Greenfield Research Center.

Innovations like this can save you time and effort in grounds care. Crews will spend less time weeding, feeding and preventing insects. They'll have more time for other jobs. What's more, Greenfield products are dependable. They give superior results and they're priced right.

Besides all this, you get technical help from your Greenfield representative. He'll help you work out a grounds program to solve your particular problem and fit your budget.

You want to switch, too?



ELANCO PRODUCTS COMPANY, A DIVISION OF ELI LILLY AND COMPANY

WEEDS TREES AND TURF, January, 1965

Send for:

THE GREENFIELD COMMERCIAL LANDSCAPING PLAN

Elanco Products Company Box 1750, Dept. WT 1-65 Indianapolis 6, Indiana

Please send full information on the Greenfield commercial landscape products and the plan which best applies to the situation I've checked: Industrial Grounds Cemeteries Parks Professional Landscaping School Grounds Airports

NAME	
COMPANY	
ADDRESS	
	STATE

Dacthal . . . one of many Diamond chemicals for dependable weed control

DACTHAL® W-75

Dacthal W-75 is the new pre-emergent herbicide which gives effective control of many annual grasses and many broadleafed weeds. Reduction in hoe bills and increase in crop yields make it economical to use. Dacthal W-75 is available for use on various vegetable crops, cotton and strawberries.

DACAMINE®

The new oil-soluble, water-emulsifiable amine with proven effectiveness on difficult-to-control weed and brush species. This new material has the safety of amines with the killing power of esters.

READY-TO-USE HORMONE FORMULATIONS

CROP RIDER® — Weed-killer formulations

LINE RIDER[®] — Brush-control formulations

Proven six-pound acidequivalent low-volatile formulations are available for the thrifty buyer. Technical 2,4-D and 2,4,5-T chemicals for formulators.

DIAMOND INSECTICIDES

Diamond dependable insecticides have long been favored because of their uniform high quality. DDT and BHC technicals are supplied in many types and forms:

DDT — 100% technical, wettable powders, dust and emulsifiable concentrates.

BHC — 45% technical, 15% technical, 12T wettable powders and dusts, emulsifiable concentrates.

Diamond Research, which discovered Dacthal, is constantly seeking new and better agricultural chemicals, and working with formulators and agricultural chemists in the development of more efficient forms and application methods. We will be glad to work with you. For information on Diamond chemicals and

technical cooperation, write to Diamond Alkali Co., 300 Union Commerce Bldg., Cleveland, Ohio 44114.




Diamond Chemicals

When Writing to Advertisers Please Mention WEEDS TREES AND TURF


ARD HAS THE RIGHT SPRAYER FOR YOU



Aeromist Sprayer-Duster



Model 99 50-100 gal. Skid-Trailer



15-30 gal. Town & Country

Hardie has just the sprayer for your particular job, whether it be for mosquito control; combatting Dutch Elm disease; nursery work; contract lawn, tree & shrubbery service . . . Whatever your need may be . . . write to Hardie today, or call your local Hardie Dealer for more information or an on-the-spot demonstration.

HARDIE SPRAYERS 4200 Wisschickon Avenue, Dept. WTT Philadelphia 29, Pa.
🗌 Have a Hardie Salesman call on me.
Literature
NAME
ADDRESS
CITY STATE
A DIVISION OF

UNIVERSAL T AMERICAN CORPORATION

TREES and

January 1965

Volume 4, No. 1

FORMERLY WEEDS AND TURF

Features:

Surfactants—How They Increase Herbicide Action By Dr. E. E. Schweizer and Dr. C. G. McWhorter
How to Use Your Soil Test Kit12
Turfgrass Portraits VII: Bahiagrass By Dr. Robert W. Schery16
Mineral Content in Fertilizers Underscored at Oklahoma's Annual Turf Grass Assn. Conference Nov. 4-6 in Stillwater By Lee Stevens
New Turf Herbicides, Sessions for Applicators On Varied Oregon Weed Program in Salem Nov. 5-6

Departments:

Editorial: Time to Spruce Up 6	
Know Your Species: Green Foxtail	
Meeting Dates	
Letters	
Classified Ads	
Advertisers Index	
Trimmings	

JAMES A. NELSON Publisher

CHARLES D. WEBB Editor

DAVID E. SCHNEIDER **Staff Biologist**

D. BUNKIN **Circulation Supervisor**

Advertising Representatives National Headquarters 1900 Euclid Avenue Cleveland, Ohio 44115 Phone: Area Code 216+771-4169

> **New York City Billingslea & Ficke 420 Lexington Avenue** Phone: 212+532-1632

Chicago Peck & Billingslea, Inc. 185 North Wabash Ave., Suite 1809 Phone: 312+DEarborn 2-0292-93

West Coast The Maurice A. Kimball Co., Inc. Los Angeles: 2550 Beverly Blvd. Phone: 213+DUnkirk 8-6178

San Francisco: 580 Market Street Phone: 415+EXbrook 2-3365


WEEDS TREES AND TURF is published monthly by Trade Magazines, Inc. Executive, editorial, and advertising offices: 1900 Euclid Ave., Cleveland, Ohio 44115. Publication office: Corner of East North St. and Cadwallader St., Fostoria, Ohio. Send all correspondence to WEEDS TREES AND TURF, 1900 Euclid Ave., Cleveland, Ohio 44115. Single Copy Price: 50 cents for current issue; all back issues 75 cents each. Foreign \$1.00. Subscription Rates: U.S. and possessions, 1 year \$3.00; 2 years \$5.00. All other foreign subscriptions, 1 year \$4.00; 2 years \$7.00. Change of Address: Three weeks advance notice is necessary for change of address. Both old and new address must be given. Post Office will not forward copies. Third Class postage is paid at Fostoria, Ohio.

Contents of this Issue (Trade Magazines, Inc., 1965

Time to Spruce Up

This is SNOW MOLD

For the answer to it and 9 other common turf problems, see below:



1. Snow mold. For more effective prevention or cure apply *Velsicol*[®] *''2-1.''* The new *''2-1''* formula has greater wettability, more stability in suspension and less foaming action than any other comparable product.

2. Dollar spot. 3. brown patch. 4. melting out (Helminthosporium). 5. gray leafspot. Velsicol® Memmi .8EC controls all these turf diseases and others. Memmi .8EC is a *liquid* mercurial that's easier to handle—needs no constant agitation, won't clog or leave sediment in the tank.

6. Plantain. 7. knotweed. 8. clover. One application of *Velsicol Banvel®* D + 2,4-D will effectively kill these and practically all the other common weeds that are found in turf. Works in warm or cold weather. Works on leaves *and* roots for complete kill.

9. Crabgrass. 10. grubs. For pre-emergence control of crabgrass plus insect control apply Velsicol *Bandane® 15G* or *Chlordane*. Both kill crabgrass before it comes up, get rid of grubs and ants. No waiting to re-seed after Bandane 15G!

For a free catalog of these and other Velsicol turf chemicals, including Pestmaster[®] Soil Fumigant-1 and Heptachlor insecticide, VELSICOL just mail the coupon.



VELSICOL CHEMICAL CORPORATION 341 E. Ohio St., Chicago, Ill. 60611		
NAME		
ORGANIZA	TION	
ADDRESS_		
CITY	STATE	

An attempt to reduce the amount of paperwork government requires of businessmen has been initiated by an organization called the National Federation of Independent Business. While we know little about this group, the current enterprise seems to us to have merit, since most businessmen often are bogged down in a ceaseless stream of forms, documents, reports, and correspondence.

At the same time, however, businessmen, in an attempt to reduce this flow of paper, must not fail to retain the necessary records, the mandatory reports, and the indispensable letters which make up office procedures in this automated age.

What seems to us the important lesson learned from the Federation's endeavor is the rather obvious one, perhaps, that a businessman must constantly analyze the amount of paperwork going on in his organization in an attempt to eliminate the superfluous while retaining the necessary.

How long has it been since you took a long hard look at your files, and the stack of incoming and outgoing mail found in every executive's office? Is it time to spruce up your system?

For example, there's a new way to answer ordinary correspondence which many companies now find useful. When a routine letter of comment or inquiry arrives in the morning mail, one that requires perhaps a sentence or two in reply, many office workers now simply write in longhand, or type, a reply on the original letter itself, have the document photocopied by one of the low-cost machines now available, and return the original to the sender, keeping the copy on file. The amount of paper retained by this system is only half that in the old method.

Forms in use in businesses today are undergoing constant revision. Smaller companies will find the counsel of the business forms supplier useful in determining how paperwork can be minimized.

It's unwise to go off on a grand crusade to abolish all letters, acknowledgements, purchase orders, and the like. It's especially important for people in vegetation maintenance and control, those who are dealing with living organisms and sometimes toxic chemicals, to keep the necessary records as a safeguard to possible litigation. If you feel adrift in a sea of paper, perhaps a few hours spent in careful analysis of your systems will pay big dividends in helping spruce up your efficiency.

WEEDS TREES AND TURF is the national monthly magazine of urban/industrial vegetation maintenance, including turf management, weed and brush control, and tree care. Readers include "contract applicators," arborists, nurserymen, and supervisory personnel with highway departments, railways, utilities, golf courses, and similar areas where vegetation must be enhanced or controlled. While the editors welcome contributions by qualified freelance writers, unsolicited manuscripts, unaccompanied by stamped, self-addressed envelopes, cannot be returned.

In Culver City, Trees Stop Growing.



Culver City, California. Five hundred trees stopped growing in Culver City this summer. These trees were treated with MH-30T* in a city-wide test. You can see the results in the photograph above. The tree on the left was treated with MH-30T. It grew only a few inches while the untreated tree on the right grew several feet.

Park Superintendent Roy Wells, who headed the project, reports: We figure MH-30T saves us from \$6 to \$8 in pruning costs on every tree we spray. And that doesn't include the huge savings in brush disposal. Next year we'll be treating more than 4,000 evergreen elms and several hundred Ficus Nitida (fig) trees. The savings to the taxpayer will be enormous."

MH-30T

Read about a revolutionary new chemical that controls the growth of trees without harming them in any way.

A new chemical can save cities and utility companies millions of dollars a year by drastically reducing the need for large scale pruning. This chemical is called MH-30T.

MH-30T actually stops cell division in trees (this is what makes them grow). You can use it to control the growth of an entire tree. Or, just the section interfering with electrical wires. The part of the tree that is sprayed with MH-30T simply stops growing. But the branches get sturdier and keep on producing leaves.

MH-30T has already been applied successfully to thousands of trees. It has been proven effective on twenty different species of trees. In most cases, MH-30T eliminates the need for at least one large scale pruning a year. Usually more. This can mean tremendous savings. For more information, simply fill out and mail the coupon.

	MH-SOT IS 0.5. Rubber's registered travemark for its mareic hydra.
Chemical Division UNIROYAL U.S. RUBBER	U.S. Rubber, Chemical Division, Naugatuck, Conn- I understand that MH-30T is for commercial use only. Please send me illustrated brochure showing how MH-30T works, and how it can reduce pruning costs by one-third or more.
NAME	Christian 200 and a strength of the second second
FIRM	TITLE
ADDRESS	
CITY	ZONE STATE

When Writing to Advertisers Please Mention WEEDS TREES AND TURF

SURFACTANTS:

How They Increase Herbicide Action

By DR. E. E. SCHWEIZER

Research Plant Physiologist USDA Agricultural Research Service Stoneville, Miss.

and

DR. C. G. McWHORTER Research Plant Physiologist USDA Agricultural Research Service Stoneville, Miss.

SURFACTANTS are widely used with agricultural pesticides to facilitate application and achieve better control of pests. The reasons for the increased efficiency resulting from surfactants remain obscure. The role of surfactants is often confusing and the terms used by farmers and other nontechnical personnel to discuss surfactants add to this confusion. Some of the terms most commonly encountered, and used interchangeably, are activator, additive, adjuvant, detergent, soap, spreader, surface-active agent, surfactant, and wetting agent. The common assumption that these terms are synonymous has contributed to the confusion that exists in surfactant terminology. Of these, additive, adjuvant, detergent, surfactant, and wetting agent are most commonly used.


The word "surfactant" is derived from the term "surfaceactive agent" and is defined by the Weed Society of America as "a material which facilitates and accentuates the emulsifying, dispersing, spreading, wetting, and other surface-modifying properties of herbicide formulations." A "wetting agent" is "any compound which when added to a spray solution causes it to contact plant surfaces more thoroughly." Thus a wetting agent is not necessarily a surfactant. An "additive" is "any material that is added to the spray solution and is not necessarily a wetting agent or a surfactant." An "adjuvant" is "that which assists, aids, or modifies" and thus might be as descriptive as any other term for reference to most surfactant-wetting agent materials. The term "detergent" is "any cleaning agent or solvent such as water or soap." Although water may be a detergent, it is not a satisfactory wetting agent because water usually does a very poor job of wetting. Lest this seem confusing, it should be remembered that to wet simply means "to cover or soak with a liquid." Spraying with water does not necessarily insure wetting.

Surfactants come in a wide variety of types. Several thousand trade name surfactants are already available. However, numerous hydrocarbon nuclei and polar functional groups are possible, so that there is no practical limit to the possible variety of surfactants. These materials are generally classified as anionic, cationic, and nonionic, depending on the electrocharge of the surface-active group. Anionic surfactants contain negatively charged atoms or groups; cationic surfactants contain positively charged atoms or groups; and nonionic surfactants are neutral. Compounds of the anionic and cationic class can be mixed with nonionics but not with one another. Anionic materials are generally used in detergents for home laundering and cleaning agents. Many emulsifying agents are mixtures of anionic and nonionic surfactants. Although cationic surfactants find many uses, their major outlet has been in various germicidal preparations such as hair shampoos and baby soaps. A few surfactants, also being marketed. are referred to as "formulated"

materials. These products contain either an anionic, cationic, or nonionic surfactant plus other materials such as alcohol, fatty acids, etc., which are supposed to aid penetration and translocation. Nonionic surfactants have been the most widely used groups, for reasons discussed below, by Midsouth farmers.

Nonionic surfactant molecules consist of two major chemical groups. One group is fat soluble (lipophilic), water insoluble (hydrophobic), and nonpolar. The second group is water soluble (hydrophilic), fat insoluble, and polar. Because neither positive nor negative ions are produced in any quantity, these surfactants have advantages over anionic and cationic surfactants. Most nonionic surfactants are not subject to hydrolysis by acidic or alkaline aqueous solutions. They do not form salts with metal ions, which make them equally effective in hard and soft waters. Because of these advantages, nonionic surfactants have received major emphasis in herbicide-surfactant research.

The more common hydrophobic portions of nonionic surfactant molecules are derived from the hydrocarbon portions of alkylphenols, aliphatic acids (especially fatty acids), and corresponding alcohols. The hydrophilic portion of the major types is conjugated chains of ethylene oxide. Long ethylene oxide chains are highly water soluble, probably due to the multiplicity of hydrogen bonds formed between the oxygen in the hydrophil chain and water of the solution. Examples of general types of nonionic surfactants are as follows:



The letter x in the hydrophil portion of the molecule denotes



Figure 1. Typical control of dallisgrass in bermudagrass from the application of 4 lb/A of disodium monomethylarsenate (DSMA) plus 0.5% surfactant.

the number of moles of ethylene oxide per mole of hydrophobe. Of the general types listed, only the alkylphenols have been retailed to farmers in quantity. The alkylphenols-including the the octyl-, nonyl-, and dodecylphenols-are important because of their ability to increase the activity of foliar-applied 3-(3,4dichlorophenyl) -1,1-dimethylurea (diuron) to weeds in cotton, and because of economy. Continued research will probably demonstrate many equally effective surfactants.

Effects of Surfactants on Herbicides

Aqueous foliage sprays of diuron without surfactant are relatively nonphytotoxic. If, however, a surfactant is included in the spray mixture, diuron is very phytotoxic at rates as low as 0.1 to 0.2 lb/A. Diuron-surfactant mixtures have been widely used in the Midsouth for weed control in cotton. The level of toxicity of these treatments is affected by the type of surfactant used. Ethoxylated nonylphenol surfactants containing 9-10 moles of




Figure 2. Diuron-surfactant mixtures prepared for the application of diuron at 0.022 lb/A in a total volume of 40 gallons of water per acre. Notice the decreased turbidity of the mixtures as the concentration of sodium lauryl sulfate is increased (McWhorter, 1963).

WEEDS TREES AND TURF, January, 1965

 Table 1. The Effectiveness of DSMA at 4 lb/A in Controlling Dallisgrass in Bermudagrass as Affected by the Addition of a Surfactant.

	No. of plants per sq. feet	Percent Reduction
Untreated	5.29	0
DSMA	2.25	58
DSMA + 0.5% ethoxylated nonylphenol type	0.53	90

ethylene oxide have generally provided maximum crabgrass control when applied in mixtures containing diuron. In these and other greenhouse studies at the Delta Branch Experiment Station, Stoneville, Mississippi, ethoxylated nonylphenol surfactants containing 15-30 moles of ethylene oxide were also effective in increasing diuron toxicity, but they were comparatively poor in increasing spray-mixture wettability.

Surfactants also enhance the activity of herbicides other than diuron. For example, surfactants increase the activity of 3,4dichloropropionanilide (DPA), used for the control of grasses in rice; 2,2-dichloropropionic acid (dalapon), used primarily for johnsongrass control; and disodium monomethylarsenate (DSMA), extensively used in the Midsouth for nutgrass and johnsongrass control. In addition, DSMA is very effective for controlling dallisgrass in bermudagrass (Fig. 1). The data in Table 1 shows that 0.5% (wt/v) of surfactant increased the control of dallisgrass by 32% over that of DSMA without surfactant.

Surfactants Can Increase Herbicidal Effectiveness

Surfactants can increase herbicidal effectiveness by: (a) improving plant coverage; (b) removing air films between spray and leaf surface; and (c) increasing foliar absorption and translocation. Surfactants improve plant coverage by reducing the surface tension and thereby increasing the total area of leaf surface that becomes wet. Increased wetting of a leaf surface results in the herbicide being spread over a larger area. Improved plant coverage and removal of air films between sprav and leaf surface can increase the

foliar absorption and translocation of herbicides.

Surfactants can also reduce the turbidity of herbicide spray solutions which may affect herbicide effectiveness (Fig. 2). It was originally believed that surfactants greatly reduced the turbidity of diuron suspensions by increasing the solubility of diuron in water. This is now known to be only partially true. Recent research with diuronsurfactant mixtures has suggested that reduced turbidity results from the formation of colloidal solutions and that diuron is incorporated into the colloid micelles by the process of "solubilization." Although additional research is needed to determine what role surfactants are performing in reducing turbidity, herbicidal activity has been greatly increased.

Even though phytotoxicity of many herbicides is increased by adding a low concentration of surfactant to the spray mixture, this may not always be beneficial inasmuch as a surfactant may eliminate species selectivity, thereby causing greater injury to a crop or turf. Therefore, judicious selection of surfactants is extremely important for the farmer, grower, or turf manager because the proper selection of surfactants and concentrations will aid in controlling resistant weed species. It will also lessen the possibility of damage to desirable plants and decrease the cost of herbicidal application.

Present, Future Surfactant Usage

The use of surfactants with herbicides to increase phytotoxicity has grown tremendously within the past four years in Mississippi. In 1960, approximately 64,000 pounds of surfactant were used with herbicides. By 1963 over 500,000 pounds of surfactant were used to control weeds in agronomic crops, pastures, and rights-of-way. This represents an increase of nearly 700%.

Although there has been a phenomenal increase in the use of surfactants to increase herbicidal action, it appears that this trend has only just begun. In the future a suitable surfactant will probably be chosen for herbicide application, just as a specific solvent is now selected for formulation of a pesticide. Indications are that such information will be available in the next few years. The problem will probably grow more complex. In the future not only will the surfactant be selected for the herbicide, but it will be selected for the particular crop and weed on which the material will be applied. Surfactants will greatly broaden the scope of herbicidal weed control, perhaps enough to surprise even those who are expecting the change.

Protect Young Trees Against Rabbits, Field Mice

Recently planted trees must be given protection during the winter months to prevent rabbits and field mice from eating the bark and causing death of the tree. This reminder is issued by the National Arborist Association, which adds that these precautions should be used to protect shrubs, too.

Ideal protection, which will last several years, is installation of a loose cylinder of ¼-inch hardware cloth, supported by stout stakes about the trunk of the tree. The cylinder should be buried a few inches in the soil to repel burrowing mice, and should be at least 30 inches high to prevent rabbits from reaching over the top.

Another method is to wrap the trunk of the young tree with aluminum foil, burlap, or treewrapping paper. Wrappings should be removed in the spring.

Chemical rodent repellents may also be used, which, to be effective, should be used according to manufacturers' directions on the container label.