

when is bluegrass a weed?



when it's annual bluegrass (*poa annua*) ...get rid of it now with **Betasan**[®]

SELECTIVE HERBICIDE

Poa annua is a weed which forms a relatively thick turf that can crowd out most perennial grasses. In the spring and fall, it grows so rapidly that lawns may become as much as 60% *poa annua*. Unlike Kentucky or Merion bluegrass, it may die and turn brown during the summer.



In some cooler areas, it behaves like a short-lived perennial because the seed germinates year round. But, in most parts of the country, the seed germinates in the fall and *poa annua* appears as a winter annual.

BETASAN, Stauffer's selective herbicide for use on lawns, gives you excellent control of *poa annua* when the seeds germinate. Application timing is important. Use it now—between mid-July and mid-September—before the new crop of *poa annua* emerges.

Send for your free copy of our BETASAN brochure and see your Stauffer distributor for full details. Stauffer Chemical Company, Agricultural Chemical Division, 380 Madison Avenue, New York, N. Y. 10017.



READ THE LABEL. HEED THE LABEL AND
GROW WITH STAUFFER CHEMICALS

Stauffer
CHEMICALS



HARDIE SPRAYERS

• YOU CAN'T MISS




Effective spraying begins and ends with a Hardie . . .

Just what kind of a sprayer do you need? . . . Well, whatever type, size, model or application, you can bet that Hardie . . . the "Sprayer-People" have it.

Without any obligation on your part, drop Hardie a line today . . . and tell them what you want to spray . . . and how much you have to spray.

We'll tell you just what you need, and we'll send you literature that will fully illustrate and describe the unit, or units that will do the job right and the price will be right too . . .

If you'd like a salesman to call, or if you'd like the name and address of the Hardie dealer nearest you, just let us know. Hardie Sprayers . . . 4200 Wissahickon Avenue, Phila. 39, Penna.



A DIVISION OF UNIVERSAL AMERICAN CORPORATION

5

REASONS WHY YOU SHOULD USE BAIRD & MCGUIRE

MALATHION



experience-tested **FORMULATIONS**

1. HIGHEST KILL-POWER
2. LOWEST COST
3. EASY TO APPLY
4. LOW TOXICITY & LOW ODOR
5. MULTIPLE INSECT CONTROL

BAIRD & MCGUIRE puts maximum kill-punch plus economy into their experience-tested Malathion formulations . . . PCO's have a wide choice of 50% emulsifiable, 5 lb. emulsifiable concentrate, 25% wettable powder and 90% solutions that are easy to apply (spray—granular—dust) with low toxicity, low odor, for indoors or outdoors . . . these formulations will solve all your mosquito and other insect control problems . . .

BUY FORMULATIONS OF EXPERIENCE . . .
BUY BAIRD'S

**Write today for complete
products price list.**

BAIRD & MCGUIRE
INCORPORATED
HOLBROOK, MASSACHUSETTS

SINCE 1910




CREATORS AND COMPOUNDERS OF THE FINEST
IN CLEANERS AND DISINFECTANTS SINCE 1910



SEED TREATMENTS

mercury based disinfectants for wheat, oats, barley, flax, sorghum, rice and cotton


METASOL



TURF FUNGICIDES

standard and custom fungicides — PMA, thiram, calomel, and cadmium formulations

METASOL



SOIL FUNGICIDES

oxyquinoline sulphate, benzoate and organo sulphur temporary soil sterilants

METASOL



FOLIAR FUNGICIDES

organo mercurials for apple scab and ornamental foliage diseases

METASOL

Quality insured with 20 years of research and manufacturing Know-how



METASOL



MERCK CHEMICAL DIVISION

MERCK & CO., INC. • 200 WAGARAW ROAD HAWTHORNE, NEW JERSEY • 07507

FORMERLY METALSALTS CORPORATION

WEEDS TREES and TURF

FORMERLY WEEDS AND TURF

August 1966

Volume 5, No. 8

Features:

Anthracnose of Sycamore and How to Protect Against It By Dan Neely	8
Pruning Mature Trees By Keith L. Davey	12
Where to Sell Those Wood Chips.....	14
Safety and the Tree Surgeon By Horace P. Bryan.....	16
Calculate Tree Canopy to Regulate Treatment Time with Mistblowers	20
Rainfall Influence on Preemergence Herbicides	22
Bluegrass Perils, All-in-one Sod Harvester Shown at Rutgers Sod and Research Field Days, June 20-22	24
Helicopter Tree Planting, Talks, Exhibits, Tours, Set for 42nd ISTC Convention in Cleveland This Month.....	30
Invert Emulsion Use in Aquatic Weed Control Reported at Hyacinth Control Meeting.....	34

Departments:

Editorial: Welcome to Cleveland.....	6
Know Your Species: Bitterweed.....	32
Meeting Dates	36
Classified Ads	44
Advertisers Index	44
Trimmings	46

Advertising Representatives

National Headquarters
1900 Euclid Avenue
Cleveland, Ohio 44115

Phone: Area Code 216+771-4169

JAMES A. NELSON
Editor and Publisher

MICHAEL I. LAH, JR.
Production Manager

D. BUNKIN
Circulation Supervisor

Chicago 60601
Peck & Billingslea, Inc.

185 North Wabash Ave., Suite 1809
Phone: 312+DEarborn 2-0292-93

New York City 10017
Billingslea & Ficke
420 Lexington Avenue
Phone: 212+532-1632

WEEDS TREES AND TURF is published monthly by Trade Magazines, Inc. Executive, editorial, and advertising offices: 1900 Euclid Ave., Cleveland, Ohio 44115.

Single Copy Price: 50 cents for current issue; all back issues 75 cents each. Foreign \$1.00.

Subscription Rates: U.S. and possessions, 1 year \$3.00; 2 years \$5.00. All other foreign subscriptions 1 year \$4.00; 2 years \$7.00. Change of Address: Three weeks advance notice is necessary for change of address. Both old and new address must be given. Post Office will not forward copies. Third Class postage is paid at Fostoria, Ohio.

Contents of this Issue © Trade Magazines, Inc., 1966

In a few weeks, over 1,000 treemen will be streaming into Cleveland for the 42nd Annual International Shade Tree Conference Convention and Trade Show. Since Cleveland is our hometown, *Weeds Trees and Turf* welcomes delegates to this active community of over one million which owes much of its beauty to the widespread use of trees that make its parks and suburban areas among the most elegant anywhere. This is truly the "Forest City."

In joint meeting with the ISTC will be the National Arborist Association. No two groups have done more to advance the science of tree care and to make Americans conscious of the need to save trees and to appreciate how much they add to our country's beauty.

Spearheading the ISTC is Dr. Lewis C. Chadwick, longtime executive secretary headquartered at Ohio State University in Columbus. Dr. Paul E. Tilford, until this past year, was the knowledgeable secretary of the National Arborist Association, who worked out of Wooster, Ohio, home of the Ohio Agricultural Research Station. His successor, Clarke W. Davis, now runs the NAA from Washington, D.C., and is already making a name for himself through his active public relations program.

The convention committee has planned a full

program that will benefit commercial arborists, municipal and utility treemen, and other vegetation controllers concerned with the protection and enhancement of trees.

This is a family affair, too. Local workers have arranged an active and fun-filled program for the ladies and the children. They will, among other things, board excursion boats to sightsee on Lake Erie which gently rolls against the northern shores of this city which is so delightful in August.

Treemen everywhere are invited to the ISTC convention at the Sheraton-Cleveland Hotel, August 28-September 2. Come. You'll be treated to a most valuable educational program, will see the newest in equipment and chemicals for tree work in the display booths, and will have a mighty fine time enjoying yourself, too. *Weeds Trees and Turf* looks forward to greeting you here at this important annual convention.

WEEDS TREES AND TURF is the national monthly magazine of urban/industrial vegetation maintenance, including turf management, weed and brush control, and tree care. Readers include "contract applicators," arborists, nurserymen, sod growers, and supervisory personnel with highway departments, railways, utilities, golf courses, and similar areas where vegetation must be enhanced or controlled. While the editors welcome contributions by qualified freelance writers, unsolicited manuscripts, unaccompanied by stamped, self-addressed envelopes, cannot be returned.



Myers

TT-29 TALL TREE MIST SPRAYER

with corrosion-proof GlasStran parts!

Developed specially for municipal maintenance operations . . . tree protection, mosquito control, leaf windrowing, brush burning, sanitation spraying, etc.

Requires only minimum care . . . air outlet case, engine cover, fuel tank-seat molded from high strength, 100% corrosion-proof GlasStran. Wide angle spray coverage . . . unit rotates 360°, outlet case elevates 15° below horizontal and 10° past vertical. Easy operation . . . all controls located at operator's console. Choice of skid-mounted or 2-wheel trailer models. Optional: power steering, seat belt, headlight accessory kit.

For complete information, see your Myers Sprayer Dealer now . . . or write today for details.

Myers . . . the finest name in power sprayers

THE F. E. MYERS & BRO. CO. SUBSIDIARY OF
400 ORANGE STREET • ASHLAND, OHIO 44805 McNEIL CORPORATION

High speed fan develops powerful air column that sends spray material to top of the tallest trees!

You say **BLUE CHIP**[®] means to fertilizer what diamonds meant to me?

That's right, Diamond Jim, just as diamonds were the trademark of your successful salesmanship, the Blue Chip tag on a fertilizer bag is the customer's assurance of a quality product that contains enough of the right kind of nitrogen.

Here's what the Blue Chip tag looks like. Be sure to ask your favorite supplier for a Blue Chip[®] fertilizer . . .



Nitroform[®] ureaform plant food contains 38% nitrogen and is one of the most efficient forms of nitrogen used today in fertilizer mixes. Released for plant use only by soil bacteria—active when growth conditions are right—Nitroform nitrogen is long-lasting and nonburning. Essentially water-insoluble, it will not leach out of the soil, and unused portions carry over as residual nitrogen until the next growing season. A synthetic organic fertilizer, Nitroform is odorless. And, Nitroform requires fewer applications for equivalent nitrogen, and less storage and handling than conventional fertilizers.

Nitroform is recommended for use on turfgrass, ornamentals, trees, and all plants that require sustained nitrogen feeding. It is easy to use and comes in two forms. Granular Blue Chip is applied by mechanical spreaders and is used in the fertilizer mixes that carry the Blue Chip tag. Powder Blue* is the first sprayable ureaform and is used on closely knit turf.



HERCULES TURF AND HORTICULTURAL PRODUCTS

FOR FURTHER INFORMATION WRITE: AGRICULTURAL CHEMICALS, SYNTHETICS DEPT., HERCULES INCORPORATED, WILMINGTON, DEL. 19899, OR CONTACT THESE SALES OFFICES: BOSTON, MASSACHUSETTS • BROWNSVILLE, TEXAS • CHICAGO (OAK BROOK), ILLINOIS • DALLAS, TEXAS • FRESNO, CALIFORNIA • GREENVILLE, MISSISSIPPI • LOUISIANA, MISSOURI • MONTGOMERY, ALABAMA • ORLANDO, FLORIDA • PHOENIX, ARIZONA • RALEIGH, NORTH CAROLINA • SAN FRANCISCO, CALIFORNIA • VANCOUVER, WASHINGTON.

® HERCULES TRADEMARK

STH66-5



By **DAN NEELY**
Plant Pathologist
Illinois Natural History Survey
Urbana, Illinois

Anthracnose of Sycamore and how to

An American sycamore with severe anthracnose symptoms. Most of the leaves were killed shortly after leaf emergence in the spring.



ONE OF the more unsightly nonlethal shade tree diseases is sycamore anthracnose. This disease is caused by a fungus that attacks leaves, twigs, and occasionally trunks of sycamore. It is most evident in May and June following opening of buds and emergence of new shoots. Young shoots are killed when the leaves are $\frac{1}{2}$ inch to 2 inches wide. An attacked tree has the general appearance of severe frost injury. Although the entire crown may be defoliated, often the lower half of the crown is affected more severely than the upper half. New leaves will appear during the summer but repeated attacks reduce growth and vigor of the tree. Occasionally trees become deformed and unsightly.

Sycamore anthracnose has been reported from almost every state in the United States in which American sycamore is a native tree. The states most frequently reporting severe occurrence are Missouri, Illinois, Indiana, Ohio, Pennsylvania, Maryland, Delaware, New Jersey, New York, Connecticut, and Massachusetts. Apparently the conditions favorable for disease development occur more frequently in these states.

Sycamore anthracnose is extremely severe in some years and almost negligible in others.



Shoot blight. Many of the young shoots on this branch were killed shortly after emerging from the buds. A few shoots were not killed and on these the leaves are normal in size.

Protect against it

It has long been observed that the disease occurs during "cold, wet springs." Recently it has been determined that spring temperature, not rainfall, is the dominant factor associated with disease severity. The temperature during the 2-week period immediately following first sycamore leaf emergence in the spring is of crucial importance. If the daily mean temperature for this 2-week period averages below 55 degrees, sycamore anthracnose is likely to be severe. As this temperature increases from 55 to 60 degrees sycamore anthracnose tends to be less severe. If the temperature for this period is above 60 degrees, few or no anthracnose symptoms will appear.

Most Susceptible Species

The species of *Platanus* subject to this disease are *P. occidentalis*, American sycamore (also called American plane); *P. acerifolia*, London plane; *P. orientalis*, Oriental plane; *P. racemosa*, California sycamore. The disease is more severe and occurs more often on *P. occidentalis* than on the other species named. However, there appears to be a variation in susceptibility within the species *P. occidentalis*. Because of severity of sycamore anthracnose on *P. occidentalis* in England, Holland, Bel-

gium, Northern France, and Germany, the common *Platanus* species grown in these countries is *P. acerifolia*, the London plane.

The fungus that causes sycamore anthracnose is named *Gnomonia platani*. Earlier the fungus was mistakenly named *Gnomonia veneta*. Unfortunately, most European and American writers have continued to call the sycamore anthracnose fungus *Gnomonia veneta*.

The fungus causing anthracnose of oak often has been

identified as the same fungus causing anthracnose of sycamore. Considering morphological, cultural, and pathogenic differences, the author feels that the fungus causing oak anthracnose is distinct and separate from the fungus causing sycamore anthracnose. The oak anthracnose fungus is *Gnomonia quercina*.

Four Developmental Stages

There are four developmental stages of sycamore anthracnose: bud, twig, shoot, and leaf blight. One or more of these may be observed during one season. The leaf blight stage probably originates following penetration of the leaf by fungus inoculum from an external source. During the summer and fall the fungus grows down the leaf veins, through the petiole, and infects the current season twig. The bud, twig, and shoot blight stages develop the following spring. They are the result of girdling action and canker formation in the infected twigs. The fungus in the canker usually remains active the second year

Bud blight. The buds failed to open because the twig tissue surrounding the buds was killed by fungus action.



Twig blight. The terminal portions of these twigs died after the twigs were girdled by the fungus.



following infection, and the canker increases in size. The repeated killing of twigs and accompanying canker formation throughout the crown causes abnormal branching and gnarled growths on many trees. Occasionally small trees are killed.

The four stages of anthracnose are described as follows:

Bud blight. The fungus present in the twig girdles and kills the tissue surrounding an individual bud before the bud expands or opens in the spring.

Twig blight. The fungus present in the twig girdles and kills the distal (tip) portion of the twig prior to bud expansion in the spring.

Shoot blight. This is similar to bud and twig blight but occurs later in the season. The fungus present in the twig girdles and kills tissue surrounding shoots that emerge from individual buds, or girdles and kills the entire twig containing many emerging shoots. Shoot blight occurs when the leaves on the emerging shoots are from 1/2 to 2 inches wide. It is often confused with frost injury.

Leaf blight. This stage occurs still later in the growing season. The first symptoms on the leaf blade are small faded, chlorotic spots. Spots that occur alongside or near veins increase in size and become necrotic (dried and shriveled). Necrotic areas along the midrib or main veins increase rapidly in size. When a major portion of the leaf blade or the petiole becomes diseased the leaf falls.

How to Protect Sycamores

Sycamores can be protected from anthracnose by application of an organic mercury fungicide. In areas where anthracnose occurs frequently, and where the American sycamore is highly valuable, the fungicide should be applied annually. In areas where sycamore is less valuable, the fungicide should be applied the year following disease occurrence to prevent excessive weakening of affected trees due to attacks during two consecutive years. Spraying of London plane is usually not warranted.

In most instances one appli-



Leaf blight. Necrotic areas along the leaf veins are characteristic leaf blight symptoms of sycamore anthracnose.



A 2-year-old canker on a twig of American sycamore.

cation of the fungicide, properly timed, is adequate. Application of the fungicide following appearance of disease symptoms is of little or no benefit. The spray must be applied in the spring when the sycamore buds are swelling and the bud caps are breaking.

Suggested organic mercury fungicides and rates of usage per 100 gallons of spray are as follows:

Coromerc, 1 1/2 pounds; Puratized Agricultural Spray, 1 1/2 pints; or Phix Apple Spray, 1/2 pound. In those unusual years when the weather remains cold for an extended period following application of the fungicide, a second application is recommended after 14 days and at two-thirds the previously suggested rate.

Why the organic mercury fungicides are effective in controlling anthracnose is not fully understood. Apparently they are able to penetrate the bark or wood of twigs in sufficient con-

centration to reduce fungus growth and prevent girdling of the buds or twigs.

The sycamore is a fast-growing tree that has been used extensively in recent years as a shade tree for home plantings. In areas where anthracnose is severe, sycamore will need special care, and planting other tree species is recommended.

Systemics Promising For Shade Trees

Systemic insecticides, which are absorbed by trees and translocated to stems and leaves, have definite advantages over spraying, according to Dr. Carlton S. Koehler, associate entomologist at the University of California, Berkeley. However, he does not expect systemics to replace conventional sprays.

Cautioning that some trees cannot be treated with systemics, the California entomologist adds that an overdose of systemic insecticide can do more harm to a tree than overspraying. He lists the advantages of systemics as their long residual action, economy, freedom from destruction of natural enemies of tree pests, simplicity of equipment required for treatment, and the ability to treat trees with systemics in unsatisfactory spraying weather.

Dr. Koehler also reports that in tests with Bidrin implantations at 6" intervals around elm trunks, aphids were controlled for 50 days and elm leaf beetles for an entire season.

T-H Improves Tedion

A new wettable powder formulation of its Tedion miticide has been announced by Thompson-Hayward. Containing 50% actual Tedion concentrate, the new W-50 formulation is said to be more economical and easier to handle than products previously available. Also marketed in an emulsifiable concentrate, W-50 is formulated for killing mite eggs, larvae, and nymphs.

Additional information on the new formulation and its uses can be obtained from Thompson-Hayward Chemical Co., Kansas City, Kans. 66110.