


Today, more than ever, Ditch Witch makes \$ense.

Dig Big!



NEW R100



4-wheel steer and tilt frame are standard.



Dig to depths of 8 feet, widths to 24 inches.

The R100 is Ditch Witch's new top-of-the-line Modularmatic! It's a big, powerful 100-horsepower model of underground construction equipment that combines proven Ditch Witch design concepts with new features that add up to the most versatile and productive machine in its class.

Exclusive Ditch Witch Modularmatic design has been utilized in the construction of the R100 vehicle; one basic vehicle can perform varying jobs simply by switching interchangeable modules. The trenching module is available with three digging booms to produce trench depths of 4- 6- or 8 feet. The 150,000 pound test digging chain can be set up with adapters and teeth to produce trench widths from 7- to 24-inches wide.

The R100 vehicle is built on a rigid, one-piece frame. It has full hydrostatic drive with 3-speed selectivity and a digging chain drive with four forward speeds and reverse. Rubber mounts isolate the operator's module from the vehicle's frame to provide operator comfort. The frame tilts to either side to provide vertical trench, even on a slope — an exclusive feature in 4-wheel-drive rubber tire mounted vehicles. Three industrial diesel engines, all in the 100-horsepower range, are offered.

Let Ditch Witch tell you more about our new top-line Modularmatic!

Ditch Witch . . . equipment from 7 - to 195-HP.


CHARLES MACHINE WORKS, INC.
P.O. Box 66
Perry, Oklahoma 73077



For More Details Circle (126) on Reply Card




TURF & INDUSTRIAL SPRAYERS




2000 Series Commercial Sprayer

A new compact design with 50 or 100 gal. rectangular polyethylene tank to save space, resist rust. Choose from high or low pressure systems for orchard, turf or pest control.



1950 Series All Purpose Sprayer

The versatile favorite offered in 100, 150 and 200 gal. tank sizes and four job-rated pump systems; features smooth mechanical agitation plus a full line of trailers and accessories.



960 Series Professional Sprayer

High performance equipment to deliver up to 25 GPM at 600 PSI. 200, 300 or 400 gal. capacity with dual stage centrifugal or piston pump systems, trailers and other equipment variations.

THE BROYHILL COMPANY
DAKOTA CITY, NEBRASKA 68731

For More Details Circle (110) on Reply Card

thiophanates, thus helping to explain its greater activity as a curative agent. Compared to the formulated parent fungicides, substantially lower amounts of the hydrolysis products MBC and EBC penetrate the foliage and move systemically in the plant. The penetration of thiabendazole is comparable to that of MBC and EBC. Although the fungitoxicity of methyl thiophanate is lower than for benomyl, it is greater than that of ethyl thiophanate because the methyl form has a slightly more rapid hydrolysis rate than the ethyl form, and because the fungitoxicity of MBC exceeds that of EBC. Thiabendazole is the least fungitoxic of the systemic fungicides discussed here.

When benomyl or thiophanate molecules are absorbed by roots, some of the chemical is stored in the root and released gradually as MBC and EBC. This provides a slow release of the fungitoxic factor, and thus long-term protection of the plant. If hydrolysis occurs outside the root, the MBC or EBC released by the soil is absorbed very rapidly and quickly passes through the plant, thus affording a shorter interval of protection. Since hydrolysis occurs in opened bags stored under humid conditions, one could expect that when these materials are used, they will be less efficient than those stored in unopened bags. This may be particularly important for granular formulations of systemic fungicides.

Application vs Efficiency

Systemic fungicides may be used as foliar sprays or as soil drenches. With the foliar application, the duration for activity of systemic fungicides is no longer than that of the contact fungicides, but because they enter the leaf and kill the fungi inside, their efficiency is somewhat greater than for the contacts. With both types of fungicides, the fungitoxicity is limited to the portions of the leaf that were actually treated. Foliar application of systemic fungicides offers no protection to roots and to newly emerging leaf tissue.

Efficient utilization of the full fungitoxic potential of systemic fungicides is achieved only through precisely controlled drenching applications. An understanding of the complexities involved in properly applying these fungicides to soil

(continued)

He has a new view of life



He saw for himself how modern coronary care units, more effective drugs and new methods of rehabilitation helped him beat the big one — heart attack.

Research scientists, many helped by your Heart Fund dollars, made these advances possible.

For greater advances in treatment and prevention . . .

Help your
Heart...

Help your
Heart Fund 
American Heart Association

Contributed by the Publisher

WEEDS TREES and TURF

get green carpet service for turf irrigation

Take advantage of the Johns-Manville "Total Support" system to spend water

like money.™ "Total Support" means people. The people who have given J-M the finest reputation for technical assistance and service can really go to work for you. These people are J-M and its nationwide distributor network who can work best with you to provide the system that's just right for the job.

"Total Support" also means you now can meet all your irrigation needs with reliable Johns-Manville products.

That means the best quality in components. Like **Buckner**® sprinklers, **Febco**® backflow preventors, **Buckner** and **Binar**® controls and **Johns-Manville** pipe. Together they give you accurate control, ease of maintenance, and efficient use of water. It all adds up to a better, more beautiful environment.

When you're in charge of the maintenance of a lush environment, make


sure you have the Johns-Manville "Total Support" system behind you. Fill out the coupon and mail it today to Johns-Manville, Box 5705-AT, Denver, Colo. 80217. We'll get behind you with the kind of quality, experience and technology you need.

spend water like money.

**J-M irrigation systems
save water, time and money**



Johns-Manville



Mail to: Johns-Manville
Box 5705-AT
Denver, Colorado 80217

I'd like to know more about Johns-Manville irrigation systems.

Name _____

Representing _____

Address _____

City _____ State _____ Zip _____

is perhaps the most important concept needed for their efficient use on a day to day basis. These fungicides adhere strongly to organic matter and clay minerals in soil. If they are allowed to dry even briefly on the leaf blade, in the thatch, or at the surface, a large amount of the efficiency will be lost. Delayed watering fails to move the chemicals from the sites where they dried to the root zone. Root density is much greater in soil below the thatch layer than in the thatch. A further complication is that the surface soil layer and the thatch are subject to periodic drying and moisture is required for the roots to absorb the fungicide. In drenching applications, it is important to know that improper drenching can leave the chemical where it is unavailable to the roots.

Special precautions are needed to prevent improper drenching applications on hot, dry days when the fungicide can dry onto the leaf blades or thatch within several minutes. It is helpful to wet the grass thoroughly before starting the application, and again during the job if the areas that were treated first are tending to dry. Drenching with one-half to one-inch of water should begin immediately after the application is completed. If a chemical residue is visible on the leaf blades when they dry after drenching, much of the efficiency has been lost. The practical difficulties with keeping the grass wet until drenching has been completed can be minimized by applying systemic fungicides very early in the morning while dew is on the leaf blades, or during a steady light rainfall.

Optimal efficiency depends upon uniformly distributing the fungicide in the soil. To aid in uniform distribution, it is advisable to minimize the penetration problems by removing excess thatch and wetting the soil thoroughly about three days before treatment. Water applied to dry soil first flows through the largest pores and passages, and then slowly wets the smaller pores. Drenching the fungicide into a dry soil may cause the chemical to be nonuniformly deposited primarily along the larger pores and passages of soil. This reduces the availability of the fungicide to many roots.

Systemic fungicides adhere more tightly to acid soils than to alkaline soils. Excess soil acidity should therefore reduce the actual amount of these fungicides that can be ab-

sorbed by roots immediately after application. Excess soil acidity also increases the rate of hydrolysis, thus further decreasing the efficiency of fungicide utilization. This is especially important for control of root-infecting fungi which require high initial concentrations of the fungicide.

The optimal time to apply systemic fungicides is just before the rapid phase of epidemic build-up. Thus, in areas where *Fusarium* blight generally becomes visually apparent by early July, the optimal application time is in mid-to-late-June when early infection is occurring. As applications are delayed beyond mid-June, the fungicide will be less efficient. The fungus which causes stripe smut lives inside living grass plants during the winter. New infections and symptom development occurs in the Spring. Optimal timing of treatments is in the late-Fall to eradicate the overwintering fungus and in the early Spring before the grass begins to green up to prevent new infections. For foliar diseases, the optimal timing is immediately after they can be seen.

Fate of Systemic Fungicides

Systemic fungicides are only translocated upward from the point of absorption. Foliar applications therefore protect against foliar-infecting fungi, but not against the root infecting fungi. The fungicides migrate toward the upper edges of leaves, and a little may even be exuded from the leaf. The fungicide that is exuded could conceivably be recycled through the plant again if it gets into the root zone before being absorbed on the thatch or soil surface. The fungicide remaining in the leaf clippings probably becomes ineffective for future protection of living plants. Some fungicide could be leached below the root zone in very sandy soils that are devoid of organic matter, but this would not be a problem in most turfgrass soils. The fungicide which is absorbed to soil degrades very slowly. About one-half of it is degraded within the first six months. This means that repeated heavy applications may accumulate to reach phytotoxic concentrations.

Fungal Resistance to Systemic Fungicides


We can expect fungi to become resistant to rather specific fungicides such as the systemics because they inhibit only one event in the

metabolism of the fungus. A simple mutation of one gene can lead to a strain of the fungus that is resistant, and selection of the resistant strain in turf could give rise to an overall loss of efficiency of the systemics in the area where resistance appeared. Since the mode of action of all current systemics is very similar, resistance to one chemical will also likely be expressed by resistance to the others. Systemics with different modes of action will be developed in the future and will alleviate this problem of cross resistance among the systemic fungicides. Contact fungicides are active against strains resistant to systemics.

Several precautions can be used to reduce the selection pressure for resistant strains. If you must use systemics, as is the case with root-infecting fungi which cause *Fusarium* blight and stripe smut, reduce your use of systemic fungicides against other diseases which can be controlled with contact materials. If high rates of systemics have not been applied for control of the root-infecting fungi, it is acceptable to use them at recommended rates against foliar diseases. When used in this manner, they should not be used repeatedly, but alternately with at least two contact materials. To reduce the selection pressure for resistant strains, it is also necessary to avoid prolonged contact of the fungus with low concentrations of the systemic fungicides. This situation exists when a systemic is repeatedly applied at low rates, either alone or in a fertilizer-fungicide mixture, for control of a disease such as dollar spot. It would be preferable to use a single curative application of a systemic soon after dollar spot appears, and then use a contact material in most of the follow-up applications.

There are no confirmed instances of "resistant-*Fusarium*" on turf in New York State. Where control is less than hoped for, a reevaluation of recommended fungicide application procedures or rates, and of cultural management practices should be considered (see Nassau County Coop. Extension mimeo C-2-7, "*Fusarium* blight of turfgrass"). The only turf pathogen which is known with certainty to have become resistant to systemic fungicides in our state is *Sclerotinia homeocarpa*. The resistant dollar spot fungus currently appears only

(continued)



**One quick
trip down and
back is all it
takes.
With Myers.**

This could be the best thing that ever happened to your turf maintenance program. With a Myers TL32TM golf course sprayer, you can cover the whole width of a fairway in just two passes. Using a two-side discharge, you can drive down the fairway and cover 70' at a time. Or, if you prefer "driving in the rough" and using a one-side discharge you can cover 40' to 50' each trip. The sprayer's powerful 32" fan sends out the fungicide spray in a ground-hugging pattern that really does the job. It keeps your fairways in tip-top playing shape and does it in a lot less time than you'd

expect! On top of that, the sprayer can be converted with the P70 attachment for tall shade tree applications to 80 ft.! The TL32 really cuts those big fairway jobs and tall trees down to size. Interested in finding out more? Ask about the TL32TM today at your Myers TurfLine dealer's.

Myers®

The name that works for you.



Myers TL32TM
with shade tree attachment

in isolated lawns in Nassau County.

Other Considerations

Systemic fungicides are toxic to earthworms and nematodes in soil. Earthworms are considered beneficial on most turfgrass soils, except under golf course putting greens. Reductions in numbers of harmful nematodes is considered to be a desirable side effect. Repeated foliar applications of systemic fungicides also reduces the populations of harmful mites in turf. When their populations decrease, so do the populations of predator mites which feed upon the harmful types. At a later time, the populations of predator mites can't increase until their food source (the harmful types) is supplied. In the meanwhile, the harmful mites can cause extensive damage to plants. This is possibly a factor in the observed decline of Kentucky bluegrass turf quality in the spring following repeated foliar applications of systemic fungicides during the previous summer. This tendency is thought to be much less serious or nonexistent with drench applications.

Summary

The systemic fungicides are a remarkable class of pesticides. Their attributes are many. But, as with every pesticide, certain restraints or precautions must be exercised in their use. Their efficiency is greatly reduced if they are improperly applied to soil. Since they are more selective than the older contact fungicides, more attention must be given to the diseases which are not controlled by the systemic fungicides. Repeated heavy applications can lead to phytotoxicity problems, or to an overall decline in turf quality. Pesticide programs which rely heavily upon this one class of fungicides can be expected to select strains of fungi which are resistant. In these cases, the only recourse is to shift back to contact fungicides, to resistant turf varieties, and to good cultural controls. Based upon these considerations, it seems wise to exercise caution in the use of systemics in the pesticide programs. If root-infecting fungi are not predominant in your turf, use mixed fungicidal programs for all diseases. Where high rates of systemics are used to control root-infecting fungi, it would appear desirable to avoid use of these fungicides for other diseases which are controlled very well by the contact fungicides.

32

Smith Turf Changes Name

Smith Turf Irrigation, Inc., of Milford, Conn., also doing business as Irrigation and Equipment Supply, has changed its name to I & E Supply, Inc., according to President Richard W. Smith.

Smith said the name change and a move to larger quarters will facilitate the firm's move into the role of a more complete irrigation and industrial supply house, specializing in sprinkler irrigation equipment, industrial pipe, valves, fittings, industrial pumps and pump packages. The move represents a change from its former role of strictly an irrigation supply house, Smith said.

Calamco and Occidental To Combine Headquarters

October ground-breaking ceremonies at the Occidental Chemical Company (Oxychem) plantsite, Lathrop, Calif., officially began construction of the combined office headquarters for California Ammonia Co. (Calamco) and Oxychem.

According to C. Martin Wilmarth, Calamco president, and James H. Lindley, vice president of Oxychem, the ever-increasing demand for nitrogen products and ammonium phosphate fertilizers has necessitated a large expansion pro-

ject involving new plants, new people and additional office facilities.

The building, to be built in three stages, is planned to accommodate future expansion as well as present employee requirements. The first stage will consist of 12,400 sq. ft. and will house over 60 people in finance, credit, accounting and data processing. When complete, the second stage will expand the ground level to 16,000 sq. ft., to which a second story will be added bringing the total size to 32,000 sq. ft. over the next decade.

Central Plains Turfgrass Elects Officers for 1975

Officers who will head the Central Plains Turfgrass Foundation for the coming year are: Larry Runyon, Kansas City, Mo., re-elected president; Monty Brown, Wichita, Kan., elected vice president; and Dr. Ray A. Keen, Kansas State University (KSU), Manhattan, Kan., reappointed secretary-treasurer. New directors include Richard Gray, Wichita, Kan., and Herman Siler, Springfield, Mo. Dr. Ron Campbell of KSU was re-elected as a director. The officers were named during the Foundation's annual business meeting, held in conjunction with the annual KSU turfgrass conference at Manhattan, Kan., in October.



Directors of the Southern Turfgrass Association are planning for the 1975 Southern Turfgrass Conference and Show, March 2-4 at the Cook Center in Memphis, Tenn. Seated (left to right): Gene Baston, CGCS, vice president; Marion Johnson, CGCS, president; Reg Perry, executive secretary; Arlin Grant (Florida); Al Frenette (Georgia); Euel Coats (Mississippi). Standing (left to right): Jim Bridges (Tennessee); Billy Smith (Arkansas); Tommie Hill (Alabama); Kayo Mullen (Kentucky); Sam Locke (Texas); Carter Huff (Missouri).

The Exciting New **TURFTYPE** Fine Leaved Perennial Ryegrass

from
Holland!

GAME

The new
revolutionary
lawngrass



GAME VARIETY **TURF TYPE** **PERENNIAL RYEGRASS**

FINE LEAVED GRASS WITH
MANY NEW
OUTSTANDING FEATURES!

A NEW VARIETY of Perennial Ryegrass . . .
GAME, was developed in the Netherlands by
a leading grass seed breeder.

GAME was bred from a selection of
flat growing plants. All the plants col-
lected showing characteristics desirable
for lawn use were tested over a three
year period under a system of close
mowing. Only those plants which sur-
vived this test and showed the best
of other traits were used in breeding
GAME.

UNDER THE highest quality controlled
conditions available, GAME is now be-
ing grown in Oregon on a Certified
Basis.

GAME is extremely drought resistant,
and in comparison to other varieties con-
tinues to show its great quality under
dry conditions.

GAME has extremely high regrowth
characteristics. It readily survives close
mowing and hard wear.



For information call or write:

Willamette Seed & Grain Co.

Bob Lowry
503—926-8883
P. O. Box 791
Albany, Ore. 97321

Max Elder
503—491-3675
P. O. Box 25
Shedd, Ore. 97377

Milton Tuck
503—926-8883
P. O. Box 791
Albany, Ore. 97321

Bob Richardson
503—491-3675
P. O. Box 25
Shedd, Ore. 97377

How Lexington, Mass., solved its caterpillar problem with no adverse side effects.

Lexington is densely populated. So Paul Mazerall, park superintendent and tree warden, had a big job on his hands.

To stop the infestation of tent caterpillars, he had to spray around schools, parks, churches and other public places. His choice of insecticide became critical.

Paul found a product that met all his requirements. A product that wouldn't endanger local wildlife or pets or people. And one which city work crews could handle easily. It was DIPEL®.

One shot did the job.

That's all it took to get rid of the tent caterpillar problem in Lexington.

Selective control made DIPEL the answer to Lexington's problem. It worked so well, in fact, that Paul plans to expand its use.

Kills gypsy moth and others.

Including cankerworm (inchworm). Spanworm. Bagworm. Fall webworm. California oakworm (California only).

But performance is just one of the reasons cities like Lexington are switching to DIPEL.

DIPEL can be mixed and applied without protective clothing. And it mixes easily with foliar nutrients. So you can do more than one job during each pass. As a matter of fact, some cities are even spraying DIPEL from aircraft.

Yes, tree wardens like Paul Mazerall have the right idea about caterpillar control. When you have a problem, spray DIPEL.

DIPEL®



AGRICULTURAL AND VETERINARY
PRODUCTS DIVISION

ABBOTT LABORATORIES, NORTH CHICAGO, ILL. 60064

GENTLEMEN: We are interested in spraying DIPEL®

Please phone me promptly. Have your representative visit. Mail literature.

Name _____ Acres _____

Title _____ Pest(s) _____

Address _____


City, Zip _____ By ground rig

Phone No. _____ By air

Company, Twosp., District, Etc. _____

SEND TO: ABBOTT LABORATORIES, 900 West Route 70, Suite 6, Marlton, New Jersey 08053





QUALITY

Rolls Royce means quality. Quality performance, the standard by which others are judged. Another name synonymous with quality and performance is Jacklin. Jacklin, one of the world's largest producers of Kentucky bluegrasses and other varieties. When ordering seeds specify JACKLIN SEED.

JACKLIN PRODUCES:

Merion, 0217® Fylking, Glade, Cheri, Nugget, S-21, Newport, Delta, Geary, Park, Six, and Troy Kentucky bluegrass; Rathdrum and Reubens Canada bluegrass; Norlea Perennial ryegrass, Public and private varieties of Creeping Red, Nova Rubra, Chewings, and Hard fescues.

And Jacklin has produced on contract:

Pennfine, N.K. 100, N.K. 200 and Pelo Perennial ryegrass. Jacklin is also a prime* contract producer of: Adelphi, Warren's A-34 and O.M. Scott's Windsor Kentucky bluegrass.

Jacklin has built a reputation of fine quality and service for over 30 years. You can count on purity of content, high quality and sure germination.

*Available only through breeder and owner outlets.

Jacklin® Seed Company

Division of the Vaughan-Jacklin Corporation

For More Details Circle (109) on Reply Card

◆ For More Details On Preceding Page Circle (112) On Reply Card

people on the move

Robert Orvis, promoted to works manager for FMC Corp.'s Outdoor Power Equipment Division.

* * *

J. Patrick Kaine, elected vice president of International Harvester. Kaine will continue to serve as president of IH's Agricultural Equipment Division.

* * *

Ronald S. Leafblad, appointed vice president, Outdoor Power Equipment, The Toro Company.

* * *

James A. Hunter, appointed control manager of the International Department of Hercules Incorporated. Hunter succeeds **William C. Prescott** who is retiring in early 1975.

* * *

Robert W. Brandt, named Forest Service Director of International Forestry.

* * *

Herbert A. Jespersen, named vice president of Outboard Marine Corporation and a division manager of OMC-Lincoln, a division of Outboard Marine Corp.

* * *

Dennis R. Wright, appointed to newly-created position of administrative manager, Vicon Farm Machinery, Inc.

* * *

Duane Ferrell, joined Thompson-Hayward Chemical Co. as director of New Pesticide Registrations. Ferrell will be responsible for planning and evaluating research with candidate pesticides.

* * *

Gary M. Cook, appointed senior vice president of Agrico Chemical Co.

* * *

David A. Markgraf, named director of marketing and sales, and **Jeffrey W. Raymond**, appointed sales representative in Florida and the Southeast for Applied Biochemists, Inc.

* * *

Wayne Kincannon, named a vice president of Diamond Shamrock Chemical Co.

* * *

Thomas B. Moorhead, elected vice president, administration, for Beker Industries Corp., a producer of chemical fertilizer headquartered in Greenwich, Conn.

* * *

Larry D. Liptac, appointed director of Agricultural Business Group, Velsicol Chemical Corp.

* * *

Ronald D. Ross, Jr. and **Joseph J. Maher**, promoted to senior analytical chemists in Ciba-Geigy's Agricultural Division. **Michael F. Sandine**, joined the Division as a senior experimentalist at the Boynton Beach Research Farm in Florida.

* * *

Richard E. Peltier, joined Disston, Inc. as plant manager of Disston's Danville, Va., operation.

* * *

Pipe & Plastics Group of Certain-teed Products Corp. named two marketing managers: **John E. Calkins**, asbestos cement and PVC pressure pipe, and **Alvin I. Leff**, non-pressure pipe products.