

granulars to the surface and soaked into the soil. If you consider that some grubs may be under the surface by more than a foot, the diluted insecticide has a long way to go and stay toxic. For this reason, more preventative adult control measures are being stressed. The reduction of insecticide effectiveness by thatch may be restrictive to control in some circumstances.

Perhaps a lesson learned by an overreliance on highly residual pesticides is the development of resistant insects. Insects have tremendous reproductive capacity. If a pesticide lasts for ten or more years in the soil, the few insects that weren't effected by the pesticide become the genetic base for future populations. This fluke resistance quickly becomes a widespread tolerance or resistance to the chemical. Hence no control. Repeat applications prove equally ineffective. Alternative insecticides then become necessary.

Similar adaptation by turfgrasses affords some hope for natural controls in the future. If researchers can identify turfgrasses which insects avoid, they can perhaps lower the dependence on chemicals for control. Dr. Jack Murray and Dr. Roger Ratcliffe at the USDA Research Center in Beltsville, MD, have detected aphid resistance in some cultivars of bluegrasses, fescues, and ryegrasses. Northrup King has been doing similar studies at its Minnesota research farm. Reinert in Florida has been checking resistance of warm season grasses to mealy bug, bermudagrass mite, and the mole cricket.

On the other hand, the Hyperodes weevil has a specific liking for *Poa annua*. Dr. Harue Tashiro of the New York State Agricultural Experiment Station in Geneva, has studied this insect closely. Confined mainly to the Long Island area, the insect has the potential to spread since *Poa annua* is common, although usually unwanted, on many eastern courses.

#### Major Turf Pests and Their Control

There are a dozen serious turf insect and mite pests in the U.S. with the potential for more. Following is an outline of the insect, current control methods, and potential solutions of problems.

#### Aphids

The only aphid of significance to turfgrass managers is the greenbug, *Schizaphis graminum*. It is a serious pest of grain crops and forage grasses. The first damage to Kentucky bluegrass was noticed by Dr. Roscoe Randall of the University of Illinois in 1970. No damage to other turfgrasses has been found.

IF  
THERE'S  
A SHADY  
SPOT WHERE  
SABRE  
WON'T GROW,  
THEN NO  
OTHER TURF  
GRASS WILL

Sabre is the new domestic variety of *Poa trivialis* which is a legendary performer in the shade. In actual use and on test plots it has proven itself superior in the shade to the fine fescues. *And that's saying something.*

Sabre is also one of the prettiest turf grasses you've ever seen. Deep green, it is fine-bladed and its mowing qualities are truly superior.

As a shade tolerant variety, it should be sown only in shady areas. It may be mixed with the fine fescues such as Highlight, Ensylva or Pennlawn, or bluegrasses such as Baron, A-34, Glade, Merit or Nugget, or used in a mixture which includes 20% turf-type perennial ryegrass such as Derby, Yorktown or Pennfine, or it may be sown alone.

When mixed with other fine turf grasses in a winter overseeding mixture in the Southern U.S., it can dramatically cut seeding rates. The addition of 20% Sabre (by weight) will cut the seeding rate of a fine fescue or ryegrass mixture by 50%.

While its normal cutting height is 1½ inches, it will prosper at lesser heights including 3/16 on a putting green.

Free of *Poa annua*, Sabre is also a disease-resistant turf grass.



**INTERNATIONAL SEEDS, INC.**

P.O. Box 168, Halsey, Oregon 97348

Telephone: (503) 369-2251 • TWX 510/590-0765

Write 117 on reader service card

# CHIPCO



**LONG-LASTING  
CHIPCO® 26019 IS  
STRONG MEDICINE.** If you're serious about protecting your turf against diseases, give it your best shot. Spray Chipco 26019 fungicide. Thousands of superintendents have switched to Chipco 26019 in the past two years. And no wonder. Chipco 26019 gives you the longest residual. You can cut the number of sprays in your program, and that cuts your costs. And Chipco 26019 prevents the major turf diseases with unsurpassed effectiveness. You'll get good results with Chipco

# CHIPCO® 26019



26019 against dollar spot (including benomyl-resistant dollar spot), Helminthosporium (leaf spot and melting out), brown patch, Fusarium blight and Fusarium patch (west of the Cascade Mountains). Chipco 26019 also suppresses grey and pink snow molds. Ask your Chipco distributor or Rhône Poulenc representative about effective, long-lasting Chipco 26019...the strong medicine in turf disease prevention. Rhône Poulenc Chemical Co., Agrochemical Div., Rhône Poulenc Inc. Monmouth Junction, NJ 08852.



Write 131 on reader service card



Four stages of metamorphosis for the Black turfgrass ataeenius.

The greenbug, in the process of sucking out plant fluids, injects fluids to break down cell walls within the plant. This fluid kills living tissues in the plant, resulting in a yellow and finally

brown appearance to affected turf. The aphids commonly infect Kentucky bluegrass under the canopies of trees.

Agricultural experts on this pest previously thought the aphid could not

overwinter north of Kentucky. However, Ohio's Niemczyk and his "Angels" have discovered reinfestations in the same locations in consecutive years. Niemczyk suggests the reinfestation is more than a coincidence of migrating adults from the South selecting the exact same location every year.

Lawn care companies are reporting increased incidence of the pest in the Midwest. Organophosphates have limited effectiveness upon this pest which can produce 20 generations in a single season. Special local needs labels in Ohio for Orthene from Chevron and Pirimor from ICI have given improved control. Randell reported good control with malathion in 1978.

Murray and Ratcliffe at USDA, Beltsville, are studying turfgrasses resistant to the sucking damage of the greenbug aphid.


#### Ataeenius

The Black turfgrass ataeenius, *Ataeenius spretulus*, has appeared sporadically on turfgrass across the U.S. In 1932, larvae of the pest were discovered in damaged turf on a Minnesota golf course. Similar damage was found in New York in 1970. Since then the number and frequency of sitings have increased and entomologists are concerned.

The ataeenius larvae feed on the roots of both bentgrass and Kentucky bluegrass. This feeding takes place after the eggs hatch in late May to mid-July. Ohio entomologists have noticed a second generation of larvae in late August and September. At the end of this period the larvae enter a short pupal stage and emerge as adults. These adults can begin laying eggs in July. The adults tend to leave the fairways for tall grassy fringe areas to overwinter.

Symptoms may be a drought appearance in June and July despite adequate irrigation. The greenbug has not been identified to warm-season turfgrass damage. Since the larva does not burrow too deeply into the soils, controls work relatively well. Prior to the expansion of the label interpretations in 1978, turf managers had a pest with no labelled control. Now, an applicator may use a pesticide against any pest not on the label if the application is to a crop, animal, or site specified on the label, unless the EPA specifically requires otherwise.

Ohio entomologists are encouraging preventative control of the ataeenius adult before it has a chance to lay eggs. Timing becomes critical in preventative control and regional entomologists should be contacted for information on timing.



### The Pickseed Winners Circle

These winners were selected for their outstanding turf quality and durability. They will withstand the ravages of man and nature, maintaining a beautiful dense turf that will resist disease, overcome summer and winter stress, and help prevent weed invasions.

These winners will reduce your maintenance costs and provide you with a better looking turf. Go with the winners.

**PICKSEED**  
PICKSEED WEST, Inc.  
801 8th, TANGENT, OREGON 97136 (503) 878-8888

Write 130 on reader service card

# Because the job demands success...



Because the job demands success, The Andersons has devoted 10 years to developing and refining our Tee-Time professional turf-care products. We have now established a network of qualified distributors and can make available to you the quality products we've been supplying to a limited market for many years.

The Andersons is a diversified agri-business committed to product quality and service.

Our company has enjoyed continuous growth in each of our divisions, accompanied by a widespread reputation for reliability, applied research, and technical innovation. Today we are the nation's sixth largest manufacturer of turf-care fertilizer products.

You can switch to The Andersons' products with confidence. If you are a professional turf manager, you're going to be very pleased with

the success you can achieve with our diverse line. We supply more than 20 products which can be selectively combined according to your particular needs to produce maximum results.

Our distributors, backed by our extensive technical knowledge, are qualified to assist you in tailoring your programs. Ask if your present supplier carries The Andersons Tee-Time products. If not, call us toll-free or write and

we'll give you the name of your nearest distributor. You'll be glad you did. Distributor inquiries welcomed.

**the professional's partner**

**The Andersons**

Lawn Fertilizer Division  
P.O. Box 119  
Maumee, Ohio 43537  
Ohio—800-472-3220  
Outside Ohio—  
800-537-3370



**Adult hairy chinchbug.** This pest may produce three generations in a single season.

### Billbugs

There are three primary species of billbug affecting turf. The bluegrass billbug is the only cool-season billbug pest. The transition zone and warm-season zones are bothered by the zoysia billbug and the Phoenix billbug which attacks bermudagrass. A possible fourth species has been discovered in Denver, Colorado.

In May and June, adults begin laying eggs near the crown on the stems of the turfgrass plant. After hatching, the billbug larvae feed on the stems, moving downward to the crown and eventually to the roots. Larvae are most abundant in July and August. Damage by billbugs can be confirmed by examining the crown of dying plants. Excessive damage in this area and the presence of a white, sawdust-like material in the root zone are good clues.

Control centers around adults in April to May and larvae from June to August. Entomologists lean toward adult control. Since the larvae move downward from the stem, to the crown, into the thatch and finally into the soil, control must change with their location. For example, soaking insecticides into the soil may leach them out of the thatch where the larvae may be at the time.

### Chinchbugs

Like billbugs, there are cool and warm-season chinchbugs. The hairy chinchbug attacks bluegrasses, fine fescues, and bentgrasses. The southern chinchbug feeds on bluegrasses, zoysiagrass, and especially St. Augustine-

grass.

Like the greenbug, chinchbugs suck out plant juices and at the same time inject digestive juices into the plant which cause death of plant tissues. Nymphs inflict the same type of damage to the turf as the adult. The hairy chinchbug may produce three generations in one year whereas the southern chinchbug may produce seven or more.

Chinchbugs take special advantage of patches of turf under summer drought stress. Eggs are laid in the lower leaf sheaths of grasses. Southern chinchbug adults are active year-round, whereas chinchbug adults are active from late March to late September.

Chinchbugs are highly susceptible to a fungus disease called *Beauveria*, especially when the turf is adequately irrigated. Proper fertilization can help the grasses resist a second or third generation.

This highly reproductive insect has exhibited some resistance to many insecticides. When resistance has been a factor, Baygon has served as a good replacement. Early treatment of the population to reduce the size of future generations is good policy.

Murray and Ratcliffe at USDA, Beltsville, have studied the resistance of Kentucky bluegrass and fescues to chinchbugs.

### Cutworms and Armyworms

These are the larvae of moths. The adult moths are active primarily at night. Armyworms are a serious south-

ern problem with some northern damage. The bronzed cutworm has been a problem in the Maryland area. The black cutworm is a northern problem.

Cutworm larvae become active in April, feeding during the night and retreating into the thatch and soil during the day. There is one generation per year. Materials applied to the plant surface during the day must remain effective to kill the chewing pests during the night. Solutions of pyrethrins or detergents can excite the larvae to the surface.

The armyworm is very similar in habit to the cutworm except that it is known to migrate during the night and can produce three generations in a single year. Control measures are the same as for cutworms. Consult regional entomologists for application timing. Early generation control is most efficient. The fall armyworm, unlike the true armyworm, can migrate northward to inflict damage.

### Grubs

Grubs are the larvae of many different beetles, including the Japanese beetle, the atenioid, May Beetle, European and masked chafers, and garden beetles. Not only do grubs damage turfgrass by feeding on roots, but they attract damage from moles, birds, and skunks feeding on them.

To identify the grub, the turf manager must examine the bottom side of the last body segment, the raster. The pattern of hairs on the raster determines which beetle the grub will be. This identification of the type of grub is needed to determine the timing of control measures and in the case of Milky Spore Disease, susceptible target Japanese beetles.


Grub damage is most evident in the spring and fall. Patches of dead or dying turf appear. Presence of moles or flocks of blackbirds provide a clue to grub infestation.

Diazinon, Dylox and Proxol are the chief controls. They remain active in the soil for only two weeks so timing is critical. pH is a limiting factor with Dylox and Proxol, since high pH will cause premature breakdown. Diazinon is attracted to organic matter in the soil, sometimes binding to thatch and other material reaching the pest. Uniform distribution and penetration to the pest's location are very important. Post-treatment irrigation is almost essential to good distribution.

Milky Spore Disease is effective only against the Japanese beetle. Use on other grubs will be useless. Overreliance on the chlorinated cyclodiene insecticides encouraged development of

# V250

...a dependable lawn plow.



When job conditions call for plowing in services, consider the Ditch Witch V250. It's the most compact lawn plow available, and it has design features like a newly designed ground drive and plow-lift systems that make it a dependable, efficient alternative to trenching. Put in power and communication cable, plastic pipe for water or natural gas services, a complete underground sprinkler system. Let us give you all the details on the Ditch Witch V250 vibratory plow.



## I need an alternative to trenching!

Please send information on:

WT&T

- the V250     other vibratory plow units
- CATV installations     Sprinkler System Installations
- Ditch Witch Full line brochure

Name: \_\_\_\_\_

Company: \_\_\_\_\_

Address: \_\_\_\_\_ Phone: \_\_\_\_\_

City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_

Mail to:

**The Charles Machine Works, Inc.**  
P.O. Box 66 • Perry, Oklahoma 73077

Write 111 on reader service card

# Ditch Witch®

Get the right  
pressure at the  
nozzle...



# Go Hypro all the way!



Big jobs or small jobs you need steady, reliable pressure at the nozzle. You get it with Hypro sprayer pumps and accessories: pressure for long discharge lines, for compact portable units, for fine spray or steady stream. That means time savings and application precision that can help you make more money. And Hypro dependability keeps your sprayer on the job, season after season. Get the facts. See your Hypro distributor or write us today.

### Hypro High Speed Centrifugal Pumps.

Belt driven. Models available for hydraulic, gear or engine drive, too. Capacities to 130 gpm. Pressures to 170 psi. Speeds to 6000 rpm.



### Hypro Piston Pumps.

The Series 5900. 20 gpm at 500 psi—plenty of output for proper agitation. Other Hypro piston pumps with capacities to 25 gpm.



Hypro offers a complete selection of sprayer accessories: gauges, strainers, agitators, ball valves and couplings, and a full line of valves and fittings.

Send for your free  
Hypro catalog now.

 **Hypro**  
A DIVISION OF LEAR SIEGLER, INC.

347 Fifth Ave NW, St. Paul, MN 55112 • (612) 633-9300

resistant grubs. Organophosphates are the main replacement. MoBay is currently working toward a label for isofenphos which shows promise for grub control.

Control of grubs is still one of the most difficult problems to conquer. Control of 75 to 80 percent is as high as can be expected.

### Ground Pearls

Ground pearls are a warm-season turfgrass pest. They feed on the roots of centipedegrass and other warm-season grasses. The tiny little yellow-white ball-shaped nymphs attach themselves to the roots with their piercing-sucking mouthparts, become adults and remain attached for two to three years. They secrete a substance which creates a hard shell around them. Turf attacked by the Ground Pearl turns yellowish-brown in irregular patterns.

Control is very difficult. This is another reason for study of effects of thatch on insecticides drenched into the soil.

### Hyperodes weevil

The *Hyperodes* weevil is a chewing insect that has a taste for *Poa annua*. Turf areas dependent upon *Poa annua* in the Northeast must control this pest or renovate to other turfgrass types. Adults emerge from overwintering locations in the spring and do minor damage to the turf. The major problem is the way the adult inserts eggs between leaf sheaths. When the insect hatches, the larva feeds within the stem and works its way downward to the crown to inflict serious damage. The larvae enters the soil to pupate and emerges eight days later as an adult.

Damage becomes obvious in late May and early June. Organophosphates applied at the time provide good control. The target area for the larva is the thatch, so soaking is not recommended.

Dr. Harue Tashiro at Cornell is the national expert on this particular pest.

### Mealy Bugs

Related to the Ground Pearl, the rhodesgrass mealy bug is a sucking insect which locates in masses at the crown or at the nodes of the grass stems. It secretes a shell for protection like the Ground Pearl. Reinert of Florida has studied the bermudagrass mealy bug and has trials for resistant turfgrass varieties.

### Mites

There are two mites of importance to turfgrass. The winter grain mite feeds

on plant juices just under the leaf surface in the late fall and winter when temperature permits. They have rasping-sucking mouthparts. Evidence of infestation is limited and sporadic, but control research may be necessary. The winter grain mite lays bright orange eggs in the thatch and soil beginning in March. The eggs do not hatch until fall when they begin to damage bluegrass and fescue.

The bermudagrass mite is a serious pest in the South where they furiously suck out the juices of bermudagrass plants. Stems die individually as they are damaged. Liquid applications tend to be more effective than granular insecticide formulations for control of mites. Organophosphates do well. A serious threat by mites is to grasses grown for seed in the Northwest.

### Mole Crickets

Although mole crickets occur as far north as Ohio, they are basically a southeastern problem. All stages feed on the roots and can weaken turf by disturbing the root zone.

Adults burrow into the soil to deposit eggs in the spring. Two weeks later nymphs emerge to feed on roots. Root-zone disturbance may be a bigger problem than root feeding says Reinert of Florida. The disturbance can uproot plants and create ridges, like miniature mole tunnels. The crickets come to the surface at night. Reinert reports good control with dursban, Baygon, Scotts Nematicide/Insecticide (ethoprop) and malathion.

### Webworms

There are more than 30 species of webworms. The tropical sod webworm creates a larger problem due to a greater number of generations per year. The adults are small, grayish-white moths which fly over the turf at dusk. Female moths drop eggs into the turf as they fly. Eggs hatch in less than two weeks as caterpillars. These larvae construct tunnels through the thatch which they line with silken webs. They feed on the grass blades nearest to their tunnels. The larva is the destructive stage, not the adult. Larvae can be irritated with pyrethrins and brought to the surface for examination.

Insecticides applied to leaf surfaces work well. One webworm, the cranberry girdler, has been found in Illinois and Michigan. It feeds in the crowns and roots and must be treated as a soil insect.

Write 116 on reader service card



# FALCON

PVP applied for

**Tall**

**Fescue**



## **New turf-type tall fescue of the 80's**

You'll like this new, fine-textured variety with higher density than most of the currently available tall fescues. Falcon is a leafy, moderately low growing turf-type variety with pleasing green color and tolerance to heat and drought. It's a good performer under minimum maintenance programs, too. Throughout its area of adaptation, mature stands of Falcon have shown tolerance to wear and moderate shade which makes it useful for home lawns and play areas such as athletic fields and parks.

**Contact us for more information.**



**E. F. BURLINGHAM & SONS**

P.O. Box 217 • Forest Grove, OR 97116

(503) 357-2141 • Telex: 36-0274

Cable: Burlingham

# Green, Green, Greener

**THE BEST SEED SHOULD DO MORE  
THAN GROW GREEN GRASS**

**ADELPHI KENTUCKY BLUEGRASS** continues to be a top performer in density, disease resistance, drought, heat and cold . . . and is completely free of noxious weeds.

**ADELPHI** has proven itself to be the greener Kentucky Bluegrass that is preferred on Parks, Schools, Cemeteries, Ball Parks and Golf Courses. Ask anyone who has used it.

For information, contact:

**J & L ADIKES, INC.**  
Jamaica, N.Y. 11423

**NORTHRUP KING CO.**  
Minneapolis, Minn 55413

**VAUGHAN-JACKLIN CORP.**  
Bound Brook, N.J. 08805  
Downers Grove, Ill 60515  
Post Falls, Idaho 83854

**ROTHWELL SEEDS LTD.**  
Box 511, Lindsay  
Ont. Canada K9V 4L9

Other International Inquiries: **NORTHRUP KING CO.**, Minneapolis, Minn. 55413



Whitey Ford and Mickey Mantle play on their favorite grass

# Adelphi

(U.S. Plant Patent NO. 3150)

**THE GREENER KENTUCKY BLUEGRASS**