


A BEAUTIFUL LIE.



A BEAUTIFUL RYE.

There's no doubt about it. REGAL is the new turf-type perennial ryegrass that combines color, vigor and density for an elegant turf.

REGAL's dark green, fine-textured leaves look beautiful alone, or blend well with other turf-type ryegrasses. It has excellent tillering characteristics, and during periods of transition with bermudagrass, helps to maintain a lush, green, uniform turf.

And REGAL cuts clean, for a beautiful lie on tees, greens, fairways and roughs. Without ragged edges, or grainy areas.



Try REGAL—bred for rapid germination, exceptional vigor, density and persistence. Good disease resistance, too.

If you'd like more information on REGAL, contact North American Plant Breeders. P.O. Box 2955. Mission, Kansas. 66205 8-5256



First in the science of seeds.™

Circle 137 on free information card

REGAL

the ryegrass that gives your course a beautiful edge.

CONTROL FACTORS OF LEAF SPOT AND THEIR AFFECT ON SYMPTOMS

By Clinton F. Hodges, Professor of Horticulture and Plant Pathology, Department of Horticulture, Iowa State University

Turfgrass pathologists have endeavored to characterize symptoms of turfgrass diseases so that turf managers can diagnose specific diseases and their causal pathogens in the field. Such efforts have served the turf manager reasonably well for the more common diseases. However, increasing knowledge of environmental and cultural factors on disease symptoms continues to provide examples of variation in disease development and symptoms. Variations in disease development and symptom expression induced by environmental and cultural factors can be the cause of considerable consternation. This is becoming evident among species of the genus *Helminthosporium* and especially for *H. sorokinianum* (*Drechslera sorokiniana*). In time, however, understanding the various effects of environment and cultural aspects on disease expression by *H. sorokinianum* may be used to the turf manager's advantage. The purpose of this presentation is to summarize our knowledge of factors that can modify the symptomatology of leaf spot (*H. sorokinianum*) on Kentucky bluegrass.

Cultivar resistance and time

The use of fungicides is the primary means by which turfgrass diseases are controlled. Resistant cultivars and appropriate cultural practices also contribute to the control of some diseases.

The multitude of Kentucky bluegrass cultivars released in recent years has increased the hope among turfmen that resistance to various diseases will be improved. Resistance to diseases of Kentucky bluegrass induced by species of *Helminthosporium* leaf spot, however, must be approached with caution. Of 30 cultivars grown from seed (and vegetative parts) at the Iowa State University Horticulture Research Station, only five cultivars (Adelphi, Merion, Olymprisp, Rugby, and S. Dakota Cert.) had less than 30% of their leaves infected by *H. sorokinianum* in the spring of their fourth year of growth.

Most cultivars show increasing levels of leaf spot damage with each successive year's growth. In this regard, observations of S. Dakota Certified in subsequent years placed it among cultivars most susceptible to leaf spot.

It seems that the longer any cultivar is grown, the greater the incidence of leaf spot. The pathogen *H. sorokinianum* is a very aggressive parasite that attacks numerous species of grasses ranging from cereals to turfgrasses. The pathogen lives on dead tissue of the lesion it produces on the leaf and may produce a phytotoxin that is responsible for killing leaf tissue. It is very difficult to achieve high levels of resistance in cultivars against pathogens of this type. Under such circumstances, any cultivar that shows less than 30% of its leaves to be infected by *H. sorokinianum* after 4 to 5 years growth may be representative of higher levels of resistance to this pathogen.



Natural selection and observant turf specialists developed the earliest resistant varieties of Kentucky bluegrass for all turf managers.

Leaf age and symptoms

The leaf spot symptoms of *H. sorokinianum* are generally described as starting with small, circular, dark brown to purplish black spots that gradually enlarge and develop tan or brown centers. These lesions also may be surrounded by a zone of chlorotic tissue. This symptomatology is most common on the youngest leaves of the shoot in spring and fall, but such symptoms are greatly modified by progressively older leaves on the shoot.

Kentucky bluegrasses generally maintain 3 to 4 leaves per shoot; the two youngest leaves most commonly show typical leaf spot symptoms, while the two oldest leaves often have enlarged spots accompanied by chlorotic streaking on leaves. Occasionally, a single small leaf spot at the leaf blade base, or at the cut ends of older leaves, can result in the entire leaf turning yellow. Such symptoms on older leaves can be confusing in field diagnosis, particularly if younger leaves show few typical leaf spots. Variations of this type in symptom development have, on occasion, led pathologists to believe that perhaps different species of *Helminthosporium* were involved. It is known, however, that *H. sorokinianum* can produce a wide range of leaf symptoms that develop in response to physiological age of the infected leaf.

It is becoming increasingly clear that leaf age must be considered in all evaluations of *H. sorokinianum* symptoms. Severity of symptoms produced by this pathogen increase on each older leaf.

Environment and symptoms

Development of *H. sorokinianum* leaf spot on Kentucky bluegrass is markedly influenced by environmental factors. Recent research has es-

established that lesion development is influenced by photoperiod and light quality. Individual lesions increase in size on leaves as day length becomes shorter; and, as day length increases lesions become smaller. Under shorter day length, brown to purplish black portions of lesion increases in size as does yellowing that surrounds lesions and complete yellowing of infected leaves is accelerated. Day length also interacts with leaf age; as day length shortens, symptoms become more acute on each older infected leaf.

These observations suggest that leaf spot symptoms should be less severe in spring with increasing day length and more severe in fall with decreasing day length. This is, in fact, a common observation for *H. sorokinianum* on Kentucky bluegrass.

Quality of light reaching infected leaves also can influence appearance of leaf spot symptoms. Infected leaves exposed to increased levels of far-red light have larger lesions and more general yellowing than infected leaves exposed to normal levels of far-red light. As light passes through upper, younger leaves of a plant the proportion of far-red reaching older, lower leaves increases. This phenomenon may, in part, explain why older infected leaves tend to become completely yellow in the fall and early winter. During this time of year shorter daylength and the far-red light reaching leaves will tend to increase the rate at which they age and at the same time increase disease severity on infected leaves.

Effects of temperature and water (rainfall or irrigation) on *H. sorokinianum* leaf spot development are difficult to separate. Leaf spot is most active during the cool periods of spring and fall. As temperature (possibly combined with wind velocity) increases and rainfall decreases in summer months, number and size of leaf spots decreases. However, our field studies have shown that if the summer is relatively wet, or if irrigation is provided, leaf spot may occur on 13 to 60% of leaves throughout summer months depending on susceptibility of the cultivar. Midsummer leaf spot seems to be more closely regulated by moisture than by temperature.

Cultural practices and symptoms

Nitrogen fertilization often increases the severity of *H. sorokinianum* leaf spot on Kentucky bluegrass. When nitrogen stimulates leaf spot it is usually reflected by an increase in size of lesion and by some general yellowing of infected leaves.

Studies recently completed in our laboratory indicate that nitrogen and leaf age interact to influence leaf spot severity (i.e. the youngest leaves of nitrogen fertilized plants may have less disease than those of nonfertilized plants, but disease is increased on the oldest leaves by nitrogen fertilization). This suggests that the subject of nitrogen fertilization and leaf spot severity is related to the physiological age of leaves and that influence of nitrogen on disease may be modified by leaf age.

Herbicides represent another cultural tool that shows some potential for influencing severity of *H. sorokinianum* leaf spot on Kentucky bluegrass. Such commonly used herbicides as 2,4-D, MCP, and dicamba may enhance leaf spot development.

Nitrogen and leaf age interact to influence leaf spot severity. Some herbicides enhance development of leaf spot as well.

and dicamba may enhance leaf spot development. Like other environmental and cultural factors that enhance leaf spot, leaf age is directly involved in the interaction. Increase in diseased tissue on youngest leaves is minimal and severity of disease increases on each older leaf. In some instances, one small lesion on an older leaf of a plant exposed to MCP or dicamba can result in a rapid straw-colored blighting of the entire leaf. It is believed that auxin-like herbicides (2,4-D, MCP, dicamba) may increase the rate of aging in older leaves and predispose them to more severe disease.

Seasonal appearance of leaf spot

It is now known that leaf age, environment, and cultural factors can influence severity and appearance of *H. sorokinianum* leaf spot symptoms. On the basis of these observations, it is of interest to speculate on how these various factors might influence leaf spot during the various seasons of the year. The following outline is a speculative summation of how leaf age, environment, and culture practices might interact with leaf spot.

Spring

A. Appearance of Symptoms

1. Leaf spots small in early spring.
2. Leaf spots larger with some leaf yellowing in mid-spring.
3. Leaf spots smaller by late spring to early summer unless irrigation is provided.

B. Factors involved

1. The rapid flush of leaves in the early spring results in a very youthful shoot which inhibits leaf spot development. By mid-spring normal senescence of leaves on individual shoots has increased and such leaves show the larger lesions and the more severe yellowing symptoms.
2. By late spring increasing day length and reduced moisture levels (assuming irrigation is not provided) results in fewer leaf infection and less severe infections.
3. Applications of nitrogen or herbicides (auxin types) between mid-spring and early summer could increase severity of leaf spot. However, it is also possible that

Continues on page 62




"Saved money."
Jim Anderson, Lost Spur Country Club
St. Paul, Minn.



"Controlled brownpatch."
Larry Bunn, Blue Hill Country Club
Canton, Mass.



"3 week control."
Chris Myers, Bloomfield Hills Country Club
Bloomfield Hills, Mich.



"Best I've ever used."
Vince Spano, Hamlet Golf & Tennis Club
Delray Beach, Fla.

Chipco® 26019 is getting the w

The word is spreading fast and it's all good. "With Chipco 26019 we've pretty well licked our disease problems and it lasts up to 21 days!" said Larry Bunn at Blue Hill C.C.

Jim Anderson, Lost Spur C.C., emphasized 21 day residual control saying "The longer time between spraying has saved money and labor to keep us within budget." And Firestone's Jim Loke said, "With 200 to 300 players per day, the long spray interval adds extra convenience."

They're all impressed with just how effective Chipco 26019 is against the toughest disease problems like dollar spot, (including benomyl resistant dollar spot) brown patch, Helminthosporium (leaf spot, and melting out) and fusarium patch in the Northwest.

Bent Tree C.C. Superintendent Warren Stringer said, "Leaf spot was epidemic in proportions here last Spring. Weeks of rain left our usual fungicides ineffective. Chipco 26019 turned the tide."



"Extra convenience."
Jim Loke, Firestone Country Club
Akron, Ohio



"What I'll use in the future."
John Monson, Broadmoor Golf Club
Seattle, Washington



"Effective against leafspot."
Warren Stringer, Bent Tree Country Club
Dallas, Texas




"Eliminated dollarspot."
Robert Williams, Maidstone Club
East Hampton, N.Y.

Word from the guys that count.

Jim Bunn noted, "we controlled dollar spot and brown patch with two applications." And Chris Myers of Bloomfield Hills C.C. said that Chipco 26019 controlled dollar spot "longer than any other fungicide he used last year."

This season, ask your Chipco distributor or Rhône-Poulenc representative about Chipco 26019 . . . the turf fungicide that outperforms anything else you can use, with about half the number of sprays. Who says so? *You, the turf care professional.* And as far as we're concerned, that's the last word.

Rhône-Poulenc Chemical Co., Agrochemical Division, Monmouth Junction, NJ 08852



Circle 166 on free information card

Toro® presents a complete built as if they



COMMERCIAL 21-4
Our new rear bagger,
4 hp, 4 cycle, hand-
propelled mower.

COMMERCIAL 25-6
Our high capacity,
6 hp, 4 cycle, self-
propelled mower.

Had it with walk
mowers that break down before
they're broken-in?

You need a line
specifically engineered to take
the everyday wear and tear
of commercial use.

You need Toro Commer-
cial walk power mowers.

From rugged frames to
brawny engines to the tiniest

detail, we build them as if
they owe you a living.

Because that's what a
commercial mower is all about.

And, with our two
newest models, the Toro line is
complete:

For light to medium
size jobs, there's our new
Commercial 21-4 side discharge
mower.

It's a lower cost working
partner to our 5 hp Commercial
21-5 self propelled and Com-
mercial 21-5 hand propelled
mowers. Both are ideal for trim-
ming and medium size mowing
applications.

For jobs that require
bagging or a fine groomed
appearance, you'll want
our new Commercial 21-4 rear

line of walk mowers owe you a living.



COMMERCIAL 21-5
Our intermediate,
5 hp, 4 cycle, hand-
propelled mower.

COMMERCIAL 21-5
Our intermediate,
5 hp, 4 cycle, self-
propelled mower.

COMMERCIAL 21-4
Our new side discharge,
4 hp, 4 cycle, hand-
propelled mower.

21" COMMERCIAL
Our lightweight but
rugged, 2 cycle, hand-
propelled mower.

bagger model.

For really big jobs,
our 6 hp Commercial 25 gives
you smooth handling over
rough terrain and reliable
service day in, day out.

And, for quick trims to
light mowing, our 21" Commer-
cial 2-cycle mower is the light-
weight with heavyweight
features.

Want more information
on any or all of our Commercial
walk mowers? Call your Toro
distributor. Or, mail the coupon.



The Professionals™

Circle 180 on free information card.

Tell me more, Toro.

I'm interested in Toro Commercial
walk mowers. Please have my
distributor call me.

Name _____

Company _____

Address _____

City _____ County _____

State _____ Zip _____

Telephone _____
area code number

Mail coupon to: The Toro Company
Commercial Marketing Dept. WTT-030
8111 Lyndale Ave. S., Minneapolis, MN 55420

DISEASE MANAGEMENT IS BASED ON FIVE GENERAL AREAS OF EFFORT

By W.C. Stienstra, Associate Professor and Extension Specialist, Department of Plant Pathology, University of Minnesota.

In plant pathology disease control, I prefer the term Disease Management. Disease management is based on five general methods or traditional areas of effort.

First, resistant lines: this is and has been a major form of plant disease control or management with all plants. It is depended upon for disease control or management in crops such as cereals, corn, beans, hay and many others. In many of the more intensive crops, disease resistance is necessary to manage and produce even with the use of fungicides. The introduction of leaf spot resistance (i.e. Merion Kentucky bluegrass) changed the turf world. Today's term is "a revolution occurred." Resistant plants are, I believe, the "natural state" and we should (1) do all we can to utilize genetic material and (2) provide favorable environments for plant growth to express genetic potential.

Pathologists in general and myself specifically have not involved themselves to the extent they should in defining optimum conditions for turf growth.

The second general method of disease management is exclusion (i.e. quarantine). The best known quarantine for plant disease is at international ports of entry where ships, planes, cars, trucks and people are crossing from one country into another. Quarantine is also accomplished on a smaller scale as in foundation seed, commercial greenhouses, and many other cases where valuable stocks and certain crops are grown in confined areas with good sanitation procedures.

In turf leaf spot management, exclusion or quarantine has limited value.

The third general method is eradication (i.e. the attempt to remove the disease organism after it has entered an area). This is very difficult on a field scale once the organism has become established. However, in some confined areas this can be accomplished. The profits obtained from such a program often are quite high. This technique again has limited use in turf leaf spot management.

The fourth general method is sanitation, a term meaning clean up of an area to reduce chances of infection. Persistence of the individual plant manager in watching and using all procedures makes the difference between success and failure. This is important in turf leaf spot management.

The fifth general method is protection, the use of chemicals (i.e. fungicides). There may be too great a reliance on chemical protectants since they are easily used, readily available and response is expected where disease threatens. Fungicides are needed in disease management and are very useful in turf leaf spot control.

The five general methods of disease management are resistance, exclusion, eradication, sanitation, and protection. What management factors should the turf managers use when developing practical methods of leaf spot management in turf? *Helminthosporium* Leaf spot is one of the most

common and serious disease complexes attacking cool season turfgrasses in the U.S. Leaf spot can be very destructive during wet humid weather or in areas where the turf is sprinkled frequently especially in early evening. The more often grass is wet and the longer it remains wet, the greater are the chance of disease. *Helminthosporium* fungi are responsible for the gradual browning, thinning and melting out of Kentucky bluegrass cultivars and other susceptible grasses. As the disease progresses, large irregular turfgrass areas are yellowed, browned and finally killed. Once *Helminthosporium* fungi become established in a turfgrass stand, they remain an ever present problem.

The disease cycle of all species of *Helminthosporium* fungi is essentially the same. They survive from year to year as mycelium in dead grass tissues or in infected leaves, crowns, roots and rhizomes. Under moist conditions, tremendous numbers of spores are produced on this debris and carried to new leaf growth by air currents, mowers and foot and vehicular traffic. The spores germinate in a film of moisture and infect the leaves. Spore germination and infection of leaves can take place within a matter of hours when conditions are favorable. New spores are produced on these infected leaves within a few days which in turn spread to new leaf parts and neighboring plants. Thus the cycle is repeated. New leaf infections may occur as long as conditions remain moist and the temperatures are favorable for germination and growth.

Best management of leaf spot begins by providing an adequate soil zone for turfgrass root growth.

Best management of *Helminthosporium* leaf spot begins by providing an adequate soil zone for turfgrass root growth. Turf areas developed with planned surface drainage and adequate internal drainage will produce a vigorous network of plants (leaves, roots and rhizomes). Plants growing under these conditions will respond best to adverse environmental stresses.

When managing to prevent disease problems, it is important to remember the disease formula (i.e. you must have the host, the causal agent and a favorable environment for disease to develop). A vigorously growing plant is best suited to express its genetic resistance and to survive the period of disease stress and to recover faster.

Best management of *Helminthosporium* leaf spot continues with the selection and use of resistant

varieties. Leaf spot resistance in the form of Merion Kentucky bluegrass so changed the turf industry that now we recognize other summer diseases of turf. Before Merion, summer disease problems were always caused by *Helminthosporium*. *Helminthosporium*-incited diseases rank among the most important fungus disorders of turfgrasses. Characterized by leaf blighting, leaf abscission, root, rhizome, stolon and crown rots, it is not uncommon in certain seasons for this group of diseases to become the limiting factor in turfgrass production.

The best control of *Helminthosporium* leaf spot is obtained through the use of resistant varieties. Many lines exhibit nearly complete resistance to *Helminthosporium* under field conditions. Select high quality, disease resistant seed or sod that is locally adapted or a mix of locally tested grass varieties resistant to one or more *Helminthosporium* diseases.

But what if you already have a turf area, and do not want to begin a major rebuilding program? What should you consider in a *Helminthosporium* leaf spot sanitation program?

First - mow the bluegrass or for that matter, rye and fescue at the recommended height. *Helminthosporium* leaf spot is most damaging when close mowing occurs, so avoid close clipping at all times and especially when the leaf spot stage is a serious threat and when conditions are favorable for disease development. Mow the grass frequently, removing no more than one third of the leaf surface with a single mowing. Removal of clippings may further reduce the available food base for *Helminthosporium* leaf spot development.

Second - water turf areas as needed, wetting the soil to a depth of 4-6 inches. Repeat every week if nature does not supply adequate amounts. Apply supplemental water during or immediately after light showers during any dry periods. The value and importance of the soil can and should be noted here. Soil that is slow to wet and soil that has an inadequate water holding capacity forces you into frequent light sprinklings, and water-logging the soil surface, which promotes disease development.

Third - fertilize based on soil tests and nutritional needs of the plant. Avoid heavy applications of water soluble fertilizers, when *Helminthosporium* leaf spot disease is expected. Fertilize to maintain a uniform level of soil nutrients in the root zone, following the local recommendations for the grasses being grown. Consider fertilizer needs beyond having the greenest turf in the area.

Fourth - reduce thatch accumulation or mat to less than one centimeter by using a power rake or an aerifier. These operations are usually accomplished in the spring, fall or both when turf growth and recovery are rapid for cool season species. Thatch reduction is a process not completed in one treatment, just as thatch accumulation occurs over the entire growing season. Management of thatch



Development of resistant lines

is a major form of plant disease control and utilizes known resistant material

accumulation and its removal is needed in most turf areas.

Fifth - consider modifying the local environment. Is the turf growing under dense shade, in areas with restricted air movement? Could trees or shrubs be removed or pruned? If so, will more light penetrate to the turf and will greater air movement over the turf speed the drying process thus reducing disease conditions?

Sixth - If *Helminthosporium* diseases are not adequately managed by cultural treatments, fungicide sprays may be needed on a responsive-preventative schedule. A responsive - preventative schedule means fungicide applications begin with the development of the disease and continues every seven to fourteen days during the period when conditions permit disease development. This spray program requires disease awareness, disease diagnosis, time, money, and for the average individual may not be practical. If only a few sprays will be applied, to be most effective, most of the time, the applications should occur in the spring, fall or both when cool wet growing conditions are common. If spraying is delayed until leaf blighting is severe, or thinning is obvious, the results will be poor.

Spraying fungicides or "all the world loves a universal green paint" is the "take two aspirins and call me in the morning" philosophy. Spraying, the easy, accepted, and indeed expected disease response, often may not be well understood. We can cite chapter and verse on preventative schedules, curative schedules, manufacturer directions, package label rates, application intervals, compatibility, safe use, pressure, volume, time of day, uniformity, nozzles, nozzle spacing, temperature, moisture, grass condition, grass potential, chemical persistence, spreaders, stickers, rainfall, past fungicide use and resistant isolates.

Continues on page 62

Wherever annual weed

"Out here, crabgrass is our biggest complaint, and Balan in the spring works like a charm."



"Around these parts, we're faced with goosegrass/crowfoot in the summer, then when our Bermuda-grass goes dormant, Poa annua's the problem. So we use Balan twice."



"Sometimes we use Balan in the early spring for crabgrass, but if goosegrass is a problem we apply Balan later to get it as it germinates."



"If you've a goosegrass/crowfoot problem, a shot of Balan early in the summer takes care of that problem; but nothing makes our courses look worse than green patches of Poa after the Bermuda's gone dormant, so we apply Balan in the fall, too."

...the solution is Balan

The annual grass problems may differ from one part of the nation to another, but the solution is the same . . . Elanco's Balan. All across the country, Balan users have proven for themselves that a regular Balan program is their most effective means of controlling the hard-to-control annual weed grasses. Balan's granular form makes it as quick and easy to apply as fertilizer.