



NO OTHER HERBICIDE COVERS THE LANDSCAPE LIKE CHIPCO® RONSTAR®.

HERBICIDE

IT'S SAFE ON THE BROADEST RANGE OF TURF AND ORNAMENTALS.

You have to be careful when you put down most pre-emergence herbicides. They're safe on turf *or* ornamentals. Not on both.

Enter CHIPCO® RONSTAR® preemergence herbicide. The only product you can use safely to weed the whole landscape: turf, woody ornamentals, ground cover, and trees.

RONSTAR herbicide solves a lot of problems. And simplifies your inventory and application. It causes no root

pruning. Needs no incorporation. Won't leach or move into untreated areas.

RONSTAR GIVES EXCELLENT WEED CONTROL.

Just one application of RONSTAR herbicide gives effective, season-long control of over twenty of your toughest annual broadleaf and grassy weeds.


FOR A GOOD-LOOKING LANDSCAPE, TAKE A GOOD LOOK AT RONSTAR HERBICIDE.

If you'd like to find a simpler, safer way to keep weeds under control, find out more about

Circle No. 152 on Reader Inquiry Card


the preemergence herbicide that covers the landscape: CHIPCO RONSTAR herbicide.

See your dealer, or write: Rhône-Poulenc Inc., Agrochemical Division, Monmouth Junction, NJ 08852.



RONSTAR SPREAD THE WORD.

Please read the label carefully, and use only as directed.



SWELL
PAT PENDING
SF
FOOPER

HOW TO TAKE OUT SURFACE FEEDERS AND GRUBS IN ONE SWELL FOOP.

To take out surface feeders like sod webworm larvae, chinch bugs, billbugs, and flea beetles, just follow these simple steps.

Step one, get some [®]OFTANOL 2 Insecticide. It's the liquid formulation of OFTANOL. If you think OFTANOL is too expensive to use on surface feeders, check out step two.

Step two, mix it at the *economical rate* and spray where the bugs are. It'll work.

The same product, OFTANOL 2, does a bang up job on grubs. Mix it at the recommended rate and spray it on. Use OFTANOL anytime, as a preventive in the spring or as a curative, whenever grubs are a problem.

Surface feeders and grubs, two serious problems with one serious solution. OFTANOL 2.

OFTANOL is also available in a granular formulation. Always read the label before use.



**OFTANOL 2.
WHEN YOU'RE SERIOUS
ABOUT SURFACE
FEEDERS AND GRUBS.**



Mobay Chemical Corporation
Specialty Products Group
Box 4913, Kansas City, Mo. 64120

OFTANOL is a Reg. TM of the Parent Company of
Farbenfabriken Bayer GmbH, Leverkusen.

Circle No. 139 on Reader Inquiry Card

construct a frass-packed tunnel from the chamber to the outer bark cortex before overwintering. This larval channel undoubtedly expedites and facilitates emergence by the adult the following spring.

Larvae pupate in May or June. Ten to 14 days later, the adult chews a D-shaped emergence hole that conforms to the shape of the thorax before emerging. If the hole is not shaped properly before the beetle tries to climb free, it may become permanently stuck in the hole.

Most adults emerge during a 4 to 6 week period in late spring and early summer and must feed at least 8 to 10 days before they are capable of laying eggs. Females are apparently discriminating egg layers because each produces only 10 to 40 eggs during a 3 to 6-week life time.

Eggs are glued to bark surfaces, usually near the v-shaped, black areas at the junction of branches with limbs or limbs with the trunk. Larvae hatch by chewing through the egg chorion at its point of attachment to the bark after a 1 to 2 week incubation period.

Larvae feed initially in the cambial zone, stimulating some trees to form a callus-like tissue that shows-up as ridges in the bark. Since larvae feed horizontally and switch directions at acute angles, the raised areas on the bark are in a zigzag pattern.

Other researchers have hypothesized that this feeding pattern minimizes vascular pressures on larvae, improving their chances of surviving the tree's natural resistance.

The primary symptom of bronze birch borer attack is dieback of the uppermost branches of the tree. When this is noticed, the bark should be inspected for the presence of narrow, raised areas in an S-shaped configuration.

If this symptom is present on birches with top-dieback, you can be certain that bronze birch borer is the culprit. If D-shaped emergence holes are detected, you know that the infestation must be dealt with immediately if there is any hope of saving the tree.

Control strategies

There are four factors that are of primary importance in developing a borer control strategy:

- 1** maintaining plant vitality,
- 2** knowing when adults begin emerging and for how long emergence continues,
- 3** knowing which labeled insecticides are most effective, and

4 achieving thorough coverage of all bark and leaf surfaces.

Maintaining plant vitality

Many people do not realize that the urban forest is a hostile environment for woody plants. Soil texture, grade, and chemistry have often been dramatically altered by activities of urbanization, and exotic trees are planted in locations for which they are not adapted.

Landscape architects and others who choose trees for specific sites should do a better job of selecting

Very little is known about buprestids and their relationship to host plants.

trees that can be expected to thrive, or at least survive, in the area where they will be planted.

Whenever possible, sites should be altered to meet the requirements of the tree to reduce long-term maintenance costs associated with weakened trees.

Recognizing that borers are opportunists that exploit weakened trees, it should be obvious that anything we can do to improve tree vitality will reduce borer infestation and damage.

The following cultural practices will improve or help maintain tree vitality, helping the tree to help itself in its battles with borers and perhaps other insects and diseases.

Pruning

Landscape trees should be pruned properly and on a timely basis following recommendations recently published by Dr. Alex Shigo of the Northeast Forest Experiment Station in Durham, NH.

Trees like birch should be pruned after August 1 to minimize bleeding and problems with canker-causing fungi. Proper pruning will improve both the confirmation of the canopy and the energy economy of the tree, leading to improved vitality.

Fertilization

In native habitats (i.e., forests there is a cycling of nutrients as leaves and branches are shed, fall to the ground, and decompose. The same process occurs with understory plants, including shrubs, forbs, and grasses, resulting in a continuous supply of nutrients.

This cycling does not occur in much of the urban forest because

leaves and other refuse are discarded annually. Competition by dense plantings of grasses further depletes soil nutrients. Consequently, urban trees will benefit from fertilization.

An inorganic, slow-release fertilizer should be used in fall after one or two hard frosts. It can be broadcast on the soil surface, injected under pressure, or placed in drill-holes. If annual fertilization is practiced, 4 pounds of actual N per 1000 square feet should be used. I recommend 6 pounds of N per 1000 square feet every other year.

Top-dressing may be the most cost-effective method, but aeration supplied by injection or drill-holes may provide additional benefits.

Mulching

Bark damaged by maintenance equipment commonly serves as an oviposition site for borers.

Whenever possible, trees that are known to be hosts of clearwing borers should be mulched so that equipment does not need to be used close to the tree trunk. Of course, mulching also reduces evaporation of water from the soil, minimizes rapid changes in soil temperature, and reduces weed competition.

Mulch should not be more than about two inches in depth to minimize chances of creating an environment at the base of the trunk where microorganisms may flourish and eventually invade the tree.

Watering

There is a common misconception that large trees have deep roots, so they are not impacted by drought.

The reality is that large trees often suffer more from periods of low rainfall than smaller trees. In any case, trees will benefit from weekly watering whenever summer or fall drought occurs. Watering should be done slowly so that it percolates to the roots rather than running-off to areas that will not benefit the target tree.

This is one tactic in which the client can participate, if they are willing.

Debilitating insects

Defoliators and sucking insects should not be allowed to reach levels that debilitate trees, thereby reducing their natural resistance to opportunistic pests like borers. A high degree of control is not necessary because low levels of pests usually do not reduce tree vitality significantly. Consequently, insecticidal soaps, horticultural spray oils, and bacterial insecticides can be used to minimize reliance on conventional insecticides.

Implementation of a tree care program that includes the above prac-

*Tough mowers...
from rough cut
to smooth finish.*

The CM Series

Extra Heavy Duty
Hydrostatic Rotary Mowers for
Commercial/Municipal Users

CM 20-60/72

Superior performance in both rough cut and refined mowing at speeds to 11 miles per hour. The "CM" gives the commercial user a tough, low maintenance machine that mows better at top speed than other mowers at half the speed. And does it with the added safety of rear discharge.

Kut-Kwick. More than forty years of experience building professional mowers for professional users.

kut-kwick

P.O. Box 984
Brunswick, Georgia 31521
Telephone 912-265-1630

Call today for complete information.
Toll-free 1-800-248-5945
In GA. 1-800-592-5945

Circle No. 233 on Reader Inquiry Card

tices will lead to a more vital urban forest and a reduction in insect pest problems. But, there will always be times when a borer infestation must be dealt with aggressively to save the tree.

Defining adult emergence

Detection of presence and delineation of the emergence period for several clearwing moth borers can be accomplished with pheromone traps. Conrel Clearwing Borer Traps (available from Conrel, 110 A St., Needham Heights, MA 02194) can be used to

capture male moths of the following borers: banded ash clearwing (looks much like lilac borer but emerges in August and September rather than May and June), dogwood borer, lilac borer, lesser peachtree borer, peachtree borer, an oak borer, and rhododendron borer.

The traps are simply placed in a convenient location in your service area and checked periodically beginning about two weeks before emergence of the target borer species is expected. Directions provided in the trap kits and color photographs make

the traps useful even to practitioners who are unfamiliar with insects.

Borer sprays are applied at specified time intervals after the first male moth is captured.

In most cases, clearwings that attack trees can be controlled with a single application of an effective insecticide, if it is timed properly.

We have been unable to detect the presence of long-range sex attraction with bronze birch borer, so monitoring adult emergence is laborious, requiring daily or weekly inspection of infested trees or bolts.

Recently, we have described a way to predict bronze birch borer emergence based on local temperature. Degree-Day ($^{\circ}\text{D}$) insect activity models are based on a threshold temperature and a starting date (the date on which you begin accumulating $^{\circ}\text{D}$ above the developmental threshold).

Careful analysis of several threshold temperatures and starting dates in Columbus and Wooster, OH, indicate that these values must be prescribed for a given geographical area.

For example, in Wooster which is about 50 miles south of Lake Erie, the best threshold temperature for predicting 10 percent adult emergence was 10°C (50°F) with a starting date of April 1. In Columbus which is latitudinally about 50 miles south of Wooster, the best threshold temperature was 8°C (46°F) with a starting date of May 1.

We don't know how well these models will fit other locations. Someone in each area of the country who wishes to develop this kind of model for predicting bronze borer emergence will need to collect infested wood in early spring, record seasonal emergence and temperature, and then work with a statistician to determine if one of our models accurately describes birch borer emergence in their area.

The importance of using a predictive model based on local temperature is that a borer population may emerge as much as two to three weeks earlier or later than usual during a particular year. This accelerated or delayed emergence is directly related to temperature and will directly influence the success of your borer control program.

Effectiveness of available insecticides

Dursban (chlorpyrifos), Turcam and Dycarb (bendiocarb), lindane, and Thiodan (endosulfan) are registered for use against borers. I have not worked extensively with Thiodan but believe the other materials to be at

continued on page 60



It's a winner!

The finest golf courses in the world are fertilized with Milorganite . . . naturally

Milorganite is easy to apply, without danger of burning, so watering-in is never necessary. It never deteriorates or hardens during storage, and Milorganite contains natural organic nitrogen, phosphorus, and iron. All are released gradually by nature's soil bacteria when and as the grass needs them to insure uniform color with even, prolonged, and healthy growth.



America's Number One Naturally Organic Fertilizer For Over 50 Years

Milwaukee Metropolitan Sewerage District
735 N. Water Street, Milwaukee, WI 53202

Circle No. 133 on Reader Inquiry Card

Here you see our finest products. But not our best work.

Our finest products are lush, green, and vigorous. Our best work is knowing we can stand behind them.

Years of field experience. Continuous research. Nationwide testing. All these are hidden in every blade that sprouts from a Northrup King Medalist® Turfgrass Mixture.

Mixtures such as Athletic Pro® II,

Northrup King Medalist Turf Products, P.O. Box 959,

Medalist North, Premium Sod Blend, Landscape Pro, and the Medalist Winter Overseeding Products. And we never stop improving them.

For help with your toughest turf problems, contact us. Then we can show you more of our best work. Listening.


And answering with turfgrass mixtures that work.



TM Minneapolis, MN 55440 ©1985 Northrup King Medalist

Circle No. 143 on Reader Inquiry Card

Here you see our finest products.
But not our best work.





Above and below the call of duty.

Only D-z-n® destroys 52 surface and sub-surface insects.

You know that insects can wreak havoc in lawns and ornamentals.

But D-z-n diazinon gets them where they live. Above and below the soil line. Sod webworms, chinch bugs, armyworms, aphids and more. Including grubs, cutworms and other sub-soil pests. With a quicker kill and longer residual control.

When safety counts.

D-z-n diazinon is so safe it's non-restricted. Use it near homes, hospitals or schools. You need no special clothing or protective equipment.

Mix D-z-n with fertilizers and fungicides. In liquid, granular or powder formulations. The choice is yours. D-z-n diazinon. From the people who invented diazinon over 20 years ago.

Ciba-Geigy, Ag Division, Box 18300,
Greensboro, NC 27419.

D-z-n. The bug stops here.

CIBA-GEIGY

©1984 Ciba-Geigy Corporation

Circle No. 245 on Reader Inquiry Card

BORERS from page 56

least as effective or better.

Chlorpyrifos is available in three formulations (Dursban 2E, 4E, 50WP) and is registered for use against all borers on all trees and shrubs. The usage rate is commonly one pound of active ingredient per 100 gallons of water applied to runoff. Currently, Dursban is the best choice for controlling clearwing borers. In my experience, a single application of Dursban, 10 to 14 days after first adult emer-

gence as monitored by a pheromone trap, will provide season-long protection.

The exception would be lesser peachtree borer that emerges from May through September in much of its range. Two or three applications at 5-week intervals may be required to control this species. Lindane is somewhat hard to find these days but is an effective borer control product with an excellent safety record.

The most commonly used formulation is the 20 percent liquid that contains 1.8 pounds of insecticide per

gallon. It can be substituted for Dursban at the one pound of active ingredient per 100 gallons of water usage rate for controlling clearwing borers. To my knowledge, products containing bendiocarb have not been evaluated against clearwing moths. We plan to do this in the near future.

Turcam 76WP (or Dycarb) is my choice for controlling bronze birch borer, although Dursban has also provided excellent control. Again, the one pound active ingredient usage rate applies. The initial thorough-coverage spray should be applied just prior to first adult emergence.

In northeastern Ohio this is usually in late May (20-25). Two repeat applications are made at two-week intervals. The spraying should be initially focused on bark of the trunk and all branches, even in the top of the tree, but leaves should also be sprayed because adults must feed for at least several days before laying eggs.

By spraying before adults begin emerging, repeating the application throughout the major emergence period, and covering both bark and leaves thoroughly we:

- 1 kill adults as they construct their exit hole,
- 2 perhaps kill eggs or larvae when they eat sprayed bark while chewing through the egg chorion at its point of attachment to the bark, and
- 3 kill adults when they contact bark or consume or contact treated foliage.

When spraying trees to control clearwing borers, application can be delayed until after oviposition, because larvae hatch and then crawl over the bark surface before choosing an entrance site.

However, with flatheaded borers, the spray must be applied before oviposition, since larvae construct their entrance hole at the point on the bark where the egg is attached. Any borer control strategy should be coordinated with a tree health care program, including removal and destruction of borer infested wood for top-killed and dead trees, and cultural practices that have already been enumerated.

Borer control procedures have now been developed that can be used by arborists and other landscape managers to minimize damage done by borers, some of the most damaging insect pests in the urban forest. **WT&T**

The author encourages all applicators of insecticides to read the container label carefully, following all instructions. Recommendations provided in this article are based on experience by the author. The author wishes to thank Rodney C. Akers for providing information on D models for predicting emergence of bronze birch borer adults.

• **FASTER ESTABLISHMENT**
• **BETTER SURVIVAL**

TREES • SHRUBS • TURF • SOD

Viterra[®]
Gelscape[™]

DESIGNED TO MAXIMIZE PROFITS FOR THE PROFESSIONAL APPLICATOR

CALL OR WRITE FOR MORE INFORMATION:
Nepera Inc., Route 17, Harriman, New York 10926
Telephone: (914) 782-8171

NEPERA  Nepera Inc., Route 17, Harriman, NY 10926
A Division of The Clorox Company

Copyright 1985

Circle No. 163 on Reader Inquiry Card