




Maximum nutrient availability

Maximum availability is indicated by the widest part of the bar



sides of the spreader.

With a liquid application method, fertilizer is either solubilized or suspended in water and sprayed on the turf. The amount of water normally varies from one to five gallons per 1,000 sq. ft.

Spray equipment can be broadly classified into either low pressure spray booms or high pressure (hydraulic) sprayers. Both types feature a tank, pump, pressure regulator, strainers or screens, and nozzle(s).

Low-pressure spray booms, as the name implies, are operated at pressures in the range of 15-60 lbs. per sq. in. (psi) and deliver one gallon or less of spray per 1,000 sq. ft. They are designed to be driven over large areas delivering the spray from a series of nozzles in distinct swaths. They are often used on golf course fairways.

High-pressure sprayers can create several hundred pounds or more of pressure. They use a hose and hand-held nozzle for directed application, and are used by lawn care companies.

TREE AND SHRUB FERTILIZATION

Landscape trees and shrubs are often subject to adverse soil and environmental conditions. Compacted soils, poor drainage, restricted root areas, highway salts, air pollutants and com-

Exact fertilizer needs are difficult to judge in interiorscape maintenance without measuring light at strategic locations.

petition from turfgrass contribute to plant stress, thus increasing the importance of regular fertilization.

Vigorous trees are more resistant to insects and disease, more attractive, and a greater asset to properties.

Only nitrogen, phosphorus and potassium are normally applied in tree fertilization. However, supplemental micronutrients such as iron and manganese may be necessary for certain species growing in alkaline or sandy soils.

Plants often respond to nitrogen applications with dramatic improvements in shoot growth and leaf color. Because nitrogen is transitory in soils and relatively large amounts are used by plants, soil analysis is not particularly useful. Heavy applications of nitrogen alone may stimulate shoot growth more than root growth, disturbing the natural root/shoot ratio.

The need for supplemental phosphorus and potassium is more difficult to determine since they do not produce a visible response except on young or newly-transplanted trees and shrubs.

Results from field studies also have been inconsistent because of differences in soil, age, condition and location of test species, and the timing and method of application. Where reliable soil tests are not available for phosphorus and potassium, most arborists use a complete fertilizer on all trees and shrubs.

Since arborists must be concerned with trees and shrubs in a wide variety of soil conditions, the most practical approach to fertilization is to provide an effective formulation for trees and shrubs in a geographic area.

Specific soil/plant deficiencies may be addressed, if necessary, on an individual basis. In most cases a 3:1:1 (or similar) ratio is satisfactory, but more potassium and/or micronutrients may be advisable in sandy or alkaline soils, particularly for ericaceous or other so called "acid-loving" plants.

Application rates

Most fertilizer recommendations are based on the size of the growing area for shrub beds or the branch spread for individual trees and shrubs.

Fertilizer recommendations based on trunk diameter can result in over-fertilization and plant damage if the root system is restricted by paved areas, foundation walls, or other obstructions.

Three pounds of actual nitrogen per 1,000 sq. ft. per year, or six pounds every other year, will keep deciduous trees and shrubs healthy and vigorous. If leaf color, annual growth or general vigor is unacceptable, six pounds of nitrogen per 1,000 sq. ft. may be applied annually.

Broadleaf evergreens, small shrubs, flowering trees and recently-transplanted or declining trees are more sensitive to fertilizer salts. These plants should receive only about one-half the recommended rate, particularly when quick-release fertilizers are applied. The risk of injury to sensitive plants may be reduced by splitting the recommended annual amount into two or more applications.

The amount of fertilizer per 1,000 sq. ft. of root area can be calculated by dividing the percent nitrogen on the fertilizer bag into the desired nitrogen

continued on page 34

NOR-AM's TURF PRODUCTS DON'T SIMPLY NOURISH YOUR TURF... THEY PROTECT IT

Now achieve fine turf with fewer problems...thanks to NOR-AM's Total Turf Management Program. This unique program brings you a broad selection of high-performance products that assure healthier turf with better root development, fewer insect problems and less disease destruction.

As a service to the industry, NOR-AM has developed a Total Turf Management Program. This program includes a turf management calendar guide, brochures on turf diseases, insect control and fertilization techniques, in addition to timely turf management reports. To enroll in this helpful program, fill out the coupon below or contact your local NOR-AM sales representative.

MAIL COUPON TO:

**NOR-AM Chemical
Company
3509 Silverside Road
P.O. Box 7495
Wilmington, DE 19803**

Attn: Communications Dept.

WWT

YES, I would like to be enrolled in NOR-AM's Total Turf Management Program. Please send me further information.

Company _____

Name _____

Title _____

Address _____

City _____ State _____ Zip _____



Nitroform®

Slow predictable release by soil bacteria provides constant 38% nitrogen when most needed for even, sustained growth.

Nitroform reduces the risk of nitrate contamination, decreases thatch build-up, is nonburning and nonstreaking, resists leaching even in sandy soil, and assures better root development.

Turcam®

An advanced broad-spectrum carbamate insecticide that controls white grubs, chinch bugs, sod webworms, mole crickets and many other turf and ornamental pests.

Turcam is odorless, works well in spray equipment, won't tie up in thatch, won't damage turf or ornamentals, and should be your product of choice when long residual control is required.

Proxol®

Economical Proxol 80SP Insecticide works fast on surface-feeding sod webworms, armyworms and cutworms, and readily penetrates thatch to control grubs. Proxol is easy to apply as a spray, has no unpleasant odor, and no long-term residual build-up in the soil.

Progress®

A selective herbicide that effectively controls annual bluegrass, crabgrass and other troublesome weed species.

Acti-dione®

Broad-spectrum family of turf fungicides that provide four-season disease control. Acti-dione has proven to be effective without creating resistance problems.

Acti-dione TGF® can be tank-mixed with Bayleton® or Daconil® to complement or improve disease control with significant cost-per-acre savings.

Banol®

Specialized fungicide that controls pythium and phytophthora in turf and ornamental plants. Preventative control with Banol stops the spread of pythium by mowers, foot traffic and surface water; it also means fewer applications, lower rates and savings in labor, fuel and equipment.

 **NOR-AM**

NOR-AM CHEMICAL COMPANY

3509 Silverside Road, P.O. Box 7495
Wilmington, DE 19803

**Total Turf Care from the
Turf Managers**

Circle No. 130 on Reader Inquiry Card

IMPORTANT: Please remember always to read and follow carefully all label directions when applying any chemical.



Spray systems such as the one shown above are used to apply nutrients to turfgrass.

NITROGEN FERTILITY LEVELS

lbs./growing month/1,000 sq. ft.

HIGH .5-1.5	
Kentucky bluegrass	bermudagrass
creeping bentgrass	
MEDIUM .4-1.0	
zoysiagrass	colonial bentgrass
tall fescue	velvet bentgrass
perennial ryegrass	St. Augustine-grass
annual bluegrass	
LOW .2-.6	
carpetgrass	red fescue
chewings fescue	
VERY LOW .0-.4	
blue grama	bahiagrass
buffalograss	centipedegrass

per 1,000 sq. ft.

For example, to determine the amount of 30-10-10 fertilizer required to apply six pounds of nitrogen per 1,000 sq. ft., divide .30 into 6, which equals 20 pounds.

Application timing

Although the roots of woody plants may elongate during the growing season, active root growth most often occurs in early spring and late fall when soil temperatures are cool and leaves offer little competition for water and nutrients.

Fertilization is most effective when supplemental nutrients are available during periods of optimum root growth. Soluble nitrogen fertilizers, because of their short residual in soils, should be applied between October and December and/or between February and April. Controlled-release nitrogen ensures availability in the root zone for a relatively long period, depending upon the nitrogen source's solubility. The application timing of these fertilizers may not be a major concern.

Application techniques

Supplemental nutrients can be supplied to landscape plants through foliar sprays, trunk injections, or applications on or beneath the soil surface. Though each method has advantages in specific situations, woody plants in most cases respond best to soil applications.

Surface applications

Nitrogen fertilizers can be applied to the soil surface, since nitrates are highly mobile and will move downward into the root zone. When fertilizing woody plants in sodded areas, surface application should be limited to no more than three pounds of nitrogen per 1,000 sq. ft. from a controlled-release source. However, since turfgrasses in the application zone may be injured or respond with undesirable succulent growth, sub-surface applications are often used on trees and shrubs in quality lawns.

Fertilizer with phosphorus should not be applied to the soil surface. Phosphorus is bound to soil particles and does not move downward to contact the absorbing roots. Surface applications of phosphorus may also stimulate annual bluegrass, which is undesirable in home lawns.

Drill hole technique

Fertilizer can be placed in the root zone by dividing the recommended amount of fertilizer equally among

drilled holes in the ground. For trees, holes should be 12 to 18 inches deep and 18 to 24 inches apart, beginning two to three feet from the trunk and going two to three feet beyond the drip line.

To prevent turfgrass injury, fertilizer should be at least four inches below the soil surface. Calcined clay, perlite, or other soil amendments can be used to fill the top of the hole or, in quality lawns, a plug of grass can be removed before drilling and replaced after adding fertilizer.

Soil injection

Liquid soil injection is a fast, economical alternative to the drill hole technique. A hydraulic sprayer operated at 150-200 psi and an injector probe inserted about 12 inches into the soil are needed. The injections are normally in a grid pattern about three feet apart within and slightly beyond the tree canopy.

Soil injection can mean more thorough nutrient distribution than the vertical hole technique, and generally can be done in about one-fourth the time.

Unfortunately, most soluble fertilizers have a high burn potential and soluble nitrogen may leach from the root zone. It may remain in the root zone for as little as six weeks.

Because of the limits of liquid soluble fertilizers, suspension fertilizers for soil injection are becoming popular. Ureaformaldehyde is particularly effective as a controlled-release nitrogen source in soil injection systems. Its release rate is not greatly affected by particle size. Suspended in water, powdered ureaformaldehyde can be injected into the soil by hydraulic pressure.

Two recent introductions, soluble methylol and methylene ureas, have a lower burn potential than urea or other soluble nitrogen sources.

Other methods

The aerofertil technique injects dry fertilizer by blasts of air into drilled holes in the soil. This method is similar to drill hole application, but provides additional aeration by breaking up heavy or compacted soils.

Fertilizer stakes or spikes are driven into the ground at intervals beneath the drip line of trees and shrubs.

Although they contain enough fertilizer, spikes are expensive and not as effective as other fertilization methods. One or two spikes per inch of trunk diameter provide only a small amount of fertilizer, not all of which comes in contact with the root system since there is little lateral distribution within the root



Dick, ProTurf already has a 15-0-30 High K fertilizer for fairways. Why did you feel you needed something for greens?

"On greens, you want a fertilizer with a very small particle size. Much smaller than our fairway product designed for the big rotary spreaders."

Why does particle size make a difference? "The smaller particles disperse evenly from the spreader, filter down into the turf, and disappear when you water. There's less chance they'll affect a putt or stick to shoes and get tracked around."

That sounds good for the golfers. "Superintendents will like it, too. In addition to being small, each particle is homogeneous. So, even at half the regular application rate, there's no chance of getting the kind of speckling you might see if you used a coarse blend. And more particles per square inch produces a more desirable, uniform greening response."

"We developed this new version of 15-0-30 High K fertilizer especially for greens. And golfers."

Dick Westfall, Scotts Research project leader (Fertilizer R&D), talks about new High K Greens Fertilizer.

Now tell me why High K will be good for my greens.

"High potassium levels have been found to enhance turf performance, particularly under stress conditions—improving drought hardiness, resistance to disease, and wear tolerance. There's also evidence that high potassium reduces winter kill in bermudagrass."

Why is there so little nitrogen? "To prevent over-stimulating top growth. For example, High K Greens can be used at double rates to build potassium levels in sand greens and you'd still be applying only 1 pound of N per 1,000 square feet. Or you can use it at regular rates on any greens that are on a low nitrogen diet. It can also be used at any time of the year."

Anything else we should know about ProTurf High K?

"If you like what it does for fairways, you'll love what it does for greens."

For more information on new High K Greens Fertilizer, call your ProTurf® Tech Rep. Or call Scotts® direct at 800-543-0006. In Ohio, call collect 513-644-2900.

Circle No. 149 on Reader Inquiry Card





Annual Nitrogen Requirement of Turfgrasses*

Species	Length of Growing Season, Months	Nitrogen per Season lbs./ 1,000 sq. ft.	Variations in Management
Cool-Season:			
sheeps & hard fescue	4- 8	0- 3	low maintenance; roughs
red fescues	4- 8	1- 3	low maintenance to good care
Kentucky bluegrass	5-12	2- 8	lawns, fairways
bentgrasses	4- 8	1- 4	medium care, lawn, fairways
bentgrass, greens	5-12	6-15	clippings removed, forced growth
Warm-Season:			
zoysia	6-10	1- 6	adequate cover
common bermuda	7-12	2- 8	most variable
St. Augustine, bahia	10-12	2- 8	warm areas, lawns
bermudagrass, fairways and tees	5-12	4- 9	good management
bermudagrass, greens	8-12	8-20	may rest over winter

*Adapted from Turf Managers' Handbook by William H. Daniel and Raymond P. Freeborg, published in 1973 by Harvest Publishing Company, New York, N.Y.

zone of most soils.

Foliate sprays, trunk injections and implants can supply limited nutrients to woody plants. They are recommended for micronutrients whose availability is reduced by alkaline soils. These methods are most effective when a single micronutrient is deficient.

FERTILIZATION OF TREES AND SHRUBS IN CONTAINERS

Growing trees and shrubs in landscape containers is common where plants are desirable but suitable planting sites limited.

They need careful attention because the reservoir of available growing media—minerals and water—is much smaller. Container soils, often wet and poorly aerated, can leach excessively and require a regular fertilization program.

In general, recommended fertilizer rates for landscape plants based on square footage have been successful in maintaining container-grown plants.

A complete fertilizer applied at an annual rate of 0.5 to 1.0 ounce of nitrogen per 10 sq. ft. of container soil surface is usually used. However, because of the wide selection in plant material and variations in container design and growing media, fertilizer needs are best determined by soil and tissue analysis.

Container plants can be fertilized by dry, foliar, and liquid application.

As with landscape plants, foliar applications are usually limited to micronutrients.

Foliar fertilization is effective where soil conditions may slow root absorption or where a quick response is needed. The spray should be contained, since some micronutrient sources can stain.

Dry fertilizers may be applied either in controlled release or quick-release form. High analysis fertilizers may be difficult to distribute evenly because of the small amount needed per container. Liquid applications of soluble or suspension fertilizers provide a uniform dosage and fast and easy distribution, but more frequent

applications may be needed because they may leach from container soil.

FERTILIZING INTERIOR PLANTS

During production, the foliage plant growth is quickened by using high rates of nutrients. These same plants grown indoors, however, usually get less light, and neither need nor will tolerate the amount of fertilizer they received in production.

Exact fertilizer needs are difficult to judge in interiorscape maintenance without measuring light at strategic locations.

Light varies from one side of a room to another, often within a few feet. Usually, the stronger the light under which foliage plants are growing, the more nutrients are needed. Recommended annual fertilizer rates can vary from as low as 0.3 grams of nitrogen per square foot for low light intensities to 3.0 grams for high intensities.

A complete fertilizer with a nitrogen/phosphorus/potassium ratio (similar to those recommended for landscape plants) is suitable for indoor plants. The highest levels of nutrients should be applied at highest growth periods, for most plants, spring and summer, when natural light is strongest.

Micronutrients are seldom recommended but may be needed when growing sensitive plants in media other than soil. The rubber plant (*Ficus elastica*) and the Areca palm (*Chrysalidocarpus lutescens*) are both sensitive to boron deficiency.

In addition, the Areca palm can also become zinc deficient. However, over-applications of micronutrients can cause toxicity problems.

Nutrient levels are determined by plant species. Plants normally grown under low levels of fertility include many ferns and fleshy plants such as Peperomia. Plants needing high nutrient levels include rapidly-growing species and large-leaf plants such as Ficus and Schefflera.

Salt, both from fertilizer and irrigation water, can build up unless the root area is periodically flushed with excess water which is allowed to drain away. This is true when plants are over-fertilized during periods of low light and/or little growth.

Since visual symptoms such as stem rot and leaf necrosis in new growth are similar to those of over-watering, the soil should be tested for soluble salts.

WT&T

What is so rare as a day in May? Especially with no leaf spot.



Leaf spot. It's a spoiler. Give it half a chance and it'll go after your grass.

To protect the beauty and playability of your turf from leaf spot, get new [®]DYRENE 4 Turf fungicide.

Now available as a flowable, it is not only easier and cleaner to use, but much more economical, too. For control of leaf spot, DYRENE 4 can be applied at *half the rate* of DYRENE wettable powder, yet will provide *twice the residual*.

DYRENE has always been an essential part of any complete program of disease control. Now, it's economical, too. Always read the label before use.



DYRENE.
IT STOPS THE SPOILER.



Mobay Chemical Corporation
Specialty Products Group
Box 4913, Kansas City, Mo. 64120

DYRENE is a Reg. TM of Mobay Chemical Corporation.

Introducing

From LESCO, the company that listens and responds to customer needs, comes a complete new line of small-particle sulfur-coated urea fertilizers.

Specifically designed for low-cut turf, the new ELITE fertilizers from LESCO provide many more particles per square foot than coarser-grade products that can cause speckling. That means consistent greening and a more uniform distribution pattern. In addition, the slow-release feeding provided by the sulfur-coated urea insures steady growth and long residual.



Shown is a half-gram each of granular sulfur-coated urea fertilizer (-6/+16) and LESCO ELITE sulfur-coated urea fertilizer.

The chance of mower pickup is virtually eliminated with ELITE fertilizers because the new small particles readily filter into the turf.



The chance of mower pickup is virtually eliminated with the new LESCO ELITE fertilizers.



Granular Sulfur-Coated Urea (-6/+16)

LESCO

ELITE

And because LESCO manufactures its new ELITE fertilizers using sulfate of potash, there's more plant-available sulfur and low potential for burn.

The fertilizer you've been waiting for is available now in six analyses: 21-0-16, 19-0-19, 15-0-30, 12-24-14, 21-4-12 and 31-0-0.




LESCO ELITE is available now in a variety of analyses.

To order the ELITE fertilizer that best fits your needs, call us toll free today.

(800) 321-5325
NATIONWIDE

(800) 362-7413
IN OHIO

Circle No. 251 on Reader Inquiry Card



LESCO ELITE Sulfur-Coated Urea

LESCO

LESCO, Inc., 20005 Lake Road, Rocky River, Ohio 44116 (216) 333-9250

THE DECIDING FACTOR

Water. Its availability may be the deciding factor when choosing a turf variety. Your choice should reflect water availability in your area.

by Dorothy F. Borland



It's national news every night. In newspapers across the country, the water shortage is reported. The potential repercussions of long-term shortages are highlighted.

Because of the shortage, methods are being explored to eliminate or ease the low water supply.

Two common methods are: (1) finding new water supplies through construction of water diversions or buying water rights; or (2) through restrictions on how available water may be used. Obtaining new water supplies is time consuming and quite expensive.

Regulating water use is an easier method. Unfortunately, too many municipalities prefer to legislate than educate on intelligent water use, especially in relation to water use in the landscape.

Of water used in urban areas, estimations indicate approximately 50 percent is used on the landscape. This usage is very visible.

In response to water shortages, some municipalities have limited the

Dorothy Borland is a turfgrass consultant with The Turf Expert in Denver.

Buffalograss provides a nice turf on this home lawn in the month of June.



The smooth brome/tall fescue mix at two maintenance levels. At left, irrigation every three days. At right, irrigation as needed, perhaps twice a month. Notice the difference in turf density.