

tions may be necessary to provide control over the entire germination period. Contact your local university or industry turf specialist for germination times of primary weeds.

### The differences

Turfgrass pre-emergence herbicides differ in soil longevity, weed control efficacy, and potential turfgrass injury. Benfen is generally considered to be shorter-lived in soils than bensulide or pendimethalin; while DCPA is considered to be intermediate. Concentration of the initial application is important in terms of maintaining pre-emergence herbicide soil concentrations.

Pre-emergence herbicides must maintain a critical soil residual level (threshold value) during the growing season that is conducive for germination of the target weed (i.e., annual bluegrass, crabgrass, foxtail, goosegrass or spurge). Applications made too early in the season may break down in the soil to levels below the threshold value. If this occurs and conditions remain favorable for weed germination, less than desirable control will be obtained.

### Maintaining levels


To obtain critical threshold values for the pre-emergence herbicide, adequate application rates must be made or repeat applications must be applied to maintain the threshold level. Initial applications will be dictated by label directions, turfgrass safety, efficacy and economics. In cases where potential turfgrass injury is a concern, light, frequent or split applications may be used to obtain control and minimize potential turfgrass injury.

Consult your local turfgrass specialist for specific information regarding the need for split applications of pre-emergence herbicides in your area. This information should increase its effectiveness and safety of your pre-emergence herbicide program.

Most turfgrass pre-emergence herbicides are effective in controlling crabgrass. These herbicides may differ somewhat in percent of control. It is wise to check with turf specialists in your region for the most efficacious materials.

### Other annuals

Pre-emergence herbicides differ in their ability to control other annual weed problems. For example, DCPA and pendimethalin effectively control prostrate spurge; while benfen and bensulide are ineffective in controlling this troublesome weed. Bensulide, at concentrations normally ap-



Pre-emergence herbicides differ in their ability to maintain effective soil residual concentrations (i.e., Herbicide A vs. Herbicide B). When values drop below the "threshold" level, effective weed control is lost. In this case, Herbicide B requires a second application to maintain effective weed control.

plied for crabgrass control, is not as effective in controlling foxtail as is DCPA or pendimethalin. Oxadiazon is more effective for goosegrass control than DCPA. Pendimethalin has good to excellent efficacy for all the annual warm-season species.

It is essential to apply pre-emergence herbicides uniformly to the area. These herbicides are bound in the soil by clay and organic matter and have limited lateral movement. Skips in application allow the target weed to escape, germinate and produce addi-

tional seed, thus disrupting turf quality. Careful application is important, making sure to follow label directions and to calibrate application equipment prior to treatment. This will give the safest, most efficacious control from the herbicide selected at the most economic rate.

Turfgrass managers should keep in mind that successful pre-emergence herbicide programs result from selecting the appropriate herbicide, and applying it uniformly at the appropriate rate and time. **WT&T**

## PRE-EMERGENCE HERBICIDES FOR WARM-SEASON TURF

by Tim R. Murphy

**S**uccessful weed control programs in warm-season turf depend on the development of a two-phase control strategy by the professional turf manager.


The first phase involves the use of cultural practices and insect and disease control programs that promote a dense, vigorous turf cover. Adequately maintained turf is less susceptible to a high level of a weed

infestation than poorly maintained turf.

Prior to the use of any herbicide, cultural practices—adequate fertilization, irrigation, cultivation and correct mowing height and interval—should be matched to the needs of a particular turfgrass species. Addition-

Tim Murphy is an extension agronomist in weed science for The University of Georgia.

# Ryan keeps your business growing strong.



Ryan equipment is a great addition to your business. Because with the Ryan line, you can do more jobs faster at less cost. There's less downtime and more worktime. And the more work you can do, the more business you'll take on.

With competitive equipment, what you save on the purchase price you'll pay in replacement costs. But Ryan is designed to go the distance. So when you're considering Ryan, consider what Ryan can do for you in the long run.

## The long-term value of the Lawnaire IV.®

The Lawnaire IV is the highest-quality, roll-type aerator on the market today. It combines speed and precision for fast and effective core aeration.

Self-propelled by a rugged 3-hp. industrial engine, the Lawnaire IV features a 38 lb. removable weight bar and a 55 lb. water drum, made of durable, corrosion-resistant polyethylene for tine penetration up to 2¾ inches. The 30 coring

tines penetrate the soil evenly every 3¾ inches across a 19-inch path. A convenient lift handle raises and lowers the machine for easy maneuverability in tight spots. A single clutch-throttle lever on the handlebar makes operation a snap.

Add Ryan's durable, low-maintenance, chain-drive design, and you've got an aerator that will keep you on the job and out of the repair shop for years to come.



**Increase your customer base with the new Mataway® Overseeder.**

Nothing is as versatile and effective for turf renovation as Ryan's new Mataway Overseeder. A powerful 10-hp. engine and extra wide 19-inch swath lets you dethatch or deep slice up to 10,000 square feet per hour. Snap out reels make it easy to change the type of blade and spacing you want. An Overseeder accessory can also be added, giving you the flexibility to power rake, overseed, or to do both at the

same time with a single piece of equipment.

It's a cost-effective way to stay one step ahead of your competition.

**A full line of Ryan reliability.**

The proven Ryan Ren-O-Thin® power rakes with interchangeable reels; the industry standard Jr. Sod Cutter with Tote Trailer; and the Tow Lawnaire for really large turf areas put you in a position for additional business. Because they're what you need to provide

professional quality turf care.

**Prove it to yourself.**

The best test of Ryan's reliability is in your own back yard. So ask the dealer nearest you for a free demonstration today. Or call toll-free: 1-800-228-4444.

**RYAN®**

**BUILT TO LAST**

6082 Cushman, OMC-Lincoln, P.O. Box 82409, Lincoln, NE 68501

ally, insect and disease problems should be eliminated. Strict reliance on herbicides without regard for the contribution of other management practices to the total weed control program does not result in a high-quality, aesthetically appealing turf.

The second phase of a weed control strategy involves the use of pre-emergence and post-emergence herbicides. Pre-emergence herbicides are applied prior to weed seed germination and generally have little, if any, activity on emerged weeds. In contrast, post-emergence herbicides are applied directly to the foliage of emerged weeds and have no or minimal residual activity on non-emerged weeds. Although the majority of herbicides may be classified as either pre-emergence or post-emergence, atrazine and simazine are exceptions. These herbicides have both pre-emergence and post-emergence activity on a wide variety of winter annual weeds.

Pre-emergence herbicides form the base of the chemical weed control program and are used primarily to control annual grasses and certain annual broadleaf weeds. This group of weeds can be controlled with post-emergence herbicides; however, two or more applications are usually necessary and a possibility exists that the turf may be discolored or injured for a short time after application.

Generally, pre-emergence herbicides are safer to apply to warm-season turf, and often only one application is necessary. Therefore, pre-emergence herbicides are the preferred method for controlling annual grass and certain annual broadleaf weeds. Post-emergence herbicides should be relied upon to control perennial grass and broadleaf weeds that are not controlled by pre-emergence herbicides.

Pre-emergence herbicides are applied in the spring for crabgrass and goosegrass control and in the fall months primarily for annual bluegrass control.

Since most pre-emergence herbicides are not effective against emerged weeds, applications must be made prior to weed seed germination. Late February to early March applications generally provide better crabgrass control than later applications. However, in the cooler, mountainous regions of the South, the spring application may be delayed until late March.

For annual bluegrass, late August to early October applications are used, depending on the geographical location. Pre-emergence herbicides require rainfall or irrigation water to

**Table 1. Directory of turfgrass pre-emergence herbicides.**

Common Name	Trade Name(s)	Formulations <sup>1</sup>	Company
Atrazine	Aatrex	80W, 90DG, 4L	Ciba-Geigy
	Purge	4L	Security
Benefin	Balan	2.5G	Elanco
	2.5 Benefin Granular	2.5G	Lesco
Benefin + Oryzalin XL		2G	Elanco
Benefin + Trifluralin Team		2G	Elanco
Bensulide	Betasan	2.9E, 4E, 3.6G, 7G, 12.5G	Stauffer
	Lescosan	4E, 7G, 12.5G	Lesco
	Weedgrass Preventer	8.5G	Scotts
	Pre-San	4E, 7G, 12.5G	Mallinckrodt
Bensulide + Oxadiazon	Bensumec 4LF	4E	PBI Gordon
	Goosegrass/Crabgrass Control	5.25G + 1.31G	Scotts
DCPA	Dacthal	75W	Fermenta
	Dacthal	5G	Lesco
Ethofumesate	Prograss	1.5E	Nor-Am
Fenarimol	Rubigan	50W	Elanco
Napropamide	Devrinol	50W, 5G	Stauffer
Oryzalin	Surflan	4AS	Elanco
Oxadiazon	Ronstar	50W, 2G	Rhone-Poulenc
Pendimethalin	Southern Weedgrass Control	2.45G	Scotts
	Turf Weedgrass Control	1.71G	Scotts
	Weedgrass Control	60DG	Scotts
	Pre-M	60DG	Lesco
	Pendimethalin	60DG	Clean Crop
Pronamide	Kerb	50W	Rohm-Haas
Siduron	Tupersan	50W	Du Pont
Simazine	Princep	80W, 90DG, 4L, 4G	Ciba-Geigy

<sup>1</sup>W = Wettable powder; DG = Water dispersible granule; L = Liquid; E = Emulsifiable concentrate; G = Granule.

**Table 2. Tolerance of warm-season turfgrasses and tall fescue to pre-emergence herbicides.**

Herbicide	Turfgrasses					
	Bahia	Bermuda	Centipede	St. Augustine	Tall Fescue	Zoysia
Atrazine	NR	I	T	T	NR	I
Benefin	T	T	T	T	T	T
Benefin + Oryzalin	T	T	T	T	T	T
Benefin + Trifluralin	T	T	T	T	T	T
Bensulide	T	T	T	T	T	T
Bensulide + Oxadiazon	NR	T	NR	NR	T	T
DCPA	T	T	T	T	T	T
Ethofumesate	NR	I	NR	NR	NR	NR
Napropamide	T	T	T	T	T	NR
Oryzalin	T	T	T	T	T	T
Oxadiazon	NR	T	NR	T	T	T
Pendimethalin	T	T	T	T	T	T
Pronamide	NR	T	NR	NR	NR	NR
Siduron	NR	NR	NR	NR	T	NR
Simazine	NR	T	T	T	NR	T

T = Tolerant; I = Intermediate tolerance, apply to dormant grass; NR = Not Registered for use.

move them into the zone of maximum weed seed germination (e.g. the upper one to two inches of the soil profile). Recommendations vary slightly among the different pre-emergence herbicides, but unless one-fourth to one-half inch of rainfall occurs within seven days, the herbicide should be irrigated in the top two inches of the soil profile.

Removal of heavy thatch by

cultivation (aerification, verticutting, etc.) prior to herbicide application will improve spray penetration through the turf canopy and increase herbicide contact with the soil. Cultivation after a pre-emergence herbicide application is not recommended since it may decrease the effectiveness of the herbicide.

In past years, there were only about four or five pre-emergence her-

There's no better value for crabgrass control than Balan. And none with more experience. It's been used more years by more golf course superintendents than any other granular preemergence herbicide.

Its easy-to-spread clay carrier is one reason why. It makes application more uniform, reduces the dust common in other carriers and also increases your equipment calibration accuracy.

With Balan, you get superior preemergence crabgrass control for eight to ten weeks with

a single application. Then you can overseed or perform other turf management jobs, without worry of reduced seed germination. If you want longer control, the Balan price allows you a second application.

Balan. The better spreader. Talk to your Elanco distributor soon. Or write:

Elanco Products Company  
A Division of Eli Lilly and Company  
Lilly Corporate Center, Dept. E-455, Indianapolis, IN 46285, U.S.A.  
Balan®—(benefin, Elanco Products Company)

Circle No. 122 on Reader Inquiry Card



## Proven Balan.<sup>®</sup> The better spreader for crabgrass control.



bicides available for use on warm-season turf. However, 16 are now registered for use on Southern turf (Table 1).

The selection of which to use should be based on (a) turfgrass tolerance and (b) the weed species composition of a particular site. Herbicide selection based solely on cost may result in possible turf injury and/or the additional expense of a follow-up post-emergence herbicide treatment.

### Turfgrass tolerance

Warm-season turfgrasses and tall fescue vary in their tolerance to pre-emergence herbicides (Table 2). For example, with the exception of spring greenup, centipede and St. Augustine have excellent tolerance to atrazine at all times of the year. Zoysia and bermudagrass are tolerant to atrazine when dormant, but may be discolored or injured if applications are made at other times of the year. Tall fescue is highly susceptible to severe injury from atrazine. Also, bahiagrass will be injured from pre-emergence applications of this herbicide.

### Summer annual weed control

Crabgrass (large, smooth, Southern) and goosegrass are commonly found in Southern turf. With the exception of atrazine, pre-emergence herbicides applied in the spring months provide good to excellent crabgrass control (Table 3).

Goosegrass tends to sporadically germinate during the long growing season of the South and is more difficult to control than crabgrass. Oxadiazon, napropamide and bensulide + oxadiazon have provided high levels of goosegrass control in tests conducted in Georgia. Benfen + oryzalin, oryzalin and the various formulations of pendimethalin have also provided fair to good control of goosegrass with a single application at recommended rates.

Split applications of oryzalin and pendimethalin are being evaluated in Georgia and usually have provided better goosegrass control than a single application.

Sandbur is an occasional problem in warm-season turf. It can be controlled with spring applications of benfen + oryzalin, bensulide, napropamide, oryzalin, and pendimethalin.

Certain pre-emergence herbicides can also be selected that provide control of prostrate spurge and prostrate knotweed.

### Winter annual weed control

Similar to summer annual weed control, pre-emergence herbicides vary

in their effectiveness on winter annual weeds. Bensulide has generally not provided the high level of annual bluegrass control that has been observed with other pre-emergence herbicides (Table 4).

Common chickweed and henbit are easily controlled by pre-emergence applications of benfen, benfen + oryzalin, oryzalin, or pendimethalin. Additionally, common chickweed can be controlled with DCPA, ethofumesate, napropamide, and pronamide. Herbicides that have effectively controlled parsley-piert are bensulide and oxadiazon. Napropamide is effective for spurweed control. Good to excellent corn

speedwell control can be obtained with benfen, DCPA, napropamide, oxadiazon and pronamide. Atrazine and simazine provide effective pre-emergence or post-emergence annual bluegrass and winter annual broadleaf weed control (Table 4).

The wide time period of application for these herbicides offers tremendous scheduling flexibility for the professional warm-season turf manager.

### Overseeded bermudagrass

On bermudagrass that is overseeded with annual or perennial ryegrass, fenarimol or ethofumesate may be used for annual bluegrass con-

Table 3. Summer annual weed control ratings for pre-emergence herbicides.

Herbicide	Crabgrass spp.	Goosegrass	Sandbur	Prostrate	Prostrate
				Spurge	Knotweed
Atrazine	P	P	P	E	E
Benfen	E	F	F	P	P
Benfen + Oryzalin	E	F-G	G	L	L
Benfen + Trifluralin	L	L	—	—	—
Bensulide	E	P	G	P	G
Bensulide + Oxadiazon	E	G	—	—	—
DCPA	E	F	F	P	G
Napropamide	E	G	G	—	L
Oryzalin	E	F-G	G	L	L
Oxadiazon	G	E	F	P	G
Pendimethalin	E	F-G	G	L	—
Siduron	G	P	—	P	P
Simazine	G	P	—	G	G

E = Excellent,  $\geq 90\%$  control.

G = Good, 80 to 89% control.

F = Fair, 70 to 79% control.

P = Poor, <70% control.

L = Weed species is listed on the herbicide label, but has not been evaluated by the University of Georgia.

— = Weed response is not known.

Table 4. Winter annual weed control ratings for pre-emergence herbicides.

Herbicide	Annual	Common	Parsley-		Corn	
	Bluegrass	Chickweed	Henbit	Piert	Spurweed	Speedwell
Atrazine	E	E	E	E	E	E
Benfen	E	G	G	P	P	E
Benfen + Oryzalin	E	L	L	—	—	—
Benfen + Trifluralin	L	—	—	—	—	—
Bensulide	F	P	P	E	P	P
DCPA	G	E	F	P	P	E
Ethofumesate	G	L	—	—	—	—
Fenarimol	G	—	—	—	—	—
Napropamide	G	E	P	P	E	E
Oryzalin	G	L	L	—	—	—
Oxadiazon	G	P	P	E	P	G
Pronamide	E	E	P	P	P	E
Pendimethalin	G	L	L	—	—	—
Simazine	E	E	E	G	E	E

E = Excellent,  $\geq 90\%$  control.

G = Good, 80 to 89% control.

F = Fair, 70 to 79% control.

P = Poor, <70% control.

L = Weed species is listed on the herbicide label, but has not been evaluated by the University of Georgia.

— = Weed response is not known.

# MUSTANG

## THE TALL FESCUE THAT LOOKS LIKE BLUEGRASS

KENTUCKY BLUEGRASS

MUSTANG TURF-TYPE TALL FESCUE

### BLUEGRASS QUALITIES WITH TALL FESCUE PRACTICALITY

You'll quickly notice Mustang's finer texture, rich dark green color and dense, uniform turf—and you'll understand why we say Mustang has bluegrass-like qualities.

But there's more to Mustang turf-type tall fescue than beauty; it's tough and durable. It's heat and drought tolerance, winter hardiness, and ability to endure low mowing heights are remarkable. Mustang even shows improved resistance to *Helminthosporium* net blotch and many other diseases.

Best of all, Mustang is practical, because it performs

extremely well under low maintenance conditions like minimum fertilization, watering and mowing. National tests and actual applications in parks, golf courses and playing fields have proven it.



**DROUGHT TOLERANCE AFTER SEVEN WEEKS NO RAINFALL.**



KENTUCKY BLUEGRASS



K-31 TALL FESCUE

That's why Mustang is becoming the favorite choice of professional turf managers, over K-31 and many other commercially available varieties.

For bluegrass qualities and tall fescue practicality, use the professional's choice—Mustang turf-type tall fescue.

Pickseed also produces



and



and other fine turf grasses available nationwide from quality seed suppliers.

Produced by:

**PICKSEED®**  
**PICKSEED WEST Inc.**

P.O. Box 888 • Tangent, Oregon 97389

(503) 926-8886


# 10 reasons why you should ask for a Continental R-Series engine

High performance, long life design — that's what the Continental R-Series liquid-cooled engines from Teledyne Total Power can offer you.

Take a look at the counter-weighted spheroidal cast iron crankshaft with five main bearings and forged steel connecting rods... right down to the cast iron cylinder block and crankcase, you can depend on the quality to keep up with your

equipment.

Ask for Continental R-Series engines from Teledyne Total Power, with over 5,000 distributors and service centers in over 90 countries ready to serve you. Send for your free "Reasons Why" brochure today: Teledyne Total Power, P.O. Box 181160, Memphis, TN 38181-1160, 901/365-3600, Telex: 462-1058 (ITT)



**TELEDYNE  
TOTAL POWER**

3409 Democrat Road • Memphis, TN 38118  
Call toll-free: 1-800-932-2858

R-11/14

tol. Ethofumesate should only be applied to bermudagrass that is completely dormant. Ethofumesate applications to bermudagrass that is not completely dormant have shown to delay greenup the following spring.

Spring applications of oryzalin, oryzalin + benefin and pendimethalin have shown to decrease the stand of overseeded ryegrass. In some years, overseeded ryegrass may persist for longer than the desired time period and provide competition to bermudagrass that is breaking dormancy. These herbicides are currently being evaluated for their possible role in improving the transition of a cool-season to a warm-season grass in overseeded golf course fairways.

## Summary

Pre-emergence herbicides represent the backbone of the chemical weed control program. Post-emergence herbicide use will be necessary if perennial grass or perennial broadleaf weeds infest the site. However, pre-emergence herbicides offer the professional turf manager more than crabgrass, goosegrass and annual bluegrass control. Careful study of weed response charts or the herbicide label will enable the selection of a pre-emergence herbicide to control these annual grasses and certain annual broadleaf weeds.

WT&T

### PRE-EMERGENCE HERBICIDE SOIL LONGEVITY

Short

benefin  
DCPA  
oxadiazon  
pendimethalin  
bensulide

Long

### PROSTRATE AND SPOTTED SPURGE CONTROL

Poor

benefin  
bensulide

Variable

oxadiazon  
siduron

Excellent

DCPA  
pendimethalin

### CRABGRASS CONTROL


Good

benefin  
bensulide  
DCPA  
oxadiazon  
pendimethalin

### FACTORS INFLUENCING PESTICIDE CONCENTRATIONS IN SOIL

- \*photodecomposition
- \*volatilization
- \*leaching
- \*microbial breakdown
- \*insoluble complexes
- \*absorption
- \*dilution
- \*mechanical removal





CHIPCO® 26019  
HAS BEEN  
CONTROLLING  
DISEASES  
FOR A LONG TIME,  
FOR A LONG TIME.

---

Years of excellent, long-lasting performance have made CHIPCO® 26019 fungicide the leader in its field.

---

When it's your business to keep turf healthy and handsome, you don't wait for disease to break out. You strike first and play for keeps.

That's why thousands of superintendents base their disease control programs on CHIPCO 26019 fungicide.

It's unsurpassed at controlling the major turf diseases, like Helminthosporium Leaf Spot and Melting Out, Dollar Spot, Brown Patch, Fusarium Blight, Red Thread, Fusarium Patch and Gray and Pink Snow Molds.

And it stays on the job longer, giving up to 28 days of protection for a lower cost per day.

Add a few extras — no phytotoxicity, low toxicity, pesticide compatibility, and no corrosiveness — and you've got a fungicide that's earned its place as the standard of the industry.

Use CHIPCO 26019 this season and you'll stay with it for a long time to come.

Rhône-Poulenc Inc., Agrochemical Division, Monmouth Junction, NJ 08852. 

**CHIPCO**  
**26019**  
Taking care of business.

CHIPCO® is a registered trademark of Rhône-Poulenc Inc.

Circle No. 166 on Reader Inquiry Card

# THE LANDSCAPE EXPO: YEAR TWO

The Landscape Expo has seen exceptional growth in just one year, making it one of the largest industry trade shows.

**M**ore than 5,000 people will visit the 1987 edition of the Landscape Exposition, March 3-5 in the O'Hare Exposition Center, Rosemont, Ill.

The 5,000 figure will double the attendees of last year's inaugural Expo, according to Expo administrators. More than 225 exhibiting companies will cover more than 50,000 sq. ft. of exhibit space, a 20 percent increase from the first show.

The Expo has the backing of the PGMS, PLCAA, ALCA and the Illinois Landscape Contractors Association.

Supporting the trade show will be a greatly expanded schedule of educational sessions (see following page) featuring some of the green industry's leading experts.

Fernando Bensusaski, managing partner in Bensusaski, Delana & Luce, kicks off the educational program with a speech about buying and leasing options for new equipment. A second talk by Bensusaski will deal with acquiring and keeping capital.

Consultants Ed Wandtke and Rudd McGary of All-Green Management Associates will discuss the logistics of setting up an employee training program, as well as what the program should include. In other sessions, the two will address business aspects of the industry such as employee relations, marketing and pricing.

Jim Leatzow, president of Leatzow Agency/Financial Guardian, and Richard Lehr, a partner with Sirote, Permutt, et. al., will deliver a pair of

two-part programs, one on preventing problems incurred by parks and playgrounds, and another on avoiding lawsuits.

In a related speech, *LAWN CARE INDUSTRY* magazine editor Elliot Maras will deliver a talk on avoiding conflict with the media. With recent problems the lawn care market has been facing with the media, this session comes at the perfect time.

Industry consultant Charles Vanderkooi will give a three-part program based on his book *Estimating & Management of Principles for Landscape Contractors*.

Dr. Martin Petrovic of Cornell University will present a session on effective turf management, and how it can help preserve groundwater supplies.

