

# Trade In The Tools Of Your Trade!

Trade in *any* broken or used cutter, trimmer, blower, or sprayer for a Maruyama product and get factory cash back for your trade!

You can depend on the uncompromising quality of Maruyama commercial equipment. It's built for long life and backed with parts & service by local qualified dealers. We make long life products. When you're looking for the very best, look for Maruyama.

**Call your local dealer today**  
**Trade in offer ends soon.**

## Maruyama's total line includes

- BRUSHCUTTERS
- BLOWERS
- HEDGE TRIMMERS
- MIST DUSTERS
- POWER SPRAYERS
- GRANULE SPREADERS
- SOIL INJECTERS
- MANUAL SPRAYERS
- FRAME MOUNT SPRAYERS
- WATER PUMPS

### U.S. Office

MARUYAMA U.S., Inc.  
15436 N.E. 95th St., Redmond, WA 98052  
P.O. Box 2167, Redmond, WA 98073  
Phone: (206) 885-0811  
Fax: (206) 885-0123

### Singapore

MARUYAMA MFG. CO., Inc.  
16 Raffles Quay  
Level 36, Hong Leong Bldg., Singapore 0104  
Phone: 321-8943 Fax: 225-9060  
Telex: RS 43382 COHQHL



Quality equipment since 1895




### Paris

MARUYAMA MFG. CO., Inc.  
34, BD. Haussmann, 75009 Paris, France  
Phone: (1) 47 70 28 52  
Fax: 011-331-45231830  
Telex: MBI 290266F

### International Headquarters:

MARUYAMA MFG. CO., Inc.  
4-15 Uchi-Kanda 3-Chome  
Chiyoda-Ku, Tokyo, 101 Japan



© 1990 CIBA-GEIGY Corporation, Turf and Ornamental Products, Box 18300, Greensboro, NC 27419. Always read and follow label directions.

Circle 144 on Reader Service Card



TO THOSE

STILL

QUESTIONING

THE BENEFIT

OF PACE

FUNGICIDE,

WE'D LIKE TO

CLEAR

THE AIR.

We'd also like to help clear up that nasty container disposal problem of yours.

Not to mention improving your Pythium control a healthy notch or two.

All of which is possible thanks to Pace® and its dust-free, water-soluble package.

It's simply the most effective Pythium control you can use. And the most responsible way you can use it.



# COOL-SEASON INSECT CONTROL GUIDE

New application methods and materials will soon result in more effective insect control. Our annual guide helps you time your treatments.

by Harry Niemczyk, Ph.D., Ohio State University

**D**uring the past year, the Turfgrass Pesticide Laboratory at the Ohio Agricultural Research and Development Center, Ohio State University, Wooster, Ohio, conducted research that further demonstrated the benefits of turfgrass to

our environment.

Researchers found that insecticides applied to turfgrasses with or without thatch showed little or no potential for leaching into the underlying soil.

The studies, based on 1988 and

1989 field applications of the pre-emergent herbicides pendimethalin, bensulide, benefin, trifluralin and oxadiazon, has shown no residues at the 10-inch soil level.

Excitement is also brewing about methods of placing insect control ma-

## SPRING

(April-May)



**Chinch bugs and billbugs**— As the warm days of spring approach, movement of chinch bug and adult billbugs increases rapidly. Generally, egg laying begins the first week of May, but can begin in mid-April if spring arrives early. Occasionally adult billbugs can be seen on sidewalks on warm April afternoons.

Generally, application of insecticides to prevent infestations of chinch bugs and billbugs should be completed by the first week in May. Applications may begin as early as the last week of March. Such applications must be made before significant numbers of eggs are laid. This time may vary as much as a week or more, depending on spring weather.

**Grubs**— Overwintered grubs return to the surface and begin feeding on turfgrass roots in April. Increased activity and damage from birds, moles, skunks and raccoons foraging on grubs can also be expected and continues through May.

Treatment should be delayed until the grubs are in the top one inch of soil. Irrigation or rainfall should follow such applications to aid in moving the insecticides to the target grub as soon as possible following application.

Although milky disease products for control of Japanese beetle grubs may be applied any time there is no frost in the soil, spring is a good time for such applications. The soil is open and frequent rains move the disease spores into the soil and thatch.

Milky disease products are primarily effective against Japanese beetle larvae. Ineffectiveness against other species is low.

Incidents of large grub infestations (June bug larvae, for example) have been increasing in cool-season areas. Locations of such infestations should be identified because reinfestation is likely every three years.

Eggs are laid in May and June, therefore treatment should be made in late summer, early fall of that year or the next spring while larvae are small. Later applications against full-grown larvae have given inadequate control in past studies.

**Black turfgrass ateniensis**— Adults of the black turfgrass ateniensis can be seen "at wing" in April and are often found in clipping catchers after early mowing of golf course greens. These adults begin laying eggs in early May, or about the time *Vanhoutte spirea* first comes into bloom. Dursban applied to a fairway at this time kills adults and prevents summer infestation of larvae. Check with local extension entomologists for the precise time if needed.

A second application, two weeks after the first, may be needed to successfully prevent infestation.

**Sod webworms**— Overwintered larvae of the sod webworm begin feeding as soon as the grass begins to grow. Usually damage is insignificant but areas that do not green up may be infested. These areas frequently have probe holes from starlings feeding on larvae.

Moth flights begin in May in northern areas. Young larvae are usually present about two weeks after the spring moth flight peaks, so treatment of young larvae can be done in May in some areas.

**Cutworms**— Moths of cutworms begin laying eggs



# **WE'D LIKE TO TELL YOU ABOUT THE BIRDS AND THE BEES.**



We'd also like to tell you about pets and other animals, and hundreds of species of ornamental plants.

Because they are the biggest reasons to use Mavrik Aquaflow.<sup>®</sup>

Mavrik<sup>®</sup> is water-based, effective at very low rates, and gentle on beneficial species. It is a non-restricted product and can even be sent through the mail!

Mavrik won't harm plants either, not even when sprayed on open blooms.

Yet Mavrik has excellent broad spectrum efficacy against most insect pests as well as mites.

So switch to Mavrik. Then if your customers are concerned about the materials you use, you can tell them about the birds and the bees too.




**SANDOZ**  
CROP PROTECTION  
CORPORATION  
DES PLAINES, IL 60018

Use pesticides effectively. Read and follow label directions carefully.

© 1989 Sandoz Crop Protection Corporation  
Mavrik and Mavrik Aquaflow are registered trademarks of Sandoz, Ltd.





Japanese beetle larvae at thatch-soil interface. This year's early spring will bring grub to the surface earlier. Watch for skunks and raccoons, which can damage turf when searching for them.

## SUMMER

(June-July)



**Chinch bugs**— Chinch bug eggs continue to hatch into June. Bright red nymphs with a center white band appear. The number of chinch bugs increases rapidly in June. Their populations peak in July and August, when northern lawns can receive severe damage. This damage is often masked by summer dormancy of turf caused by drought. Hot, dry conditions are ideal for chinch bugs.

During August the nymphs molt into adults that mate and lay eggs, thus producing a second generation. Some northern areas have only one generation per year.

**Billbugs**— Bluegrass billbug larvae feed on grass stems during June and move to the plant crowns, roots and rhizomes during July. This feeding pattern causes brown spots that frequently resemble the symptoms of some fungus diseases. Symptoms are also often masked when the turf is dormant from drought.

The larvae usually move deeper into the soil under dry soil conditions. During late July and August the larvae burrow deeper into the soil to pupate and transform into adults.

**Grubs**— By June, grubs have stopped feeding and are in the pupal stage, three to four inches in the soil. Beginning mid-June and continuing through mid-July, the adults of various species emerge and burrow into the soil to lay eggs. Hatching and appearance of young larvae occur during July and August.

Extreme heat and drought during the summer may cause some grubs to move deeper in the soil. Under such conditions, irrigation several hours before treatment and a thorough soaking afterward is advisable.

**Black turfgrass ataenius**— Eggs laid by beetles during May hatch in June and the larvae immediately begin feeding on turf roots and thatch.

From late June to mid-July, symptoms of injury include wilting in spite of irrigation. In July, larvae move deep into the soil, pupate and emerge as adults. These adults lay eggs during August, producing a second generation in some states. The second generation larvae are capable of damaging turf. States farther north have only one generation.

**Sod webworms**— Damage from sod webworm larvae occurs occasionally in most of the cool-season turf region. Injury is more common in Midwestern states, usually in July and August. Older sod fields or areas with heavy thatch are good candidates for infestation. There are generally one or two generations per year, depending on the species.

**Cutworms**— Cutworm larvae continue to cause damage to golf course greens from June through August. These larvae pupate in the soil or thatch and emerge as moths that lay eggs for additional generations.

**Fall armyworm**— The fall armyworm is seldom a problem of cool-season turf.

**Greenbug**— Damaging populations of greenbugs can occur from June through August. Populations and incidents of damage frequently vary from area to area, even within the same city.

Symptoms of injury include turf under the dripline of trees and in open areas having a burnt orange color. When symptoms are apparent, numerous aphids (40 or more) may be seen on a single grass blade. Close examination of damaged turf is necessary because the aphids are small. If left untreated, a heavy infestation can kill the turf.

## SPRING from page 44

on golf course greens and other turf areas in the spring. These eggs hatch, producing larvae that feed on grass blades during the night. The black cutworm is the most common species on cool-season turf.

While visible damage is uncommon on home lawns, damage can be significant on golf course greens in late May.

**Greenbug**— Greenbug eggs begin hatching as early as April, but significant infestations do not develop until later in the year.

Greenbugs are also brought into the region from the south on upper air winds. Aphid numbers are too low to detect in lawns at this time.

**Winter grain mite**— Damage from this mite is often first noted in March or April when turf areas are receiving spring fertilizer applications. Winter grain mites are identifiable by eight bright red legs and a dark body.

By late May, the mites will have laid their eggs and died. Mites do not appear again until the eggs hatch in October.

**Clover mites**— Incidents of visible damage to home lawns are often seen in April in several Ohio cities and Denver. Usually a nuisance pest in and around homes, the clover mite occurs in large numbers (5,000 per square foot) across entire lawns and on turf next to building foundations. Symptoms of injury were the same as the winter grain mite. Turf next to foundations may be killed.

The clover mite has a slightly pink body and eight pale-colored legs. The first pair of legs is extremely long and protrude well out in front of the mite. The absence of bright red legs distinguishes the clover mite from the winter grain mite.

"I was hesitant to try it at first. I thought, surely, there would be some injury to my flowers. But Surflan convinced me otherwise. It didn't harm them at all when we sprayed directly over the top.

"Surflan wasn't so gentle to the weeds. We used to have a real problem with crabgrass. We'd have to go in and pull them all by hand. It was so time-consuming. I figured I was paying close to \$50 for each thousand square feet just for labor.

"But with Surflan, the job took only

minutes. And cost me less than \$3.00 for what used to cost me almost \$50.

"Surflan also took care of our oxalis and chickweed. And many other problem weeds, too. For almost eight months."

Prove Surflan for yourself. See your Elanco distributor and ask for a free Surflan brochure. Or call toll-free: **1-800-352-6776.**

Elanco Products Company  
A Division of Eli Lilly and Company  
Lilly Corporate Center  
Dept. EM-455, Indianapolis, IN 46285, U.S.A.  
Surflan®—(oryzalin, Elanco)



Circle No. 112 on Reader Inquiry Card

## **"Are you kidding? Spray herbicide over the top of my flowers?"**

**Surflan proved I could."**

Cynthia Harper  
Color Burst, Atlanta, Ga.

Ritz-Carlton, Buckhead  
Atlanta, Georgia



terials, such as insecticides or nematicides, biological control agents, entomogenous nematodes, insect growth regulators (IGRs), milky disease and others, right where the pest is, without leaving residues on the surface. This approach will re-

duce the application rate necessary for control since placement is directly in the zone of pest habitation. Another significant advantage of this method is that the potential for pesticide runoff will be significantly reduced, if not eliminated.

The equipment needed for subsurface placement of granular products on golf courses is already available and will be tested in 1990. Equipment now used for subsurface placement of liquid polymers has adapted for application of liquid

## FALL (Sept.-Oct.)



**Chinch bugs**—In the northern U.S. the second generation of chinch bug is at peak numbers in September. Nymphs complete their development to adults in late October. Most chinch bugs overwinter in the turf, but some move to protected areas before winter.

Generally, infestation levels at this time are not high enough to warrant the use of insecticides. Early fall rains and infection by a parasitic fungus (*Beauveria* spp.) usually provide sufficient control.

**Billbugs**—During September, billbug adults that developed from summer larvae are often seen on sidewalks, driveways, or other paved surfaces. Before winter, these adults seek shelter in thatch, along sidewalk edges or near foundations and overwinter. Many, if not most, overwinter in turf.

In some areas a partial second generation may occur. Larvae of this generation have been known to cause

visible damage in September and October.

**Grubs**—Most species of grubs are in the third of their three stages of development and are feeding actively. When soil temperatures decrease in late October, the larvae burrow deeper into the soil to overwinter. If soil temperatures remain warm, larvae stay at the surface and continue feeding. Severely cold winters have little effect on survival.

**Black turfgrass ataenius**—By September, adults of the current generation begin to fly into protected areas, such as golf course roughs, to overwinter. Larvae that have not completed development to adults before the first frost are killed.

**Sod webworm**—Northern sod webworm larvae are small and cause little if any damage in the fall. Late in the fall the larvae construct a cocoon-like shelter in which they overwinter.

**Greenbug**—Severe infestations of greenbug have been known to occur as late as the first week of December. Areas having a history of infestation should be re-examined when mild temperatures extend late into fall. Heavily-infested turf will not survive through winter.



Japanese beetle larvae infected with milky disease (top), and a healthy grub. Subsurface placement of milky disease products should significantly increase their effectiveness (Photo courtesy Dr. M. Klein, USDA-ARS).

## LATE WINTER (Nov.-Dec.)



**Chinch bugs and billbugs**—In northern zones, chinch bugs and billbugs overwinter as adults in thatch or sheltered sites near buildings. They can become active during warm days in March. Infestations of hairy chinch bugs and bluegrass billbugs also occur in zoysia, Kentucky bluegrass and fine fescues.

**Grubs**—The larvae of this group of pests normally overwinter six inches or deeper in the soil. If spring comes early, grub activity can be expected along with skunks and raccoons, which tear up the turf in search of them. Moles, which feed on grubs and earthworms, also become active at this time.

**Black turfgrass ataenius**—This golf course pest overwinters as an adult in the soil under debris in roughs or other protected areas. A few may be seen flying about on warm afternoons in early March. Usually this activity begins when crocus starts blooming and intensifies as the bloom of red bud appears.

**Greenbug**—The only stage of the greenbug known to overwinter in northern states is the egg. Shiny black eggs deposited the previous fall may be found adhering to grass blades, fallen tree leaves or other debris.

**Sod webworms**—The most common sod webworm species overwinter as larvae in the thatch or upper inch of soil. Feeding does not resume until hibernation (diapause) is broken by early spring warmth. LM



# For Poa Annua Control, Betasan® Easily Outscores The Competition.

In fact, the performance of Betasan® in controlling germinating Poa Annua so outdistances other preemergent herbicides, there almost is no competition. The figures shown are the average percent of control as reported in a three-year university\* study. The results are clear.

But almost total Poa Annua control isn't the only reason Betasan is your best herbicide buy. Betasan can be used safely on more turfgrasses than all other competitors. Also, Betasan is the only preemergent herbicide safe and labeled for use on sensitive species such as bentgrass and fescues, as well as other major golf course grasses.

Most turf professionals already know that Betasan is unbeatable against crabgrass. And now you know even more reasons why the long-lasting protection of Betasan is such a dominating force on any course.

For information, ask your dealer or ICI representative, or call the ICI Information Line at 1-800-759-2500.



**Professional Products**

	380	440	395	224	550	437	208	390	600	3624
BLUE	349	414	367	192	512	398	180	355	563	3330
WHITE	1	2	3	4	5	6	7	8	9	3042
HOLE	4	4	4	3	5	4	3	4	5	36
PAR	<b>90% CONTROL</b>									
<b>BETASAN®</b>	<b>54% CONTROL</b>									
<b>BALAN®</b>	<b>40% CONTROL</b>									
<b>DACTHAL®</b>	<b>36% CONTROL</b>									
<b>RONSTAR®</b>										
<small>Betasan is a trademark of ICI Americas Inc.                      Balan is a trademark of Eanco Products Company.                      Dacthal is a trademark of Fermenta.                      Ronstar is a trademark of Rhone-Poolex, Inc.</small>										



**Betasan®**  
 Selective Herbicide  
**No. 1 Against Poa Annua**

\*Rhode Island University test. Details available upon request.

Circle No. 251 on Reader Inquiry Card

and biological control materials. More walk-behind units for both liquid and granular products are being manufactured or are on the engineer's drawing board.

At OARDC, excitement is mounting about the materials we have to

evaluate in 1990.

Being able to track a pest's life cycle is still at least as important as selecting an insecticide for their control. This guide points out the seasonal occurrence of some important cool-season pests to be on the

lookout for this year. Best control will come when you apply insecticides during the pest's most vulnerable stages, which are outlined here. No endorsement of products is intended, nor is criticism implied of those not mentioned.

## HOW'S YOUR KNOWLEDGE OF TURF INSECTS?

### Test Yourself or Your Crew

1. Bluegrass billbugs overwinter as \_\_\_\_\_.

adults pupae  
larvae eggs

2. A pesticide with an LD<sub>50</sub> of 75 is safer than one with an LD<sub>50</sub> of 1000.

\_\_\_\_\_ True  
\_\_\_\_\_ False

3. Sod webworms generally overwinter as \_\_\_\_\_.

adults pupae  
larvae eggs

4. \_\_\_\_\_ percent of all insects are considered pests.

Less than 1% 5%  
1% 10%

5. The reason for poor grub control with pesticides is \_\_\_\_\_.

resistance  
too much rain  
insecticide did not reach pest  
UV breakdown

6. The Japanese beetle overwinters as \_\_\_\_\_.

egg pupae  
larvae adults

7. Grubs eat \_\_\_\_\_.

thatch soil  
grass roots thatch, soil and roots

8. Dursban is an \_\_\_\_\_.

organophosphate pyrethroid  
carbamate chlorinated hydrocarbon

9. Masked chafer adults fly \_\_\_\_\_.

during the daytime at mid-day  
early morning at night

10. Match these raster drawings with the following insects:

- a. Black Turfgrass Ataenius
- b. European Chafer
- c. Masked Chafer
- d. May-June Beetle
- e. Japanese Beetle



11. \_\_\_\_\_ 12. \_\_\_\_\_ 13. \_\_\_\_\_



14. \_\_\_\_\_ 15. \_\_\_\_\_

If you got:

- 14 right - SHARP!
- 13 right - WELL DONE!
- 12 right - not bad
- 11 right - OK, but ...
- 10 or less right - more training needed

15. b  
14. d  
13. a  
12. c  
11. e  
9. at night

5. insecticide did not reach pest  
6. larvae  
7. thatch, soil and roots  
8. organophosphate

1. adults  
2. False  
3. larvae  
4. Less than 1 percent