

Service

Service. You give it to your customers and you expect it from your suppliers.


We understand how important service is to you. That's why you can depend on Special Products for prompt delivery, quality products, answers to your technical questions, experienced sales representatives and an organization from top to bottom that puts you first.

"Great Service." It sets you apart from your competition. It sets us apart from our competition.

SPECIAL PRODUCTS

A **UAP** COMPANY

ON THE MOVE



© 1991 CIBA-GEIGY Corporation, Turf and Ornamental Products, Box 18300, Greensboro, NC 27419. Always read and follow label directions.

Ciba-Geigy Corp., P.O. Box 18300, Greensboro, NC 27419

**THE ONLY
OTHER ACTIVE
INGREDIENTS
YOU'LL NEED TO
HELP PREVENT
12 MAJOR TURF
DISEASES.**

It's as effective a fungicide as you can buy. Banner® lasts for up to 4 weeks, at rates of only 1-2 ounces per thousand square feet.

To make the most of it, however, proper diagnosis is crucial. So, too, is a complete understanding of your golf course's disease history.

Knowing this, you can vary individual doses to precisely what you need.

Want the cost savings of prescriptive control? Get Banner. As they say, it's a no-brainer.





POST-EMERGENCE WEED CONTROL IN WARM-SEASON GRASSES

Post-emergence herbicides provide the turfgrass manager with viable options to control weeds during the entire year.

by Tim R. Murphy, Ph.D., University of Georgia

Weed control is the process of limiting a weed infestation so that a turfgrass can properly grow, develop and be aesthetically appealing.

A balanced turfgrass weed control program uses a combination of cultural, mechanical and chemical weed control practices. When properly maintained, warm-season turfgrasses are highly competitive with weeds. Adherence to recommended fertility programs, water requirements, mowing heights and schedules and controlling diseases and insects will dramatically improve the success of the chemical weed control program.

The use of herbicides, in the absence of approved cultural and mechanical practices, will not result in a high quality, warm-season turfgrass.

Turfgrass managers can choose from two types of herbicides to control weeds in warm-season turfgrasses. Pre-emergence herbicides form the base of the chemical weed control program. They are primarily used for the control of crabgrass, goosegrass, annual bluegrass and certain annual broadleaf weeds. Post-emergence herbicides are generally used to control weeds that are not controlled by pre-emergence herbicides, or in the event of a pre-emergence weed failure, as a reliable backup.

Post-emergence herbicides offer several advantages over pre-emergents.

Spot treat, or as needed


Post-emergence herbicides can be applied on a "spot treatment" or "as-

needed" basis, directly to a weed infestation; pre-emergence herbicides are usually applied to the entire turfgrass area.

Spot treatments of post-emergence herbicides are less expensive than blanket applications of pre-emergence herbicides. Post-emergence herbicides will control many problem annual and perennial weeds not controlled by pre-emergence herbicides.

Low rates of most post-emergence herbicides may be used on newly-sprigged or sodded warm-season turfgrasses.

In areas that are scheduled to be overseeded or renovated, the majority of post-emergence herbicides can be used up to one month before renovation. The time interval from application to seeding, sprigging or sodding operations for pre-emer-



Our Mid-size Classic continues to impress long after the pricetag has been discarded.



When you can have Ransomes quality at a competitive price, there's no reason to accept second best. ■ The Bob-Cat Mid-Size is a legend in the commercial mowing market and the Classic continues to impress. A true commercial machine throughout, with the "value-added" features you want. ■ Available in 36" and 48" cutting widths and variable speed, the Bob-Cat Classic features a Kohler 12.5 HP overhead valve engine for better fuel economy. Includes wide tires for superior climbing and three grass catcher options. And, it all comes to you at an affordable price, along with 48 hour parts delivery. Guaranteed. ■ Call your Ransomes dealer today or call us at 414-699-2000 to arrange a free demonstration. And step up to the best, a Bob-Cat Classic.



Where great ideas start.

*Does not include sales tax, freight, or dealer prep; suggested price listed in US dollars. Price may vary with dealer and region.

©RANSOMES INC., A RANSOMES COMPANY, 1991, ONE BOB CAT LANE, JOHNSON CREEK, WI 53038

TABLE 1. Common and trade names of warm-season turfgrass postemergence herbicides.

Common Name	Trade Name(s)	Company(s)
asulam	Asulox	Rhone-Poulenc
atrazine	Aatrex, Others	Ciba-Geigy, Others
bentazon	Basagran	BASF
bromoxynil	Buctril	Rhone-Poulenc
2,4-D	Numerous formulations are available.	Vertac, Lesco, Fermenta, Others
2,4-D + dicamba	Eight-One, Phenaban 801	Lesco, PBI/Gordon
2,4-D + dichlorprop	Weedone DPC Amine, Weedone DPC Ester	Rhone-Poulenc
2,4-D + mecoprop	Lescopar, Phenomec 2+1, 2 Plus 2	Lesco, PBI/Gordon, Fermenta
2,4-D + mecoprop + dicamba	Trimec Classic, Trex-san, Three-Way	PBI/Gordon, Sierra, Lesco
2,4-D + mecoprop + dichlorprop	Weedestroy Triamine, Weedestroy Tri-Ester	Riverdale
dicamba	Banvel, Dicamba 4	Sandoz, PBI/Gordon
diclofop-methyl ¹	Illoxan, Hoelon	Hoechst
diquat ²	Diquat	Valent
DSMA	Numerous formulations are available.	Vertac, Vineland, Others
ethofumesate	Prograss	Nor-Am
fenoxaprop	Acclaim	Hoechst
glyphosate	Roundup	Monsanto
imazaquin	Image	American Cyanamid
mecoprop	Mecomec, Lescopex	PBI/Gordon, Lesco
mecoprop + 2,4-D + dicamba	Southern Trimec	PBI/Gordon
MCPA + mecoprop + dichlorprop	Weedestroy Triamine II, Weedestroy Tri-Ester II	Riverdale
metribuzin	Sencor Turf	Mobay
metsulfuron	DMC	O. M. Scott
MSMA	Numerous formulations are available.	Fermenta, Platte, Others
MSMA + 2,4-D + mecoprop + dicamba	Trimec Plus	PBI/Gordon
pronamide	Kerb	Rohm-Haas
sethoxydim	Vantage	BASF

¹ Diclofop-methyl currently has a state label for use in Georgia, Florida and South Carolina. Current labels should be consulted as use directions and trade names vary between states.

² Diquat has a state label in Arkansas, Louisiana, Mississippi, Oklahoma, Tennessee and Texas for winter annual weed control in dormant bermudagrass.

Source: The author

TABLE 2. Warm-season turfgrass tolerance to postemergence herbicides.

Herbicide	Turfgrasses					
	Bahia-grass	Bermuda-grass	Centipede-grass	Carpet-grass	St. Augustine-grass	Zoysia-grass
asulam	NR-S	T ¹	NR-S	NR-S	T	NR
atrazine	NR-I	I	T	NR-I	T	I
bentazon	T	T	T	NR-I	T	T
bromoxynil	T	T	T	NR-I	T	T
2,4-D	T	T	I	I	S-I	T
2,4-D + dicamba	T	T	S-I	S-I	S-I	T
2,4-D + dichlorprop	T	T	I	I	I	T
2,4-D + mecoprop	T	T	S-I	I	S-I	T
2,4-D + mecoprop + dicamba	I-T	I-T	S-I	S-I	S-I	T
2,4-D + mecoprop + dichlorprop	T	T	I	I	I	T
dicamba	T	T	I-T	I	S-I	T
diclofop-methyl	NR	T	NR	NR	NR	NR
DSMA,MSMA	NR-S	T	NR-S	NR-S	NR-S	I
fenoxaprop	NR-S	NR-S	NR-S	NR	NR-S	T
glyphosate ²	S(D)	S(D)	S	S	S	S
imazaquin	NR-S	T	T	NR-I	T	T
MCPA + mecoprop + dichlorprop	T	T	I	I	I	T
mecoprop	T	T	S-I	I	S-I	T
metribuzin	NR-I	T	NR-S	NR-S	NR-S	NR-S
metsulfuron	NR-S	T	NR-T	NR	T	NR-T
pronamide	NR-S	T	NR-I	NR	NR-I	NR-I
sethoxydim	NR-S	NR-S	T	NR-I	NR-S	NR-I

T = Tolerant at labeled rates; I = Intermediate tolerance, use at reduced label rates; S = sensitive, do not use this herbicide; D = Dormant; NR = Not registered for use on this turfgrass.

¹ Labeled only on 'Tifway' (419) bermudagrass and St. Augustinegrass.

² Bahiagrass and bermudagrass are tolerant to glyphosate when completely dormant.

Source: The author

differently to the same herbicide. For example, Meyer zoysiagrass has better tolerance to MSMA than Emerald or Matrella.

The most important factor in selecting a post-emergence herbicide is the tolerance of the turfgrass to the herbicide. Refer to the specific label to determine if the herbicide may be used on a particular turfgrass species.

Weed species. Similar to turfgrasses, weed species vary in their susceptibility to herbicides. Correct weed identification is a prerequisite for selecting an appropriate herbicide. Weed identification manuals and identification assistance is available at many county extension service offices. Several chemical companies also distribute excellent weed identification guides.

Time of application. The time of year that a herbicide is applied can influence turfgrass tolerance. For example, dormant bahiagrass and bermudagrass have excellent tolerance to Roundup. However, severe injury will occur if this herbicide is applied to semi-dormant or actively-growing bahiagrass or bermudagrass.

The risk of injury from post-emergence herbicides is also greater during the spring green-up process (transition from winter dormancy to active growth) than when the turfgrass is fully dormant or actively growing.

Post-emergence herbicides such as 2,4-D + mecoprop + dicamba (Trex-san, Trimec Classic, Three-Way and others) have been shown to slightly decrease the quality of Tifway bermudagrass when applications were made three weeks before or during spring green-up. Image can also cause slight to moderate delays in green-up if applications are made during spring transition.

Research has shown that the decrease in turfgrass quality that may result from using post-emergence herbicides

gence herbicides varies from 1½ to 4 months.

Herbicide selection

Many post-emergence herbicides are available to control weeds in warm-season turfgrasses (Table 1). They all have different qualities.

Turfgrass tolerance. The warm-season turfgrasses dramatically vary in their tolerance to post-emergence herbicides (Table 2). Bermudagrass has good tolerance to MSMA and DSMA; however, carpetgrass, centipedegrass and St. Augustinegrass are severely injured by these herbicides. With the exception of bahiagrass and carpetgrass, warm-season turfgrasses have excellent tolerance to Image.

Additionally, cultivars within a species may respond

IT'S EASY TO SPOT A LAWN THAT HASN'T BEEN TREATED WITH DYRENE.

Leaf spot is a dead give-away for any lawn that's been treated with something other than DYRENE® Fungicide. That's because only





DYRENE can give you this kind of effective, long-lasting protection.

In fact, DYRENE widens your window of application with as much as 28 full days of control. So you

have the time to reach all your customers before the leaf spot takes hold, or spreads out of control.

Ask your distributor about DYRENE today. Without it, you could find yourself in a pretty tight spot. For more information, contact Mobay Corporation, Specialty Products Group, Box 4913, Kansas City, Missouri 64120. Or call (800) 842-8020.



Mobay 

A Bayer USA INC COMPANY

Circle No. 118 on Reader Inquiry Card

during green-up persists for two to six weeks after application. Also, in turfgrasses that are severely infested with weeds, better turfgrass growth eventually results due to the elimination of the thick cover of weeds.

In the event that a dense weed population necessitates using a post-emergence herbicide during green-up, use only the lowest recommended or one-half the recommended rate. Low or one-half rates will minimize herbicide injury to the turfgrass.

Application frequency. For some weed species, a repeat application is necessary to effectively control the weed. For example, two applications of MSMA + Sencor, at a 7- to 10-day interval, are necessary to control goosegrass. In contrast, small crabgrass can often be controlled with a single application of MSMA. However, large, well-tillered crabgrass usually requires two applications of MSMA or DSMA, each at a 7- to 10-day interval.

Ornamental tolerance. Turfgrass herbicides are commonly applied to sites containing ornamental plantings. Ornamentals may be injured by spray or vapor drift or by root absorption of the herbicide. Vapor drift is the movement of herbicide vapors from the intended site of application.

Ester formulations of the phenoxy herbicides (2,4-D, dichlorprop) easily volatilize during warm temperatures and can injure sensitive ornamentals by vapor drift. Ester formulations

Problem weed management

Bahiagrass: Repeat applications of MSMA or DSMA at 7- to 10-day intervals will control bahiagrass in MSMA/DSMA tolerant turfgrasses. In bermudagrass and St. Augustinegrass, DMC will effectively control bahiagrass. In centipedegrass, repeat application of Vantage (formerly Poast) at 10- to 14-day intervals will suppress bahiagrass growth and seedhead development.

Dallisgrass: It is believed that most pre-emergence herbicides will control dallisgrass that arises from seed.

Established dallisgrass can be controlled in bermudagrass or zoysiagrass with repeat applications of MSMA or DSMA. Applications should be made to actively-growing dallisgrass. Also, a non-ionic surfactant at 0.25 percent v/v is recommended with MSMA or DSMA for dallisgrass control. Staying on the application schedule (2 to 4 applications, each at a 7- to 10-day interval) will be required to control dallisgrass. Shortening the application interval to five days may help on sites where dallisgrass has been difficult to control with MSMA or DSMA.

Nutsedge: Basagran will provide good control of yellow nutsedge, but not of the purple variety. Monthly applications of MSMA or DSMA in tolerant turfgrasses during the late spring and summer months can be used to suppress the growth of both species. With the exception of bahiagrass and carpetgrass, Image can be used in warm-season turfgrasses for yellow and purple nutsedge control. The addition of MSMA to Image generally improves nutsedge control in MSMA tolerant turfgrasses. A repeat application, six to eight weeks after the first application, of Image or Image + MSMA will be required to control nutsedge during the summer

months.

Prostrate spurge: Control requires repeat applications of two-way or three-way broadleaf herbicides. In bermudagrass, low rates of Sencor (0.125 to 0.25 lb. AI/acre) will effectively control emerged prostrate spurge.

Virginia buttonweed: Rapidly becoming the number one problem broadleaf weed in southern

suppress the growth of Virginia buttonweed. Research conducted in Mississippi has shown that applications of Ronstar or Princep at the time of the 2,4-D + dichlorprop application increased the control of Virginia buttonweed.

The increase in control with Ronstar and Princep is believed to be the control of Virginia buttonweed plants that arise from seed. Turfgrass managers should be aware that Princep is not labeled on southern turfgrasses in the months of June, July and August. However, depending upon the geographical location, a mid-April or May application of Princep may assist in the control of Virginia buttonweed.

Wild garlic: A perennial that appears in turfgrasses in the mid- to late-fall months. Fall (November) and winter (January-February) applications of 2,4-D or two-way or three-way products that contain a phenoxy herbicide or dicamba will control wild garlic. The fall plus early winter treatment program will need to be repeated for two to three consecutive years to effectively eliminate this weed from turfgrasses. Late fall to early winter applications of Image has provided good to excellent control of emerged wild garlic. Image slowly kills wild garlic and treated plants will remain visible in the turfgrass for an extended time after application. Mowing one to two weeks after the Image application will remove a significant portion of the wild garlic foliage and improve the overall appearance of the turfgrass. Image should be applied to emerged wild garlic in the late fall or early winter after the first killing frost or the onset of winter dormancy of the warm-season turfgrass.

In St. Augustinegrass and bermudagrass, DMC will also effectively control wild garlic.

—Dr. Murphy □



turfgrasses. A warm-season perennial, Virginia buttonweed reproduces by seed, cut plant pieces, and fleshy roots.

Research has shown that 2,4-D + dichlorprop (Weedone DPC amine and ester) has provided better Virginia buttonweed control than other two-way or three-way broadleaf herbicides. Monthly applications of 2,4-D + dichlorprop will be needed during the summer months to

"This Baby Will Steal Your Heart."

"When you put as much into something as I have with this new 30-H, it gets close to you. But I think it's the kind of machine you're going to love. It's got a Yanmar diesel that turns out 30 horses for all the power you'll ever need. The 72" deck gives you the big cut you want for wide area mowing. Talk about smooth. The hydrostatic transmission really gets the job done. And our special rear-wheel steering leaves a "0" uncut circle. Naturally, the 30-H has all the dependability we build into every Gravelly. Let me tell you, this baby will steal your heart."

Ask your local Gravelly dealer to introduce you to the powerful, new 30-H today. **GRAVELLY**

Jan Miller

Jan Miller, Director of Engineering
Gravelly International, Inc.
Clemmons, North Carolina

Fall In Love With A Gravelly

Gravelly International, Inc., P.O. Box 5000, One Gravelly Lane, Clemmons, NC 27012 • 919-766-4721 • Telefax: 919-766-7545

Circle No. 221 on Reader Inquiry Card

Suggestions for use

Post-emergence herbicides are applied after annual weeds emerge or when new growth of perennial weeds appear in the turfgrass. Follow these guidelines for better weed control and improved turfgrass tolerance.

1. Small, actively-growing weeds.

Perennial and annual weeds that are growing under good soil moisture conditions at moderate air temperatures are easier to control with post-emergence herbicides than weeds that are stressed due to adverse environmental conditions. Target the application to coincide with good soil moisture conditions at air temperatures of 60 to 90°F. Applications on cold, wintery days, or to drought-stressed weeds will result in poor weed control.

2. Do not apply post-emergence herbicides to turfgrasses and weeds that are stressed due to high air temperatures or drought.

The tolerance of warm-season turfgrasses to post-emergence herbicides decreases at air temperatures greater than 90°F, and when turfgrasses are drought-stressed. Additionally, turfgrasses growing under high soil moisture, high relative humidity and high air temperatures, (less than 90°F) have a lower level of tolerance to post-emergence herbicides than turfgrasses growing under similar moisture and humidity conditions but at cooler temperatures.

Herbicides that contain 2,4-D; dicamba; mecoprop; dichlorprop; MSMA and DSMA should not be applied at high air temperatures since there is a high risk of increased turfgrass injury.

Follow label guidelines.

3. Repeated applications at low rates will generally improve weed control and turfgrass tolerance.

Single applications at high rates generally cause more turfgrass injury than repeat applications at low rates. Additionally, single, high rate applications often do not control the weed, particularly perennial weeds. The repeat application is usually made at intervals of 7

to 14 days after the first application, or when regrowth of the weed is noted. Refer to the label for information regarding repeat treatments.

4. Coordinate mowing schedules.

A general recommendation is to delay mowing three to four days before or after a post-emergence herbicide application. The delay prior to application will increase the leaf surface area of the weed and result in better spray coverage and control. The delay after application is necessary to allow adequate time for herbicide absorption and translocation in the target weed species.

5. Do not apply post-emergence herbicides immediately before rainfall or irrigation.

The effectiveness of most post-emergence herbicides is better when rainfall or irrigation does not occur for 6 to 24 hours after application. Rainfall or irrigation immediately after application can wash the herbicide from the treated weed foliage and decrease control.

6. Use surfactants and crop oil concentrates according to label directions.

The effectiveness of many post-emergence herbicides is enhanced by the addition of a crop oil concentrate or surfactant to the spray mixture, particularly under less than ideal spray conditions. However, indiscriminate use of surfactants or crop oil concentrates can increase the risk of turfgrass injury. Check the label for crop oil guidelines.

7. Calibrate spray equipment, train operators.

Weed control failure can be linked to improper calibration of spray equipment.

The tolerance of warm-season turfgrasses rapidly decreases at elevated or higher-than-recommended rates of post-emergence herbicides. Training assistance is available through most county extension service offices and spray equipment company representatives.

—Dr. Murphy □

should not be used during the warm months of the year when environmental conditions are favorable for volatilization. Spray drift damage can be minimized by spraying when the wind velocity is less than 5 mph, and selecting a nozzle tip and spray pressure that produces large spray droplets.

Due to their soil residual characteristics, Aatrex and dicamba (Banvel, Dicamba 4) can injure broadleaf ornamentals via root uptake. Injury to ornamentals is most likely to occur on sandy soils when a heavy rainfall immediately follows application of these herbicides. The potential of these herbicides to injure ornamentals via root uptake can be prevented by avoiding their use over the rootzone of shrubs and small trees.


Spring break important

Post-emergence herbicide use should be avoided during the spring transition of warm-season turfgrasses and during periods of extremely high air temperatures.

winter, late spring and early summer months. The majority of turfgrass post-emergence herbicides are more effective when applied to small weeds than when applied to large weeds. Waiting until weeds are in advanced growth stages will usually require that the highest recommended rate be used, and often, a repeat application.

In the event of a pre-emergence herbicide weed control failure, post-emergence herbicides can be relied upon for a complete chemical weed control program. Due to the necessity of repeat applications and temporary turfgrass injury, most managers prefer to use a combination of pre-emergence and post-emergence herbicides to control turfgrass weeds.

Cultural practices that favor vigorous turfgrass growth, the use of pre-emergence herbicides at recommended times of the year, and timely use of post-emergence herbicides will enable the turfgrass manager to achieve the goal of a high quality, attractive turfgrass that is relatively weed-free. LM



Agronomist Tim R. Murphy specializes in weed science at the University of Georgia Cooperative Extension Service.