

# New **low odor** technology

## Same outstanding performance

*"The low odor is an advantage with our high-end commercial accounts where customers are coming and going all day, as well as with our high-end residential accounts who don't need to close their windows or postpone outdoor activities to avoid the pesticide odor."*

Tom Mieyal, General Manager, of a major Cleveland, Ohio lawn care company.

Trimec® Classic Broadleaf Herbicide is now a "low odor" product. This new characteristic is not created with a masking agent. Advancements in our production technology have enabled us to manufacture Trimec® Classic as a low odor product. If you've been using a "cheaper" three-way and adding a masking agent, now would be an excellent time to switch to quality -

**TRIMEC CLASSIC**  
BRAND  
BROADLEAF HERBICIDE



**G** pbi / Gordon  
CORPORATION

An Employee-Owned Company

**1-800-821-7925**

[www.trimec.com](http://www.trimec.com)



# Northern style

*Turfgrass that is established and maintained properly, is healthy turfgrass and doesn't provide "room" for weeds to compete and spread.*

By FRANK S. ROSSI, Ph.D.

**T**he coexistence of turfgrass and weeds is the basis of the age-old tip for controlling weeds in turf: maintain a healthy, dense stand of turf that prohibits weeds from establishing.

This is based on two important ecological concepts space and competition. Therefore, the foundation of a turfgrass weed management program must be to implement management programs that favor the competitive advantage of turf, while minimizing bare areas where weeds can invade.

Understanding the ecological concepts then aids the turfgrass manager when addressing the aesthetic and functional expectations. For example, while we understand the environmental benefits of a well managed lawn turf, essentially lawn management is focused on a high aesthetic demand; the lawn has to look good! In contrast, sports turf or golf turf is focused equally on aesthetic and functional quality, in that, not only does it have to look good, put it must sustain regular surface disruption that results in gaps where weeds can invade.

It follows then that a lawn turf should be easier to keep weed free because less gaps should occur, yet, it seems almost the opposite. Lawn managers and homeowners



spend much more time managing weeds than a sports or golf turf manager. Why? Most likely it is related to the regular attention paid by sports and golf turf managers to maintain a dense vigorous stand under regular surface disruption. In contrast, many lawn managers visit the lawn intermittently, and often, do not have direct control over the key cultural practices like mowing and watering that sustain a dense healthy turf.

#### **Where Does Weed Management Start?**

These ecological principles will always work in your favor as a turf manager if the proper decisions are made during turfgrass establishment regarding site preparation, soil modification, turf selection, and establishment procedure. Visualizing an integrated weed management program as a triangle (see next page), it becomes clear that proper site assessment, soil modification,

#### **Ground ivy control research finds timing is more important factor.**

and species selection form a solid foundation for the life of the stand. Whereas, mistakes at establishment, limits options to pesticide use and consequently an unstable foundation, evident in the inverted triangle.

Continuing to visualize an integrated weed management approach, the primary cultural practices of mowing, watering and fertilizing should focus on maximizing root growth. The ability to establish and sustain a healthy root system will always make the turf more forgiving of environmental, biological and traffic stress. For example, maintaining a higher than usual height of cut (3" or greater) will promote deep rooting and shade the soil surface. Irrigating judiciously, so as to avoid a moist soil surface



that will encourage weed seed germination. Finally, the most efficient fertilization programs for cool season turf is focused in the fall. This approach is based on maximizing energy production under cooler temperatures without the surge in top growth associated with spring conditions. Still, if turf density is low in spring, a fertilization will be needed to increase density (to fill the space with turf!).

#### Characterize Your Weed Management.

In keeping with traditional Integrated Pest Management (IPM) programs, it is vital to map and monitor weed populations (as well as turf species). Maintaining records of populations over time provides unique insight into the response of the turf and the weeds to different environmental conditions and management programs. For example, you may have decided to change a fertilizer program to a more water soluble source and notice from your mapping that patches of annual bluegrass have become established. Following a dry year, you may notice increase in clover populations. Over time this information can assist with weed management programs.

Still, the cornerstone of an IPM approach is the establishment of thresholds. Weed thresholds are slightly different than for other pests, in that we must have an aesthetic and a functional threshold. An aesthetic threshold for weeds could be defined as the point at which the number of weeds in a turf reduces the visual quality below an acceptable level. Subsequently, the functional weed threshold is the point at which the number of weeds present reduces the functional quality (soil stabilization, traction, elasticity, etc.) below an acceptable level.

As you might imagine establishing thresholds can be subjective. Do you establish visual weed thresholds with "curb appeal"? That is do you evaluate the visual quality from a distance, or by standing over the turf? What season do you determine

threshold? Is there any tolerance for weed invasion or must the turf be 100% weed free? The same questions arise for functional quality. How many dandelions or prostrate knotweed plants can you have in a soccer field before the players game is affected? How many patches of clover are tolerated in a golf course rough area before you hear complaints of playability? How many crabgrass or broadleaf plantain plants can a lawn tolerate before soil movement is increased?

Viewing weed management from this perspective will challenge the turf manager to communicate with their clientele to assist in establishing thresholds. It has been my experience that this dialogue with our clientele (homeowners, athletes, coaches, and golfers) is rarely conducted. We assume it must be 100% weed free and do not share the responsibility of establishing thresholds with the persons who receive our service. It follows then that when questions regarding pesticide use arise, we are unprepared to explain how we conduct an IPM program for weeds. Furthermore, we must have a realistic basis for decision making prior to implementing a management program designed to reduce or eliminate weed populations. This will become more important as chemical pesticide use becomes more restrictive.

#### Timing is everything

The old adage that "timing is everything for a successful life" rings true when considering chemical weed control. First, we must decide whether we will chose a preventative approach with preemergence herbicides, or a curative approach with postemergence herbicides. Clearly the preventative approach requires less labor and knowledge, as monitoring and mapping efforts are not used reveal threshold populations. This approach is widely used, primarily as a result of the "zero threshold" that exists for weed populations. The curative approach might be more labor and



Think of weed management as a triangle. At its base it starts with proper site and species (or variety) selection. Then comes



necessary cultural practices like watering, fertilizing and mowing. Finally, if weeds remain a problem, we use herbicides as tools.

knowledge intensive, in that it will require timely monitoring (you'll have to know when weeds might emerge or flower) as well as knowledge of how to identify the major cool season weeds. Additionally, research has indicated that effective postemergence control relies on proper timing, often based on the growth stage of the weed.

Effective preemergence control, usually of summer annual grass weeds such as crabgrass and goosegrass, relies on understanding the seed germination and emergence time. In essence, this can be viewed as "seed bank management". Regarding germination, not all weed seeds in the soil will germinate every year. Plants have de-



veloped dormancy mechanisms for a percentage of their seeds that keep them viable for many years (waiting for the right time and space). For example, research indicates less than 50 percent of the crabgrass seed produced in the previous fall will germinate in the spring. However, turf areas with a history of crabgrass invasion will have a high percentage of seed viable from several years of production. The management question is then, how long do you have to use a preemergence herbicide before the seed bank is depleted? I am unaware of research that might address this question.

Emergence time has recently been investigated by researchers at the University of Maryland and currently here at Cornell. This research provides new insight into the timing of emergence and the length of time that crabgrass seeds germinate in a season. In a dense turf under a medium maintenance program, soil temperatures greater than 73° F were required for significant emergence. In addition, monitoring the emergence time with base 50° F growing degree days (GDD) accumulated from March 1, the Maryland study found that 25% of the total emergence occurred by 200 GDD, 75% by 600 and 100% by 1100. This suggests that under average turf density, germination and emergence can occur for 10 to 12 weeks depending on the season. This has enormous implications on timing of preemergence applications in the early spring. Simply, if applied to early, the effectiveness of the preemergence herbicide may dissipate and allow breakthrough.

Postemergence timing has received increased attention over the last decade as new materials have become available, hard to control weeds have emerged, weed tolerance levels have declined, and the use of preemergence herbicides come under question. First and foremost, the key to effective postemergence control is proper identification, followed by understanding weed biology so as to time applications for maximum effectiveness.

Interesting research on yellow nutsedge

control has indicated that effective long term control requires a systemic herbicide to eliminate the surface vegetation (the leaves) and to be translocated to the forming nutlet (perennial storage organ), thus preventing it from emerging the following year. Consequently, the most effective application timing, based on the information above, is late June, early July when day length begins to shorten. This event apparently triggers the translocation of energy in the nutsedge plants from the leaves to the nutlets. Effective materials such as Basagran and Manage will provide long term control applied at this time.

Successful control of perennial broadleaf weeds is best accomplished in the late summer, early fall. Certain materials such as Confront, have been shown to provide activity on broadleaf weeds down to 37° F. Still, most of our clientele is not willing to wait until this time of year for control. Therefore, effective timing of spring applications is needed. Researchers at Purdue University have reported that spring control of dandelion is most effective when based on a simple growing degree day (GDD) model. They concluded that formulation of Weedone and time of application were critical for maximizing control. If using the ester formulation of Weedone, it is best applied around 100 to 150 base 500 F GDD, while the amine formulation should be applied much later at 250 to 300 GDD. This is useful not only for control, but also for minimizing potential vapor drift resulting from applying the ester formulation later in the spring when conditions could favor volatilization.

Historically hard to control weeds such as ground ivy (creeping charlie, gill-over-the-ground, etc) could also be more effectively controlled using some research on improved timing. A three year study at the University of Wisconsin-Madison indicated that any herbicide formulation that included 2,4-D as an active ingredient would control ground ivy if applied at full bloom in the spring or immediately following the first frost in the late summer early fall. The results were dramatic, however,



Researchers at Purdue University have reported that spring control of dandelion is most effective when based on degree days.

there does not appear to be an easy explanation. Unfortunately, wild violet, another hard to control weed, in the plots was not controlled using the same approach.


#### Developing an integrated approach

Turfgrass managers should regularly strive for more resource efficient management programs that incorporates a knowledge-based approach of biology and ecology to maximize turf health and maintain adequate density. This information is then utilized through effective monitoring and mapping of plant populations and taken together serve as the basis for weed control decisions. As the industry continues to improve our overall communication skills, we will have a more open dialogue for establishing thresholds and making more cooperative decisions regarding weed management that will include herbicide use and possibly biological/organic based approaches as well. **LM**

Frank Rossi is an assistant professor of turfgrass science and extension turfgrass specialist at Cornell University



# INSURANCE IS ONE THING YOU SHOULDN'T HAVE TO WRESTLE WITH.



## INTRODUCING THE NEW BUSINESS PACKAGE POLICY (BPP)\* FROM FLORISTS' MUTUAL.

We're proud to present the most flexible business package policy the Floral/Horticultural industry has ever seen. Now buildings, greenhouses, contents, crops, computers, business income and public liability are available in a single policy. We can custom-tailor the BPP to meet your unique needs.

Perhaps best of all, you'll be covered by Florists' Mutual, the industry's insurance specialist for more than a century. Call us today for a free coverage evaluation and custom-tailored

Business Package Policy proposal.



Growing With You For  
More Than A Century

500 St. Louis Street, Edwardsville, IL 62025-1573

1-800-851-7740

FAX 1-800-233-FMIC

<http://www.plantnet.com>

\*May not be available in all states.



## Southern style

*Winter broadleaf weeds are an opportunity or a headache for the turfgrass manager. If control is late, customers should expect multiple applications and slow results.*

By BERT McCARTY

**E**arly spring is a great time to be in the southern United States. Everyone is stirring after a long chilly winter, and the blooming dogwoods, azaleas, redbuds, and others plants help renew the faith of warmer times ahead. However, for the professional turfgrass manager, late winter and early spring often rings in the new season with a deluge of irate customers calling about winter weeds.

This time is when winter annual weeds really begin to grow, eventually producing flowers. By their nature, winter weeds usually do not die until hot temperatures of early summer. Until then, these weeds are an eyesore for homeowners and because of their maturity, difficult to control for the professional turfgrass manager.

Usually, multiple applications are required to control these which really drive up labor costs. One also needs to be formulating and applying a strategy for pre-emergence crabgrass control at this time.

With numerous customers, the logistics of performing these duties really separate the turf pros from the Johnny-come-latelies.

### A weed management program

Weed management is an integrated process where good cultural practices are employed to encourage desirable turfgrass ground cover as well as the intelligent selection and use of herbicides (Table 1).

The first step to successful weed management is proper identification. Due to mowing of flowers and seedheads, turfgrass managers are forced to identify weeds on the basis of the vegetative structures, such as ligules, leaves and stems. An excellent weed identification guide is *Weeds of Southern Turfgrasses*, a publication available through Clemson University for \$8.00 per copy. Call 864-656-3261 for ordering information. This publication is also available through the Cooperative Extension Services in Georgia, Alabama, and Florida.

Winter weeds germinate in late summer through early fall when daytime temperatures consistently drop in the 70's. These grow throughout the winter months, and flower or produce seedheads during late winter and early spring. Winter weeds are sneaky in that they blend with the turf in the fall and early winter months and do not become noticeable until late winter when growth spurts, along with seedheads and flowers, produce a ragged appearing turf.

### Scouting

Information on which weeds, where they occur, and relative level of occurrence are needed in making informed management decisions on if and which control option(s) to consider. Scouting simply means breaking the service area into logical sections or units and determining which weeds are present and at what level. Normally for lawns, these sections are broken into front, back and side yards. Due to visibility, front lawns generally require control implemen-

tation first followed by the sides, and finally, by the back yard section. Golf courses are sectioned into tees, fairways, greens, and roughs for each hole. Roughs receive least attention for weed control while greens and tees receive the most. Weed cover patterns can be as elaborate as estimating the percent weed cover for each unit or more realistically, involve a rating system of being widespread, spotty or in a single patch. Threshold levels needed before treatment justification is generally determined by the owner or manager of the turf site.



Henbit (*Lamium amplexicaule* L.) has greenish to purplish, tender, four-sided stems.

Optimum scouting timing for winter weed control is in early fall (September/October) with a follow-up in early spring (March/April). The fall scouting allows early detection and pressure level assessment of each. The early spring scouting indicates which weeds were not controlled and where they can be expected to occur the following winter season.

*cont. on page 48*

**Open letter  
to all  
Grounds  
Professionals:**



**PGMS**

***It's more than OK to come home!***


You have a home with your fellow grounds professionals at PGMS. In fact, PGMS is the only place for grounds professionals as individuals.

You deserve your own society, your own place, where you can relate to other grounds pro's — as individual professionals, not just as an employee of some organization.

In fact, PGMS is more stable than most organizations. We've been around for more than 85 years, and are continuing to build for your interests. Serving the individual professional is the only reason PGMS was established in 1911, and will remain our only focus into the next century and beyond.

Again, this is YOUR place. Here are some of the reasons why:

- Annual Conference and Green Industry Expo
- Monthly Newsletter
- Training Manuals\*
- Local Branches
- Certified Grounds Manager Program
- Groundskeeper Certification
- Discounts — Rental Cars, Books
- Complimentary Green Industry Magazines
- Membership Directory
  - Personal ID (membership card, decals etc.)
  - Awards



Includes the three most respected operating manuals for grounds professionals: Grounds Maintenance Estimating Guidelines, Grounds Maintenance Management Guidelines and Grounds Management Forms & Job Descriptions.

Overall, PGMS provides the opportunity for you to network with, learn from, help instruct, etc. those with your same interests, needs and problems.

**The question remains:** If you are a grounds professional, why are you trying to do it alone, or simply as an employee of some organization? Don't you deserve the advantages of joining with fellow grounds professionals in PGMS? In the '90's, more than ever before, employment and individual betterment statistics indicate that each professional had better be an individual agent.

Why not take a closer look by requesting more information? Just complete and mail the following form, and information will be rushed to you.

To: PGMS  
120 Cockeysville Road, Suite 104  
Hunt Valley, Maryland 21031 Tel. (410) 584-9754 FAX (410) 584-9756



**I want to grow with PGMS. Please send more information explaining how.**

Name \_\_\_\_\_ Orgn. \_\_\_\_\_

Street \_\_\_\_\_

City/State/Zip \_\_\_\_\_

Telephone \_\_\_\_\_



## TABLE 1. WINTER WEED MANAGEMENT SCHEDULE FOR WARM SEASON TURFGRASSES

### LATE SUMMER

To build root carbohydrates, apply winterizing fertilizer which supplies adequate potassium and mow at the upper recommended mowing height. Apply preemergence herbicides for annual bluegrass if scouting the previous spring warrants it. Scout and map areas with crabgrass to pinpoint herbicide needs for the following spring.

### EARLY/MID FALL

If needed, apply postemergence herbicides for annual bluegrass control.

### MID FALL

If needed, apply postemergence herbicides for broadleaf weed control.

### EARLY WINTER

Reapply postemergence herbicides for broadleaf weeds and annual bluegrass, if necessary.

### WINTER

Calibrate and repair sprayers; evaluate the previous year's weed control strategies; plan for the upcoming year's strategy.

### LATE WINTER

Apply preemergence herbicides for crabgrass control. Apply postemergence herbicides for broadleaf weed control for new customers.

### EARLY SPRING

Apply preemergence herbicides for goosegrass control. Repeat broadleaf weed control application, if necessary, for new customers. Follow-up scouting for remaining winter weeds for formulating the upcoming fall control strategies.

### SUMMER

Maintain optimum fertility, watering, and mowing height and frequency to encourage healthy, thick turf stands which helps discourage weeds and other pests.

cont. from page 46

### Herbicide Selection and Use

Preemergence herbicides are applied prior to weed seed germination and prevent development of the germinating seed. If applied after germination, preemergence herbicide effectiveness greatly diminishes. Preemergence herbicides should be activated by  $\frac{1}{2}$  to  $\frac{1}{4}$  inch rainfall or irrigation after application.


Postemergence herbicides generally are effective only on visible weeds. Young (two to four-leaf stage) and actively growing weeds are the most susceptible and require least amount of herbicide. At this stage, herbicide uptake and translocation are favored and weeds have less developed, more tender root systems. Waiting later results in poorer translocation of applied materials in plants, more difficulty controlling mature plants, and possible setback of turf during green-up.

Postemergence herbicides should only be used when weeds are actively growing. This primarily occurs with temperatures between 40 and 80°F. Applications outside this temperature range are too slow to either be effective or result in excessive turf damage.

### Broadleaf Weed Control

Preemergence broadleaf weed control is provided by the herbicide Gallery. Gallery must be applied before broadleaf weeds germinate. Gallery should be tank-mixed with another preemergence herbicide such as prodiamine (Barricade), dithiopyr (Dimension), pendimethalin (Pendulum), or oryzalin (Surflan) if annual bluegrass or other grasses are expected.

Atrazine (AAtrex) and simazine (Princep T&O) are the backbone products for postemergence winter weed control for warm-season turfgrasses such as centipedegrass, St. Augustinegrass, zoysiagrass, and bermudagrass. These materials should be used in mid fall (October/November) for optimum control timing. A follow-up application may be needed three weeks later for total control. These herbicides become less effective when applied after January. Also, if these are applied during spring



Chickweed (*Stellaria media* (L.) Cyrillo) is a mat-forming, many-stemmed winter annual.

green-up, temporary set back of turfgrasses should be expected.

For those broadleaf weeds that these herbicides do not effectively control, a single or combination application of 2,4-D, 2,4-DP, MCPP, MCPA or dicamba are needed (Table 2). These herbicides are selective, systemic, foliar-applied herbicides. Several considerations are noted before using one or more of these materials. First, few broadleaf weeds, especially perennials, are controlled with just one of these herbicides. A two or three-way combination is generally necessary for satisfactory results. Control also depends on the maturity of the weed. Younger weeds are easiest and cheapest to control. Applications should ideally be initiated in November to take advantage of these younger, more succulent plants. Waiting until March or April to attempt control requires sequential applications spaced 10 to 14 days apart. This increases labor and herbicide costs and equipment wear and tear. Later applications also may delay turfgrass green-up and require longer for herbicides to work.

Until recently, these herbicide combinations were used as the main control chemicals for broadleaf weeds. New chemistry such as triclopyr and clopyralid (Turflon and Confront) have been intro-

cont. on page 51



TURF-SEED'S PREMIUM PERENNIAL

# RYEGRASS

FOR PERMANENT TURF AND WINTER OVERSEEDING

Variety / Blend

Characteristics

<b>Alliance Brand</b>	Traffic, heat and drought tolerant / A blend of top quality Turf-Seed perennial ryegrasses
<b>BrightStar II (2M3)</b>	A top variety in the 1995 NTEP trials. A unique leafy, decumbent, late maturing cultivar
<b>BrightStar</b>	Tops 1994 NTEP ryegrass trials / Extremely dark color / Dwarf growth habit
<b>Catalina</b>	Low growth profile and reduced clippings / 78% endophyte / Tested as GH-94
<b>Chaparral</b>	Very dense, low-growing, dark green turf. Lowest vertical growth in turf. Tested as 2DLM
<b>Charger</b>	Good low fertility and cool weather performance / Improved disease resistance
<b>Charger II</b>	Dark, dense, dwarf turf ranks high in winter overseeding trials. Experimental code 2QM.
<b>Citation III</b>	Excellent heat and wear tolerance / Improved disease resistance / Rich, dark green color
<b>Manhattan 3</b>	Very dense growth habit / Rich color / Good mowing qualities / Contains endophyte
<b>Navajo</b>	Dark green, disease resistant turf / Excellent for overseeding / Good shade tolerance
<b>Quickstart</b>	Heat tolerant / Establishes quickly / Disease resistant / Excellent for overseeding
<b>Roadrunner (2ET)</b>	Dark blue-green color ideal for winter overseeding and permanent turf / Contains endophyte
<b>Sunrye (246)</b>	Dark green color / Dwarf growth habit / Contains endophyte / Disease resistant

*Turf-Seed, Inc. uses the full recommended rate of Apron® XL™ LS seed treatment. Help your grass seed get a good, healthy start.*

Produced and  
Marketed by



TURF-SEED, INC. / PO Box 250, Hubbard, OR 97032 / 800-247-6910 / 503-651-2130 / FAX 503-651-2351

Circle No. 144 on Reader Inquiry Card



TURF-SEED'S PREMIUM PERENNIAL

# RYEGRASS

FOR PERMANENT TURF AND WINTER OVERSEEDING

*Perennial ryegrasses from Turf-Seed, Inc. represent the latest in research and development technology and express a sound investment in turfgrass agronomics.*

*Our varieties and blends typically produce:*

- Rich, dark green color with enhanced disease resistance
- Quick-to-establish, dense tillered turf with improved mowing qualities
- Excellent turfgrass ratings in NTEP Trials, plus California, Arizona, and North Carolina overseeding trials
- Most contain high levels of endophyte for natural above-ground feeding insect resistance

#### **Establishment**

Turf-Seed perennial ryegrasses establish quickly in bare soil and will germinate in six days with

#### **SEEDING RATES FOR PERMANENT TURF**

Athletic fields, playgrounds	.8-10 lbs./1000 sq. ft.
Golf course fairways, roughs, and cart paths	.8-10 lbs./1000 sq. ft.
Parks, trailer courts, and camp areas	.8-10 lbs./1000 sq. ft.
Low maintenance areas	.3-5 lbs./1000 sq. ft.

#### **WINTER OVERSEEDING RATES**

Greens	.30-40 lbs./1000 sq. ft.
Tees	.20-30 lbs./1000 sq. ft.
Fairways	.500-700 lbs./acre
Roughs	.450-600 lbs./acre
Athletic fields, playgrounds	.10-15 lbs./1000 sq. ft.
Home lawns, parks, civic areas	.8-10 lbs./1000 sq. ft.

favorable conditions. Dark color and a height of 2 to 3" can be expected in two weeks, with the first mowing at about 21 days. Limited use can be expected in 30 days. Our advanced varieties can be mowed at 1/4" and less for Winter overseeding greens and tees in the South.

Our perennial ryegrasses perform exceptionally well as a monostand, and when mixed 1/3, 1/3, 1/3 with Turf-Seed's Kentucky bluegrass and fine fescues for year-round home lawns, parks, and other shady areas subject to wear and cold. Seed this mixture at 5 lbs. per 1,000 sq. ft., or 400 lbs. per acre. Keep seeds moist until seedlings emerge in 21 days.

#### **Fertilization rate**

Due to the naturally dark-green color of Turf-Seed ryegrass varieties and blends, they should require less fertilizer than is customary. Three to 6 lbs. of nitrogen per 1000 sq. ft. per season should be adequate.

#### **Blue Tag Certified Program**

Turf-Seed ryegrass blue tags earn cash for turfgrass research. Hundreds of thousands of dollars have been contributed to university research and scholarship programs as well as turfgrass councils in the purchaser's name. Call Turf-Seed, Inc., or ask your seed dealer for details.

PRODUCED AND  
MARKETED BY

TURF-SEED, INC. • PO Box 250 • Hubbard, OR 97032 • 503-651-2130 • FAX 503-651-2351 • 800-247-6910

© Turf-Seed, inc.

Circle No. 149 on Reader Inquiry Card